



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ





شبكة المعلومات الجامعية



شبكة المعلومات الجامعية

التوثيق الالكتروني والميكرو فيلم

جامعة عين شمس

التوثيق الالكتروني والميكرو فيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأفلام قد اعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15 – 20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of
15 – 25c and relative humidity 20-40 %



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



شبكة المعلومات الجامعية



بالرسالة صفحات
لم ترد بالأصل

**EFFECT OF SOME DIFFERENT UNILATERAL
DISTAL EXTENSION PROSTHODONTIC
TECHNIQUES ON THE ACTIVITY OF
TEMPORALIS AND MASSETER
MASTICATORY MUSCLES.**

Thesis

Submitted to the Faculty of Oral and Dental Medicine Cairo
University in Partial Fulfillment of the Requirement for Master
Degree in Crown and Bridge Fixed Prosthodontics

By

KHALED MAHMOUD KHOLIEF
B.D.S. (Cairo)

Faculty of Oral & Dental Medicine
Cairo University

2000

B71A

SUPERVISORS

Prof. Dr. Fattouh Abd allah Ramadan.

***Professor and Chairman of
Crown and Birdge Department.***

**Faculty of Oral and Dental Medicine
Cairo University**

Dr. Suzanne Mahmoud El-Sayed.

Associate Professor of Crown and Bridge

**Faculty of Oral and Dental Medicine
Cairo University**

٢. د. سوزان محمود

د. فاضل
د. فاضل
د. فاضل

د. فاضل



ACKNOWLEDGEMENT

First of all, I thank God for blessing this work until it has reached its end, as part of his generous help throughout my life.

I wish to acknowledge my sincere appreciation and gratitude to Prof. Dr. *Fattouh Abd allah Ramadan*, Professor and Head of Crown & Bridge Department, Faculty of Oral & Dental Medicine, Cairo University, for his continuous encouragement and valuable support.

I would like to express my deepest thanks and sincere gratitude to Dr. *Suzanne Mohmoud El Sayed*, Associate Professor of Crown & Bridge, Faculty of Oral & Dental Medicine, Cairo University, for planning this work, for her kind supervision, eminent guidance, enthusiastic support and great help. I must admit that I consider myself fortunate to have worked under her supervision.

CONTENTS

	Page.
Introduction	1
Review of Literature	4
Aim of the Study	21
Materials and methods	22
Results	50
Discussion	91
Summary and Conclusion	98
References	100
Arabic Summary	

LIST OF FIGURES

Figures No.	Title	Page
Fig. (1) :	<i>Preparation of the second premolar and the first molar showing the sub-gingival finishing line</i>	27
Fig. (2) :	<i>The finished bridge on the cast.</i>	27
Fig. (3) :	<i>The fixed bridge after cementation in the patient's mouth.</i>	28
Fig. (4) :	<i>The design of the removable partial denture</i>	32
Fig. (5) :	<i>The finished removable partial denture on the cast.</i>	32
Fig. (6) :	<i>The finished removable partial denture in the patient's mouth.</i>	33
Fig. (7):	<i>The ART precision attachment.</i>	37
Fig. (8) :	<i>The milling surveyor</i>	37
Fig. (9) :	<i>The finished cast restoration after application of the porcelain.</i>	38
Fig. (10) :	<i>The finished cast restoration in the patient's mouth.</i>	38
Fig. (11) :	<i>The finished removable part</i>	39
Fig. (12) :	<i>The male part of the attachment in its place in the removable part.</i>	39
Fig. (13) :	<i>The finished partial denture with precision attachment on the cast.</i>	40
Fig. (14) :	<i>The finished partial denture with precision attachment in the patient's mouth.</i>	40
Fig. (15) :	<i>The Cadwel Excel electromyogram.</i>	45
Fig. (16) :	<i>Patient position and electrodes placement.</i>	47
Fig. (17) :	<i>Changes of mean volt values of masseter muscle in the edentulous area in patients treated with fixed bridges (group A)</i>	65
Fig. (18) :	<i>Changes of mean volt values of temporalis muscle in the edentulous area in patients treated with fixed bridges (group A)</i>	66

Figures No.	Title	Page
Fig. (19) :	<i>Changes of mean volt values of masseter muscle in the edentulous area in patients treated with removable partial dentures (group B)</i>	67
Fig. (20) :	<i>Changes of mean volt values of temporalis muscle in the edentulous area in patients treated with removable partial dentures (group B)</i>	68
Fig. (21) :	<i>Changes of mean volt values of masseter muscle in the edentulous area in patients treated with partial dentures with precision attachments (group C)</i>	69
Fig.(22) :	<i>Changes of mean volt values of temporalis muscle in the edentulous area in patients treated with partial dentures with precision attachments (group C)</i>	70
Fig. (23):	<i>Changes in power values of masseter muscle in the edentulous area in patients treated with fixed bridges (group A)</i>	85
Fig. (24):	<i>Changes in power values of temporalis muscle in the edentulous area in patients treated with fixed bridges (group A)</i>	86
Fig. (25) :	<i>Changes in power values of masseter muscle in the edentulous area in patients treated with removable partial dentures (group B)</i>	87
Fig. (26) :	<i>Changes in power values of temporalis muscle in the edentulous area in patients treated with removable partial dentures (group B)</i>	88
Fig. (27) :	<i>Changes in power values of masseter muscle in the edentulous area in patients treated with partial dentures with precision attachments (group C)</i>	89
Fig. (28) :	<i>Changes of power values of temporalis muscle in the edentulous area in patients treated with partial dentures with precision attachments (group C)</i>	90

LIST OF TABLES

Tables No.	Title	Page
Table (1) :	<i>Mean volt values (in milli volt)of masseter and temporalis muscles in patients of group A for maximum clenching in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the fixed prosthesis.</i>	56
Table (2) :	<i>Mean volt values (in milli volt)of masseter and temporalis muscles in patients of group B for maximum clenching in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the removable partial denture.</i>	57
Table (3) :	<i>Mean volt values (in milli volt)of masseter and temporalis muscles in patients of group C for maximum clenching in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the partial denture with precision attachment.</i>	58
Table (4) :	<i>Mean volt values (in milli volt)of masseter and temporalis muscles in patients of group A for soft food (banana) chewing in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the fixed prosthesis.</i>	59
Table (5) :	<i>Mean volt values (in milli volt)of masseter and temporalis muscles in patients of group B for soft food (banana) chewing in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the removable partial denture.</i>	60
Table (6) :	<i>Mean volt values (in milli volt)of masseter and temporalis muscles in patients of group C for soft food (banana) chewing in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the partial denture with precision attachment.</i>	61
Table (7) :	<i>Mean volt values (in millivolt)of masseter and temporalis muscles in patients of group A for hard food (carrot) chewing in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the fixed prosthesis.</i>	62

Tables No.	Title	Page
Table (8) :	<i>Mean volt values (in milli volt) of masseter and temporalis muscles in patients of group B for hard food (carrot) chewing in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the removable partial denture.</i>	63
Table (9) :	<i>Mean volt values (in milli volt) of masseter and temporalis muscles in patients of group C for hard food (carrot) chewing in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the partial denture with precision attachment.</i>	64
Table (10) :	<i>Power values (in micro watt) of masseter and temporalis muscles in patients of group A for maximum chenching in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the fixed prosthesis.</i>	76
Table (11) :	<i>Power values (in micro watt) of masseter and temporalis muscles in patients of group B for maximum clenching in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the removable partial dentures.</i>	77
Table (12) :	<i>Power values (in micro watt) of masseter and temporalis muscles in patients of group C for maximum clenching in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the partial denture with precision attachment.</i>	78
Table (13) :	<i>Power values (in micro watt) of masseter and temporalis muscles in patients of group A for soft food (banana) chewing in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the fixed prosthesis.</i>	79
Table (14) :	<i>Power values (in micro watt) of masseter and temporalis muscles in patients of group B for soft food (banana) chewing in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the removable partial dentures.</i>	80

Tables No.	Title	Page
Table (15) :	<i>Power values (in micro watt) of masseter and temporalis muscles in patients of group C for soft food (banana) chewing in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the partial denture with precision attachment.</i>	81
Table (16) :	<i>Power values (in micro watt) of masseter and temporalis muscles in patients of group A for hard food (carrot) chewing in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the fixed prosthesis.</i>	82
Table (17) :	<i>Power values (in micro watt) of masseter and temporalis muscles in patients of group B for hard food (carrot) chewing in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the removable partial dentures.</i>	83
Table (18) :	<i>Power values (in micro watt) of masseter and temporalis muscles in patients of group C for hard food (carrot)chewing in dentulous and edentulous area before treatment and during a follow-up period of 2 months after receiving the partial denture with precision attachment.</i>	84

INTRODUCTION