







شبكة المعلومـــات الجامعية التوثيق الالكتروني والميكروفيا.



جامعة عين شمس

التوثيق الالكتروني والميكروفيلم



نقسم بللله العظيم أن المادة التي تم توثيقها وتسجيلها على هذه الأفلام قد اعدت دون آية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

40-20 في درجة حرارة من 15-20 منوية ورطوبة نسبية من

To be kept away from dust in dry cool place of 15 – 25c and relative humidity 20-40 %









BIINY

PROGNOSTIC AND PREDICTIVE VALUE OF POSTERIOR CHEST LEADS IN ACUTE INFERIOR MYOCARDIAL INFARCTION

A thesis submitted for partial fulfillment of master degree in Cardiology

BY
Alallah El Said El Beheery
(M.B.B.CH)

Under supervision of

Prof. Dr. Ahmed Abd El-Moneim Mohammed Professor of cardiovascular medicine

Banha university


Prof. Dr. Hesham Mohammed Abo El-Enein

Professor of cardiovascular medicine
Banha university

Dr. El-Sayed Abd El-Khalik Mahmoud

Lecturer of cardiovascular medicine
Banha university

2001



A thesis submitted for partial fulfillment of master degree in Cardiology

BY
Alallah El Said El Beheery
(M.B.B.CH)

Under supervision of

Prof. Dr. Ahmed Abd El-Moneim Mohammed

Professor of cardiovascular medicine
Banha university

Prof. Dr. Hesham Mohammed Abo El-Enein

Professor of cardiovascular medicine
Banha university

Dr. El-Sayed Abd El-Khalik Mahmoud

Lecturer of cardiovascular medicine
Banha university

2001

(ريسم المؤلمة الركي الركيم)

اقر أ ريسم ريكي الركي الأليم الأربسان من على المؤلم المؤلسان ما المربي القلم المؤلسان ما المربي المؤلسان ما المربي المؤلسان ما المربي المؤلسان ما المربي المؤلسان المنازرة ١٠٥٠)

ACKNOWLEDGMENT

First and almost thanks to 'Allah' prayness and greeting upon His Prophet.

I am deeply indebted to Prof. Dr. Ahmed Abd El-Moneim Mohammed professor in cardiology Benha faculty of Medicine for his thanks and faithful support. I can not express may deep thanks and gratitude for his advice and genuine help.

I would like to express may deepest thanks to Prof. Dr. Hesham Mohammed Abo El-Enein assistant professor of cardiology and cardiovascular medicine, Benha faculty of Medicine for his supervision upon this work guidance and endless cooperation Throughout the course of this work..

I do not find words to thank Dr. El-Sayed Abd El-Khalik Mahmoud Lecturer in cardiology Benha faculty of Medicine to whom I feel most indebted for his essential guidance help up on Practical part of this work endless support and continuous encouragement.

Finally I would like to thank my parents and my wife for their personal support and encouragement.

Dedication

To soul of my father,

To my mother,

To my wife and

To my sons

Contents

Introduction and time su	Page
Introduction and aim of the work	1
Review of literature	4
*Pathology& Pathophysiology of AMI	4
*Radiological anatomy of coronary arteries	14
*ECG & myocardial infarction	27
*Correlation of ECG and angiography.	31
* Posterior myocardial infarction	33
*Limitation of E.C.G	33 37
Risk stratification of AMI-	•
Patients and methods.	57
Results.	73
Discussion .	82
	124
Summary and conclusion.	133
References	136
Appendix.	166
Arabic summary.	168

LIST OF TABLE

Table no	Table	page
Table (1)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding age	89
Table (2)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding sex	90
Table (3)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding history of relevant disease	91
Table (4)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding risk factor	92
Table (5)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding clinical data	93
Table (6)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding complication	94
Table (7)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding number of complicated patients	95
Table (8)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding admitted ECG patterns	97
Γable (9)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding CPK	98

Table (10)	Comparison between patients with & without	100
	STE in posterior chest leads (PCL) regarding EF.	100
Table (11)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding MR	102
Table (12)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding wall motion abnormality	103
Table (13)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding posterior & lateral wall motion abnormality versus others.	104
Table (14)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding Inferior &normal wall motion abnormality versus other	105
Table (15)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding affected arteries	107
Table (16)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding percentage of culprit artery from total lesion of each group	109
Table (17)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding circumflex and right coronary artery	110
Table (18)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding number of vessels	111
Table (19)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding site of lesion in coronary artery	113

Table (20)	Validity of STE in PCL in relation to posterior wall motion abnormality in diagnosis of posterior myocardial infarction in echocardiography.	115
Table (21)	Validity of R/S ≥1 in relation to PWMA in diagnosis of posterior myocardial infarction in V1-V2	116
Table (22)	Validity of STD in relation to PWMA in diagnosis of posterior myocardial infarction in precardial leads V1-V3	117

.

.

LIST OF FIGURES

Figure no	Figure	page
Figure (1)	Distribution of the studied patients.	88
Figure (2)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding complication	96
Figure (3)	Comparison between patients with & without STE in posterior chest leads (PCL) CPK	99
Figure (4)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding EF	101
Figure (5)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding wall motion abnormalities.	106
Figure (6)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding affected arteries.	108
Figure (7)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding number of vessels affected	112
Figure (8)	Comparison between patients with & without STE in posterior chest leads (PCL) regarding site of coronary lesion	114
Figure (9)	ECG of AIMI without STE in PCL	118
Figure (10)	. Echocardiography of AIMI patient EF 68%	119
Figure (11)	Angiography of AIMI showing total occlusion of right coronary artery at its middle	120