

IMMUNOHISTOCHEMICAL STUDY OF THE EXPRESSION OF OCT-4 IN BLADDER UROTHELIAL CARCINOMA

Thesis submitted for partial fulfillment of

The M.Sc. degree of Pathology

By

Amal Mohammed Salah El-Din Ibrahim Asar

M.B.B.Ch

Supervised by

Prof. Dr. Samya Mohammed Gabal

Professor of Pathology

Cairo University

Prof. Dr. Noha Abdel Hafez Helmy Mohammed

Professor of Pathology

National Research Centre

Dr. Sara El-Sayed Khalifa

Lecturer of Pathology

Cairo University

2016

TABLE OF CONTENTS

ACKNOWLEDGEMENT	VIII
ABSTRACT	IX
INTRODUCTION	1
AIM OF THE WORK	3
ANATOMY AND HISTOLOGY OF THE URINARY BLADDER	4
EPITHELIAL LESIONS OF THE URINARY BLADDER	7
CANCER STEM CELLS	40
MATERIALS AND METHODS	47
RESULTS	51
DISCUSSION	86
CONCLUSION	96
REFERENCES	97
ARABIC SUMMARY	X

LIST OF TABLES

Table 1	Incidence and mortality of Bladder Cancer in Egypt per sex	7
Table 2	Stage 0a and 0is	16
Table 3	Sage I	16
Table 4	Stage II	17
Table 5	Stage III	18
Table 6	Stage IV	19
Table 7	Histological parameters useful in the diagnosis of flat lesions with atypia	22
Table 8	Clinical outcomes of patients with atypical urothelial proliferations of the urinary bladder based on the 1998 WHO/ISUP classification	24
Table 9	Morphologic patterns of carcinoma in situ	25
Table 10	Histologic features of papillary urothelial lesions.	28
Table 11	Variants of invasive urothelial tumours	33
Table 12	Staining protocol and relevant reagents, temperatures and durations	49
Table 13	Characteristics of the study cases	51
Table 14	Grades of the studied bladder urothelial carcinoma	53
Table 15	Morphology of the studied bladder urothelial carcinoma	54
Table 16	Stages of the studied bladder urothelial carcinoma	55

Table 17	Frequencies of superficial and deep bladder urothelial carcinoma without and with variant morphology	56
Table 18	OCT-4 expression as detected by H- score in studied bladder urothelial carcinoma	58
Table 19	Oct-4 expression in age groups of studied cases	59
Table 20	Oct-4 expression according to sex of the studied cases	60
Table 21	OCT-4 expression according to grade of bladder urothelial carcinoma	61
Table 22	OCT-4 expression according to the different histopathologic types of urothelial carcinoma.	62
Table 23	Level of Oct-4 expression according to the studied morphologies of urothelial tumours.	64
Table 24	Level of Oct-4 expression according to the studied morphologies of urothelial tumours.	65
Table 25	OCT-4 expression according to morphological variants of bladder urothelial carcinoma	66
Table 26	OCT-4 expression according to stage of bladder urothelial carcinoma	67
Table 27	Level of Oct-4 expression in the different stages of studied cases	68
Table 28	OCT-4 expression in superficial and deep bladder urothelial carcinoma without variant morphology.	69
Table 29	H- score level among positive superficial and deep urothelial tumour cases without variant morphology	70
Table 30	OCT-4 expression according to Belharzial association	71

LIST OF GRAPHS

Graph 1	Age distribution of studied cases	52
Graph 2	Percentage distribution of studied cases by gender	52
Graph 3	Distribution of studied specimens according to grade	53
Graph 4	Frequency of studied morphologies of bladder urothelial carcinoma	54
Graph 5	Stage frequency of studied cases	55
Graph 6	Frequencies of superficial and deep urothelial carcinoma cases	57
Graph 7	Oct-4 expression in studied cases	58
Graph 8	Oct-4 expression in age groups of studied cases	59
Graph 9	Oct-4 expression in sex groups of studied cases	60
Graph 10	Oct-4 expression as per grade of the studied cases	61
Graph 11	Oct-4 expression in different studied histopathological types	63
Graph 12	Level of Oct-4 expression in studied histopathological types	64
Graph 13	Oct-4 expression in studied stages of urothelial carcinoma	67
Graph 14	Level of Oct-4 expression in the studied stages	69
Graph 15	Frequency of belharzial infection in studied cases	71

LIST OF FIGURES

Figure 1	Cancer stem cell model proliferation, regenerating more stem cells and differentiating into the heterogenous tumour population.	41
Figure 2	Invasive urothelial carcinoma infiltrating muscle surrounded by desmoplastic stroma with lymphoplasmacytic infiltrate	72
Figure 3	Infiltrating cohesive nests of urothelial carcinoma within the muscularis propria, stage T2	72
Figure 4	Urothelial carcinoma infiltrating the muscularis propria with multiple vascular emboli	73
Figure 5	Sheets of infiltrating urothelial carcinoma dissecting through muscle bundles of muscularis propria, stage T2	73
Figure 6	Urothelial carcinoma with evidence of perivascular fat extension, stage T3	74
Figure 7	Nests of urothelia carcinoma infiltrating the prostate, stage T4, showing pleomorphism and scattered prominent nucleoli	74
Figure 8	High magnification of the vascular emboli	75
Figure 9	Papillary urothelial carcinoma; high grade, stage Ta	75
Figure 10	Papillary urothelial carcinoma; low grade, stage Ta	76
Figure 11	Urothelial carcinoma in-situ	76
Figure 12	Sarcomatoid variant of urothelial carcinoma	77
Figure 13	Squamoid variant of urothelial carcinoma	78
Figure 14	Micropapillary variant of urothelial carcinoma	78

Figure 15	Infiltrating urothelial carcinoma with glandular differentiation	79
Figure 16	Infiltrating undifferentiated urothelial carcinoma	79
Figure 17	Seminoma control with positive nuclear staining for Oct-4, intensity score 3	80
Figure 18	Papillary urothelial carcinoma showing positive nuclear staining for Oct-4, where there are clusters of positive cells (inset), intensity score 3	80
Figure 19	Infiltrating urothelial carcinoma showing nuclear positivity for Oct-4, intensity score 3	81
Figure 20	Infiltrating urothelial carcinoma showing nuclear positivity for Oct-4 in a single malignant cell, intensity score 2	81
Figure 21	Infiltrating urothelial carcinoma, showing positive nuclear staining for Oct-4 in two cells, intensity score 2 and score 1	82
Figure 22	Infiltrating urothelial carcinoma showing positive nuclear staining for Oct-4, intensity score 2 and score 1	82
Figure 23	Infiltrating urothelial carcinoma showing positive nuclear staining for Oct-4, intensity score 1	83
Figure 24	Infiltrating urothelial carcinoma showing positive nuclear staining for Oct-4, intensity score 1	83
Figure 25	Infiltrating urothelial carcinoma negative for Oct-4 staining, score 0	84
Figure 26	Infiltrating urothelial carcinoma negative for Oct-4 staining, score 0	84
Figure 27	Infiltrating urothelial carcinoma showing negative tumour cells for Oct-4, score 0, with concomitant Oct-4 positivity in the desmoplastic reaction, intensity score 3	85

LIST OF ABBREVIATIONS

AJCC: American Joint Committee on Cancer

CIS: Carcinoma In-Situ

CSC: Cancer Stem Cell

DAB: Di-Amino Benzidine

H&E: Haematoxylin and Eosin

ISUP: International Society of Urological Pathology

TNM: Tumour, Nodal involvement, Metastasis

UBC: Urinary Bladder Cancer

WHO: World Health Organisation

Acknowledgement

As my first practical research, am greatly indebted to all the professors and colleagues who helped me through the entire process. It was really challenging and I wouldn't have endured it if not for the support I received.

Am thankful to Professor Dr/ Samya Gabal; professor of Pathology, Cairo University for her immense facilitation and her keen concern; she always edited my work promptly and was keen that all steps were complete as soon as possible; without compromising the scientific approach and scientific quality of the material.

I need to pay a special gratitude to Professor Dr/ Noha Abdel-Hafez Helmy; Research professor of Pathology, National Research Centre, for suggesting the point of research while considering my field of interest. Her care and support never ceased throughout the steps of the research; starting by complementing my missing or rare cases from her work flow, following up on the immunostaining, and finally her concern to teach me how to scientifically think and how to record my observations and formulate research questions or hypotheses.

Many thanks go to Professor Dr/ Fatma Adly Morsy; Research Professor of Histochemistry, National Research Centre, for her continuous encouragement and follow up.

Thanks are due to Dr/ Sara El Sayed Khalifa; Lecturer of Pathology, Cairo University, who critically edited the work and reviewed it meticulously time after time, without her efforts and observations, this work wouldn't have been produced with the current quality.

I am greatly indebted to Professor Dr/ Elia Anis Isaak, professor of Pathology, Cairo University, who contributed to this work by his guidance, his scientific suggestions and by facilitating the practical steps of the work; both case collection and immunostaining; I am really thankful to have worked under the mentor of such a great professor.

I am also thankful to Professor Dr/Samir Atlam, Professor of Public Health, Tanta University, who endured with me a lot while writing the results chapter; his patience in interpretation of the raw data and conducting the statistical tests is greatly appreciated.

Finally I would like to thank my family who endured a lot of stresses throughout this process, both mentally and physically; and never ceased to support me throughout the journey.

ABSTRACT

Bladder carcinoma is a significant problem in Egypt; being the 10th in the worldwide rank of incidence for both sexes. The disease is characterized by multifocality, multiple morphologies coexisting together and multiple recurrences after excision. A cancer stem cell model was proposed; that there is a subpopulation of tumour cells capable of resisting conventional therapies and surviving treatment to facilitate recurrence and metastasis. Oct-4 is a pluripotency marker of stem cells; which has been found to be associated with worse prognosis in multiple somatic tumours such as breast, brain, skin and thyroid.

We tested 84 urothelial tumour specimens including all grades, stages and some variants of urothelial carcinoma for the expression of Oct-4. Two sections were prepared per case for histologic evaluation and immunohistochemical staining by Oct-4 monoclonal antibody. The immunostaining was evaluated semiquantitatively through obtaining an H-score for each case.

All cases took up the Oct-4 stain except for low grade tumours and CIS cases. The highest percentage of positive cases was seen in variants of urothelial carcinoma (19.1%). Oct-4 expression was significantly associated with grade ($p=0.03$), histopathologic type of urothelial tumour ($p<0.001$), stage ($p<0.001$), and tumour progression ($p=0.001$). Conversely; the level of Oct-4 expression was not found to be associated with any of the studied parameters.

Our study proves that Oct-4 is expressed in bladder urothelial carcinoma, and its presence regardless of the level of expression seems to play a role in guiding tumour grade, stage, progression and histopathological type; which calls for considering Oct-4 as a novel prognostic marker that could be implemented in target therapies for urothelial carcinoma.

INTRODUCTION

Recent work in cancer research shows mounting evidence supporting the idea that tumors, similar to normal adult tissues, arise from a specific stem-like cell population, the cancer stem cells (CSCs), which are considered as the real driving force behind tumor growth, the ability to metastasize, as well as resistance to conventional antitumor therapy (**Dimov et al., 2010**).

Stem cells are unspecialized cells that undergo unlimited self-renewal and multilineage differentiation to become specialized cells. They usually constitute a small percentage of the total cell population in any tissue and are usually indolent; dividing very slowly (**Stricker and Kumar, 2010**). It is generally accepted that CSCs might arise from transformed stem cells or progenitor cells that have regained self-renewal activity; due to the autonomous cell-cycle control, alterations of cellular stress-control mechanisms, and interference of signaling pathways; as a result of accumulated genetic and epigenetic changes (**Dimov et al., 2010**).

Extensive research in the field of cancer stem cells; in trying to isolate them and characterize their surface markers, showed that both normal stem cells and CSC share common markers and biological behavior (**Hatina and Schultz, 2012**).

The stem cells of individual urothelial carcinomas appear to differ considerably and may contribute to the heterogeneity of this disease. The presence and quantity of urothelial carcinoma stem cells in each case may thus carry important clinical information that might allow a rationale stratification of urothelial cancer patients and probably pave the way for CSC targeted therapy. Many stem cell markers of normal urothelium have been the target of research in recent years, mainly CD44 and basal-type cytokeratins such as CK5 and CK 17, being usually expressed in the proposed normal urothelial stem cells (**Hatina and Schultz, 2012**).

However, in cancer biology, the relationship between embryogenesis and oncogenesis has long been a prevailing theme. Broadly speaking, cancer cells are similar to very early embryonic cells, which are immortal, undifferentiated, and invasive. It is important to study genes associated with embryogenesis and

tumorigenesis. The *POU* (Pit-Oct-Unc) family transcription factor member OCT-4 (POU5F1) acts as a key regulator of pluripotency in early stages of mammalian development. It has been shown that a critical amount of OCT-4 is required to sustain self-renewal of embryonic stem (ES) cells, and any up-regulation or down-regulation induces divergent cell fates (**Cheng et al., 2008**).

AIM OF WORK

1. To understand whether OCT-4 is expressed in bladder urothelial carcinoma and its variants.
2. To predict the likely association of OCT-4 with the grade and stage of the tumour.
3. To delineate whether OCT-4 is a significant prognostic marker calling for special aggressive treatment postoperatively for patients with limited disease and/or target therapy.

ANATOMY OF THE URINARY BLADDER

The urinary bladder is an intrapelvic, extraperitoneal muscular structure designed to act as a reservoir of urine that allows controlled voiding whenever necessary. The adult empty bladder has the shape of an inverted pyramid and is enveloped by the vesical fascia. The superior surface faces superiorly and is covered by the pelvic peritoneum. The posterior surface, also known as the base of the bladder, faces posteriorly and inferiorly. Relations of this surface are of particular importance since it is this wall that harbours most of the bladder neoplasms. A growing neoplasm in this wall will directly infiltrate these posterior relations; which are the rectum separated by the seminal vesicles and the ampulla of the vasa deferentia in males and the uterine cervix and the proximal portions of the vagina in females. The remainder of the pyramid is made up of two inferolateral surfaces, which together with the base of the bladder join to form the bladder neck. The bladder bed (structures on which the bladder neck rests) is formed posteriorly by the rectum in males and the vagina in females. Anteriorly and laterally it is formed by the internal obturator and the levator ani muscles as well as the pelvic bones (**Moore, 2013**).

The “apex” or “dome” (the most anterior part of the bladder) is located at the point of contact of the superior surface and the two inferolateral surfaces. It marks the point of insertion of the median umbilical ligament (area where urachal carcinomas occur) (**Reuter, 1997**). The ureters enter the bladder through a tunnel traversing the muscosa and mucosa. Each ureteric orifice is a small slit-like opening, and they are located about 2-3 cm apart and 2 cm above the internal urethral meatus in the adult bladder. The triangular region bound by these three openings in the posterior wall of bladder is called the trigone (**Engel, 1997**).

HISTOLOGY OF THE URINARY BLADDER

The urinary bladder consists of a meshwork of thick muscle bundles that surround a highly vascular lamina propria lined by stratified epithelium (Reuter, 1992). The mucosa; or urothelium, is 5-7 cell layers thick in contracted bladder, 2-3 cells thick in distended bladder and is composed of 3 cell types: basal, intermediate, and superficial (umbrella) cells. The umbrella cells comprise a single layer of cells, and are relatively large and overlay multiple underlying cells in the intermediate cell layer. In the past their presence in histological preparations used to indicate a normal maturation process and thus lead to the exclusion of possible malignancy in a suspicious lesion, however, it is now known that they could be observed on the surface of low grade papillary urothelial neoplasms and often in carcinoma in situ (CIS). They appear elliptical with abundant eosinophilic cytoplasm and often binucleation or prominent nucleoli; and are p63 negative and stain positive for CK 18 and CK 20. Moreover, they contain trilaminar (asymmetric) unit membrane composed of two dense layers of unequal thickness and a central lucent layer, and apical plaques containing uroplakins (**Castello-Martin et al., 2010**). Intermediate urothelial cells are cuboidal to low columnar with well defined borders and amphophilic cytoplasm rich in glycogen; nuclei are regularly arranged, ovoid with long axis at right angles to surface; chromatin is finely granular and has small nucleoli usually with no mitotic figures. Basal urothelial cells are more cylindrical and can be flat when bladder wall is stretched; some have longitudinal nuclear grooves. Both cell layers stain positive for p63, CK 5, CK 10 and CK 14 (**Koss, 1985**);(**Castello-Martin et al., 2010**).

Previously the bladder was thought to lack a muscularis mucosa and so the layer of fibrovascular connective tissue beneath the mucosa was termed lamina propria. However it is now well established that this layer contains wisps of incompletely differentiated smooth muscle fibres, that could occasionally be hyperplastic thus resembling the well developed muscularis propria. The importance of this observation is related to the staging of bladder carcinoma, and the significance attributed to tumour invasion of the muscle; where invasion of the muscularis propria carries a clinical significance far greater than invasion confined to muscularis mucosa, which calls for caution when interpreting biopsy specimens. A useful clue is that; in contrast to the well defined, rounded fascicles of the