

# 





ثبكة المعلومات الجامعية





## جامعة عين شمس

التوثيق الالكتروني والميكروفيلم



نقسم بللله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأفلام قد اعدت دون آية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15-20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of 15-25c and relative humidity 20-40 %



ثبكة المعلومات الجامعية







## STUDY OF RISK FACTORS OF OTITIS MEDIA AMONG CHILDREN

Thesis

SUBMITTED FOR PARTIAL FULFILMENT OF M.D. DEGREE

IN CMMUNITY MEDICINE

By Lamia Ahmed Basiony M.B., B.Ch., M.Sc.

Supervised By

Prof. Dr. Taghreed Farahat

11

Professor and Head of Community, Family and Occupational Medicine Department Faculty of Medicine, Menousiya University

Prof. Dr. Gaafer Abdel-Rasoul

Professor of Occupational Medicine
Faculty of Medicine, Menoufiya University

Prof. Dr. Laila Shehata Dorgham

Professor of Public Health and Vice Dean for Environmental Affairs Faculty of Medicine, Menoufiya University

Prof. Dr. Mohamed El-Sharnouby

Professor of ENT and Dean of Faculty of Medicine Menoufiya University


Dr. Mabrouk Ghoneim

Lecturer of Microbiology
Faculty of Medicine, Menouriya University

B 1-019

FACULTY OF MEDICINE MENOUFIYA UNIVERSITY

1999



«إن السمع والبصر والفؤاد كل أولئك كان عنه مسئولاً» سرة الإسراء (آية ٣٦)

«فلهو الذي أنشاكم وجعل لكم السمع والأبصار والأفئدة فليلً ما نشكرور.»

سورة الملك (آية ٢٣)

صدق الله العظيم

To my hsband To my parents

#### Acknowledgement

First and foremost thanks to God for his help and powerful support to full this work.

I wish to express my sincere appreciation and deep gratitude to my professor and supervisor **Prof. Dr. Taghreed Farahat,** Head of Community, Family and Occupational Medicine Department, Faculty of Medicine, Menoufiya University for her generous help, kind moral support, enthusiastic encouragement, precious comments and suggestions.

My sincere thanks, deep appreciations to **Prof. Dr. Laila Sh. Dorgham**, Professor of Public Health Affairs, Faculty of Medicine, Menoufiya University for her continuous support, fruitful supervision, great cooperation and continuous guidance.

To Prof. Dr. Gaafar Abdel-Rasoul, Professor of Occupational Medicine, Faculty of Medicine, Menoufiya University, I owe more than can be expressed for his sincere help, wide guidance, close supervvision, and for his great effort and particularly in the statistical analysis of this work and time he freely gaive to finish this work.

My profound and deep thanks to **Prof. Dr. Mohamed El-Sharnouby**, Professor of ENT, and Dean of Faculty of Medicine, Menoufiya University for his continuous encouragement, generous help, valuable suggestion and precious advice.

I am deeply indebted to **Dr. Mabrouk Ghoneim** Lecturer of Microbiology, Faculty of Medicine, Menoufiya University for his fidelity, sustained support, unlimited help, generous advice and precious ideas.

I wish to express my sincere appreciation and deep thanks to **Prof. Dr. Mohamed Kamel El-Sobky,** Professor of Occupational Medicine, Faculty of Medicine, Menoufiya University for suggesting the subject of this work and his a lot of illuminating discussions he gave throughout this work.

My sincere prays to God to treat the spirit of **Dr. Yousry**Moustafa with patience and kindness for his help,
encouragement, and enthusiastic support at the beginning of this
work.

I am deeply indebted to my colleagues in ENT, Audiology and Community Departments for their help and sympathy which will never be forgotten.

Lastly, special warm heartly acknowledgement to my husband and parents and my friends, without their love, tolerance, cooperation and encouragement, this work would never been possible.

#### LIST OF ABBREVIATIONS

OM Otitis media

ME Middle ear

MEE, MEF Middle ear effusion, middle ear fluid.

OME Otitis media with effusion

ENT Ear, nose and throat

OPC Outpatient clinic

ASM Acute suppurative otitis media

SOM Secretory otitis media

CSOM Chronic suppurative otitis media

HL Hearing loss

ET Eustachian tube

SES Socioeconomic standard

RSF Respiratory syncytial virus

URTI Urinary tract infection.

PMEE Persistent middle ear effusion

ETD Eustachian tube dysfunction.

NP Nasopharyngeal

DC Day care

CHL Conductive hearing loss

SNHL Sensorineural hearing loss

#### **CONTENTS**

	PAGE
• INTRODUCTION	1
• REVIEW OF LITERATURE	3
- Anatomy of the Ear	3
- Definition, Terminology and Classification	5
- Risk Factors in Otitis Media	15
- Aetiology and Pathogenesis of Otitis Media	32
- Microbial Aspects of Otitis Media	41
- Diagnosis of Otitis Media	56
- Complications and Sequelae of Otitis Media	68
- Prevention of Otitis Media	73
• AIMS OF THE WORK	77
• HYPOTHESES	78
• SUBJECTS AND METHODS	79
• LIMITATIONS OF THE STUDY	97
• RESULTS	98
• DISCUSSION	133
• SUMMARY AND CONCLUSIONS	166
• RECOMMENDATIONS	170
• REFERENCES	172
• APPENDIX	222
• PROTOCOL	

• ARABIC SUMMARY

#### LIST OF TABLES

	,	'age No
(1):	Number and percent distribution of ear diseases among children (aged from birth to 12 years) recorded in ENT, OPC during the academic year 1995-1996.	99
(2):	Number and percent distribution of patients and controls according to sociodemographic characteristics.	103
(3):	Number and percent distribution of patients and controls according to family history of OM, father's smoking and smoking index.	104
(4):	Number and percent distribution of patients and controls according to past history of allergic rhinitis, chronic tonsillitis, adenoid hypertrophy and chronic sinusitis.	105
(5):	Number and percent distribution of patients and controls according to past history of upper and lower respiratory tract infections.	106
(6):	Number and percent distribution of patients and controls according to some health behaviors.	107
(7):	Number and percent distribution of patients and controls according to prematurity, history of antibiotic misuse and previous operations (tonsillectomy and adenoidectomy).	108
(8):	Multiple regression results (full model) of occurrence of OM on some studied variables.	110
(9):	Stepwise regression results for factors significantly affecting the occurrence of OM.	111

(10):	Number and percent distribution of OM patients according to laterality, affected side, age, sex, season.	114
(11):	Relation between age at first attack and some studied factors.	117-118
(12):	Relation between recurrent attacks and some studied factors.	120-121
(13):	Number and percent distribution of children by ears according to hearing impairment in relation to type of OM.	124
(14):	Number and percent distribution of culture results in acute and chronic OM.	128
(15):	Number and percent distribution of bacterial isolates in acute and chronic OM.	129
(16):	Number and percent distribution of isolated organisms according to age of the children.	130
(17):	Antimicrobial sensitiity of isolated Gram-positive organisms.	131
(18):	Antimicrobial sensitivity of isolated Gram-negative organisms.	132

### LIST OF FIGURES

		Page No
1)	The difference in the angle of the eustachian tube between infants and adults.	1 4
2)	Etiology and pathogenesis of otitis media.	32
3)	Eustachain tube dysfunction	33
4)	Amploid 720 tympanometer.	85
5)	Five typical tympanogram types.	85
6)	Interacoustic audiometer AD17	87
7).	Studied risk factors according to odds ratio	109
8)	Role and share of factors significantly affecting occurrence of otitis media.	e 112
9)	Different types of otitis media patients according to laterality, age, sex and season	115
10)	Relation between age at first attack and recurrence rate (≥ 3/year)	e 122
11)	Hearing impairment in different types of otitis media.	125