### The Change In Mineral Content Of Teeth After Exposure To Different Types Of Bee Honey

### **Thesis**

Submitted to the Faculty of Dentistry, Ain Shams University in partial fulfillment of the requirements for the master's degree in pediatric dentistry.

Presented by

Marwa Abdel Kader Ahmed Taha

B.D.S, 2004

Faculty of Dentistry, Ain Shams University 2012

### Prof. Dr. Amr Mahmoud Abdel Aziz

Professor and Head of Department of Pediatric
Dentistry and Dental Public Health,
Faculty of Dentistry
Ain Shams University

### Dr. Hisham Imam Mahmoud

Assistant Professor, Laser Applications in Metrology, Photochemistry and Agriculture Department
National Institute of Laser Enhanced Science
(NILES)
Cairo University

# التغيير في المحتوى المعدني للأسنان بعد التعرض لأنواع مختلفة من عسل النحل

رسالة مقدمة لكلية طب اسنان – جامعة عين شمس

توطئه للحصول على درجة الماجستير في طب اسنان الأطفال

مقدمة من

الطبيب/ مروة عبد القادر أحمد طه بكالوريوس طب و جراحة الفم و الاسنان-جامعة عين شمس

۲ . . ٤

كلية طب الأسنان جامعة عين شمس

# <u>المشرفون</u>

اد. عمرو محمود عبد العزيز أستاذ و رئيس قسم طب أسنان الأطفال و الصحة العامة كلية طب الفم و الأسنان جامعة عين شمس

أ.د.م./ هشام إمام محمود أستاذ مساعد بقسم تطبيقات الليزر في القياسات و الكيمياء الضوئية و الزراعة المعهد القومي لعلوم الليزر جامعة القاهرة

#### **Summary**

This study was conducted to assess the change in the mineral content of enamel after the consumption of different types of bee honey that was collected directly from bee farms as well as commercially available types.

Clover and Citrus honey were gathered from 2 different bee farms and from a local store (commercially available: Imtenan®). All 4 types of honey were analyzed regarding their sugar content (by FTIR) and their pH detected (by pH-meter).

Each type of honey was then combined to a Ration (consisting of Poultry feed and Barley grains). The honey /Ration ratio was 1:1, as well as the Barley grains/Poultry Feed ratio was also 1:1

30 Sprague-Dawley rats, Helwan were used to conduct the experiment. The rats were just weaned (3 weeks old), weighing from 30-35 grams. They were grouped into 5 groups (6 rats per group), 4 of which were according to the type of honey introduced and the 5<sup>th</sup> was a control group. The food and water were supplied ad libitum for 6 weeks.

The rats were then sacrificed, the mandibles extracted then defleshed and hemisectioned. After that each mandible was sagittally sectioned. Deionized water was used as a transporting medium for all of the specimens to prevent them from dehydration and for standardization between them.

The specimens were then investigated by scanning electron microscope in conjunction with energy dispersive x-ray analysis system (SEM-EDX) for elemental analysis. In most of the specimens, the 1<sup>st</sup> fissure of the 2<sup>nd</sup> molar was chosen for mineral content analysis.

# بسم الله الرحمن الرحيم

# 

إلا ما عُلمتنا إنك أنت

العليم الحكيم

صدق الله العظيم سورة البقرة "آية 32"

#### This work is dedicated to....

My beloved husband and beautiful children who went through a lot to help me overcome every hardship.

My parents, sister and brother, who have been a constant source of support; emotional, moral and in every aspect of life.

Dr. Osama I. El Shahawy, Assistant Professor,
Pediatric and Dental Public Health Department,
Cairo University, for teaching me the meaning of
"Pediatric dentistry".

And last but definitely not least, my professors, colleagues and friends at the Pediatric Dentistry Department, Ain Shams University. Thank you for your constant support and help whenever needed.

### **Acknowledgement**

Praise be to **ALLAH**, with whose Grace good deeds are accomplished. **ALLAH**, whatever I or any of your creatures are blessed with is from You alone, as there is no associate with You. So all praises and thanks are due to You.

I would like to thank **Dr. Amr Abdel Aziz,** Professor and Head of the Pediatric Dentistry and Dental Public Health Department, Faculty of Dentistry Ain Shams University, for his advices, most valuable remarks, meticulous revision, and expert embellishing of this manuscript.

I would like also to thank **Dr. Hisham Imam** Assistant Professor, Laser Applications in Metrology, Photochemistry and Agriculture Department National Institute of Laser Enhanced Science (NILES), Cairo University.

My deepest gratitude to **Eng. Mohammed Zain**, National Institute of Laser Enhanced Science (NILES), Cairo University, for all his efforts and help with the measurements.

With heartfelt gratitude, I would like to sincerely thank **Dr. Monazah Gamal El Din Khafagi,** Assistant Professor, Spectroscopy Department, Physics Division, National Research Center, to whom I am indebted for her support and help with all the remarks, suggestions, measurements and results, whom without, this work would never have been accomplished.

## **List of Contents**

List of Tables	i
List of Figures	iii
List of Abbreviations	vii
Introduction	1
Review of Literature	4
Aim of the Study	31
Materials and Methods	32
Results	60
Discussion	71
Summary	<b>79</b>
Conclusions	81
Recommendations	82
References	83
Arabic Summary	

## **List of Tables**

Table No.	Table Title	<b>Page</b>
1.	Grouping of different types of bee honey	37
2.	Rats' grouping and distribution.	42
3.	Sugar analysis of Group A (Clover honey)	61
4.	Sugar analysis of Group B (Citrus honey)	61
5.	Sugar analysis of Group C (Clover Imtenan®)	61
6.	Sugar analysis of Group D (Citrus Imtenan®)	61
7.	pH of the honey groups	64
8(a).	Comparison between control and test groups regards calcium weight percent (One-Way ANOVA).	65
8(b).	Comparison between control and test groups regards calcium weight percent. One-Way ANOVA Post Hoc Tests, *=significant ≤ 0.05 and **= significant ≤0.01	66
9(a).	Comparison between control and test groups regards phosphorus weight percent (One-Way ANOVA).	67
9(b).	Comparison between control and test groups regards phosphorus weight percent. One-Way ANOVA Post Hoc Tests, *=significant ≤ 0.05 and **= significant ≤ 0.01	68

10(a).	Comparison between control and test groups regards fluoride weight percent One-Way ANOVA	69
10(b).	Comparison between control and test groups regards fluoride weight percent. One-Way ANOVA Post Hoc Tests, *=significant ≤ 0.05 and **= significant ≤0.01	70

## **List of figures**

Figure	<u>Title</u>	<b>Page</b>
<u>No.</u> 1.	Factors affecting mineral content of teeth	19
2(a).	Pilot, Occlusal view	33
<b>2(b).</b>	Pilot, Sagittal view	33
3	SEM-pilot	34
<b>4(a).</b>	Citrus fruits (Oranges) and their flowers	35
<b>4(b).</b>	Bee collecting the nectar from a Citrus flower	36
<b>4(c).</b>	Bees storing honey in a Honeycomb	36
5.	Grouping of different types of bee honey	37
6.	FTIR 6100 Jasco	39
7.	pH-meter HI 211	39
8.	Poultry Feed	41
9.	Barley Grains	41
10.	Sprague-Dawley rats	43

Rist of Figures
-----------------

11.	Deionized Water	44
12(a).	Control, Occlusal view	45
12(b).	Control, Sagittal view	45
13(a).	Group A, Occlusal view	46
13(b).	Group A, Sagittal view	46
14(a).	Group B, Occlusal view	47
14(b).	Group B, Sagittal view	47
15(a).	Group C, Occlusal view	48
15(b).	Group C, Sagittal view	48
16(a).	Group D, Occlusal view	49
16(b).	Group D, Sagittal view	49
17.	S150A Sputter Coater	51
18.	Gold coating procedure	51
19.	Gold coated Sagittaly-sectioned mandibles	52

20.	Samples on holder before entering SEM-EDX system	52
21.	SEM-EDX system	53
22.	EDX result on the computer screen	54
23.	SEM-Control	55
24.	SEM-Group A	56
25.	SEM-Group B	56
26.	SEM-Group C	57
27.	SEM-Group D	57
28.	FTIR absorbance graph of Honey group A (Clover honey)	62
29.	FTIR absorbance graph of honey group B (Citrus honey)	62
30.	FTIR absorbance curve of honey group C (Clover Imtenan®)	62
31.	FTIR absorbance curve of honey group D (Citrus Imtenan®)	62
32.	FTIR absorbance curves of the 4 spectra of the 4 honey groups.	63
33.	Comparison between control and test groups regards calcium weight percent	65

34.	Comparison between control and test groups regards phosphorus weight percent	67
35.	Comparison between control and test groups regards fluoride weight percent	69