

# **Functional Properties of Selected Ingredients of Some Food Stuff.**

**By**

**Safaa Moustafa Abd El Fatah Faid**

B.Sc. Fac. of Specific Education (1995)

M.Sc. (2002); Food Sci. and Nutrition.

Faculty of Specific Education.

Ain Shams University

A thesis submitted in partial fulfillment

of

The requirements for the degree of

**DOCTOR OF PHILOSOPHY**

in

Home Economics

**( Food Science and Nutrition )**

Department of Home Economics

Faculty of Specific Education

Ain Shams University

**2005**

Approval Sheet

## **Functional Properties of Selected Ingredients of Some Food Stuffs.**

By

**Safaa Moustapha Abd El Fatah Faïd**

B.Sc.(Home Economics Dept.) Faculty of Specific Education ,  
Ain shams University ,(1995)

M.Sc.(Food Science and Nutrition ) Faculty of Specific  
Education,Ain shams University,(2002)

This thesis for Ph.D. degree has been approved by :

**Prof. Dr. M Amin. Abd-Allah**

Prof. of Food Sci; Food Science Dept. Faculty of Agric. Ain Shams  
University. (Former Dean of the Faculty of Specific Education  
(1994- 1997).

**Prof. Dr. Farouk M. El Tellawy**

Prof. of Food Science and Technology. Chairman of the Home  
Economics Dept., Faculty of Specific Education,Ain Shams University.

**Prof. Dr. Sanaa M. El Bendary**

Prof. of Nutrition and Vice Dean of the Faculty of Specific Education  
Ain Shams University.

**Prof. Dr. Mohamed Youssef**

Prof..of Food Science.Food Science and Technology Dept. Fac. of  
Agric., El- Shatby.Univ. of Alexandria.

**Prof. Dr. Yehia Heikal**

Prof. of Food Engineering Food Sci. Dept. Fac. of Agric Ain Shams  
University.

**Date of Examination : 3 / 8 / 2005**

# **Functional Properties of Selected Ingredients of Some Food Stuffs.**

By

**Safaa Moustapha Abd El Fatah Faïd**

Assistant Lecturer; Home Economic Dept.,  
Fac. of Specific Education-  
Ain Shams University

## **Under the Supervision**

**Prof. Dr. M Amin. Abd-Allah**

Prof. of Food Sci; Food Sci. Dept., Faculty of Agric. Ain Shams  
University. (Former Dean of the Faculty of Specific Education  
(1994-1997)).

**Prof. Dr. Farouk M. El Tellawy**

Prof. of Food Science and Technology .Chairman of the Home  
Economics Dept., Faculty of Specific Education, Ain Shams  
University

**Prof. Dr. Sanaa M. El Bendary**

Prof. of Nutrition and Vice Dean of the Faculty of Specific  
Education Ain Shams University.

**Dr. Magda Ali Rakha**

Under Secretary of State, Central Health Labs for Laboratory  
Services, Ministry of Health and Populations

2005

## ACNOWLEDGEMENT

I would like to express my deepest gratitude and appreciations to **Prof. Dr . M. Amin Abd – Allah**, Professor of Food Science, Food Sci. Dept., Faculty of Agriculture. Ain Shams University and the Former Dean of the Faculty of Specific Education (1994 – 1997), for offering valuable suggestion and advising in planning through the supervising of the thesis.

I would like to offer my especiall thanks to **Prof. Dr. Farouk El-Tellawy**, Professor of Food Science and Technology and Chairman of the Home Economics Dep., Fac . of Specific Education, Ain Shams University for supervising , following up the work , helps and support during the preparation and finishing of this study .

Words can never express my heartily thanks to **Prof. Dr. Sanaa El – Bendary**, Prof. of Nutrition , and Vice Dean ; Faculty of Specific Education; Ain Shams University, for supervising, following up , and completion of the work.

I wish also to offer my deeply grateful to **Dr. Magda Ali Rakha** of Under Secretary of State, Central Heath Labs for Laboratory Services, Ministry of Health and Populations for supervising and help offered through analyzing samples that giving me the power to complete this work .

Thanks are given to the Dept of Home economics for giving me the chance to achieve such Ph.D. program.

My deepest thanks are also extended to the private companies for the facilities given through the work:

- National Company for Producing Corn products .10<sup>th</sup> of Ramadan City.
- Cadbury Egypt Group for Food Industries Company.10<sup>th</sup> of Ramadan City .
- Abo El- Dahab Milling Company, 6<sup>th</sup> of October City.

# CONTENTS

|  | <b>Page</b> |
|--|-------------|
| List of Tables   | xiii        |
| List of Figures  | xix         |
| List of Abbreviations  | xxii        |
| <b>Introduction</b>  | <b>1</b>    |
| <b>Review and literature</b>   | <b>7</b>    |
| <b>Part I: Major and minor chemical constituents of the investigated samples .</b>                       | <b>7</b>    |
| <b>1- Major constituents</b>   | <b>7</b>    |
| I:1- Moisture  | 7           |
| 1:2- Crude and dietary fibre.  | 8           |
| 1:3- Protein content   | 8           |
| 1:4- Carbohydrates and ascorbic acid (Vitamin C ).   | 10          |
| <b>2- Minor constituents</b>   | <b>11</b>   |
| 2:1- Minerals and heavy metals   | 11          |
| 2:2- Amino acids   | 13          |
| 2:3- Fatty acids   | 14          |
| <b>Part II: Physicochemical aspects of ingredients of the tested food stuffs.</b>                        | <b>17</b>   |
| 2:1- Water absorption and sorption isotherm.   | 17          |
| 2:2- Watability and water activity   | 18          |
| 2:3- Bulk density , Foaming capacity and emulsion .  | 19          |
| 2:4- Rehydration , modulus of fineness and viscosity properties.   | 23          |
| <b>Part III : Specified parameters measured in the tested samples ( pesticide residues and nitrate )</b> | <b>25</b>   |
| 3 : 1- Pesticide residues  | 25          |
| 3 :2- Nitrate content .  | 28          |
| <b>Part I V : Threshold pattern of the predominant groups main ingredients in the tested samples.</b>    | <b>31</b>   |
| 4 :1- Predominant groups in meat products.   | 31          |
| 4 :2- Predominant groups in legumes and vegetable.   | 34          |

|  |           |
|--|-----------|
| <b>Part V :Technological aspects and organoleptic evaluation of the tested samples .</b> | <b>36</b> |
| <b>Materials and methods</b>   | <b>41</b> |
| <b>Materials</b>   | <b>41</b> |
| <b>Methods</b>   | <b>42</b> |
| <b>I- Technological methods</b>  | <b>42</b> |
| I:1- <i>Kishk</i> samples .  | 42        |
| I:2- <i>Cake</i> sample.   | 44        |
| I:3- Faba bean paste ( <i>Bosara</i> )   | 46        |
| <b>II-Analytical methods</b>   | <b>47</b> |
| <b>2:1- Chemical analysis.</b>   | <b>47</b> |
| 2:1:1- Moisture , ash , minerals ,heavy metals , fat ,protein and crude fiber.           | 47        |
| 2:1:2- Amino acids .   | 48        |
| 2:1:3- Fatty acids.  | 49        |
| <b>2:2- Physical analysis</b>  | <b>50</b> |
| 2:2:1- Water adsorption  | 50        |
| 2:2:2- Sorption isotherm   | 50        |
| 2:2:3- Wettability   | 51        |
| 2:2:4- Water activity "aw"   | 51        |
| 2:2:5- Bulk density  | 51        |
| 2:2:6- Foaming capacity and foaming stability  | 51        |
| 2:2:7- Emulsion capacity and stability   | 52        |
| 2:2:8- Gelation properties   | 52        |
| 2:2:9- Rehydration ratio   | 52        |
| 2:2:10- Modulus of fineness.   | 53        |
| 2:2:11- Viscosity.   | 53        |
| <b>2:3- Pesticide residues</b>   | <b>54</b> |
| <b>2:4- Infrared spectrum ( functional groups )</b>                                      | <b>54</b> |
| <b>2:5- Sensory evaluation of the proposed samples</b>                                   | <b>54</b> |
| <b>III- Statistical analysis in term of Regression and analysis of variance</b>          | <b>55</b> |

|  |     |
|--|-----|
| <b>Results and discussion</b>  | 56  |
| <b>Part I: Major and minor constituents of the investigated samples</b>            | 56  |
| <b>1-Major constituents.</b>   | 56  |
| 1:1- Moisture.   | 56  |
| 1:2- Carbohydrate contents.  | 58  |
| 1:3- Ether extract and protein content:  | 64  |
| 1:4- Ash and crude fibers.   | 68  |
| 1:5- Minerals and heavy metals.  | 70  |
| 1:5:1- Minerals:   | 70  |
| 1:5:2- Heavy metals.   | 77  |
| <b>2- Minor constituents.</b>  | 82  |
| 2:1- Specified parameters based on amino acids.                                    | 82  |
| 2:1:1- Amino acids and ideal protein concept .                                     | 82  |
| 2:1:2- Essential amino acids :   | 86  |
| 2:1:3- Biological evaluation :   | 88  |
| 2:1:4- Side chains amino acids :   | 89  |
| 2:2- Fatty acids.  | 93  |
| 2:2:1- Saturated and mono unsaturated fatty acids:                                 | 93  |
| 2:2:2- Identified poly unsaturated fatty acids:                                    | 96  |
| 2:3- Ascorbic acid (vitamin C):  | 98  |
| <b>Part II: Physicochemical aspects of ingredients of the tested food stuffs .</b> | 102 |
| 2:1- Water absorption :  | 102 |
| 2:1:1- Water absorption in green bean and lentil.                                  | 102 |
| 2:1:2- Water absorption in germinated faba bean and fenugreek.                     | 103 |
| 2:1:3- Water absorption in kaki and strawberry.                                    | 104 |
| 2:2- Sorption isotherm in:   | 106 |
| 2:2:1- Sorption isotherm in luncheon and sausage.                                  | 106 |
| 2:2:2- Sorption isotherm in lentil and green bean.                                 | 108 |
| 2:2:3- Sorption isotherm in kaki and strawberry.                                   | 110 |
| 2:2:4- Sorption isotherm in germinated faba bean and                               | 111 |

|  |            |
|--|------------|
| fenugreek.   |            |
| 2:3- Wettability.  | 114        |
| 2:4- Water activity.   | 116        |
| 2:5- Bulk density.   | 120        |
| 2:6- Foaming capacity.   | 122        |
| 2:7- Emulsification .  | 129        |
| 2:8- Gelation properties.  | 134        |
| 2:9- Rehydration Ratio.  | 135        |
| 2:9:1- Rehydration of luncheon and sausage.  | 135        |
| 2:9:2- Rehydration of kaki and strawberry.   | 138        |
| 2:9:3- Rehydration of legume samples (lentil, green bean,<br>germinated fenugreek and germinated faba bean). | 141        |
| 2:10- Modulus of the tested samples.   | 145        |
| 2:10:1- Modulus of fineness in lentil and green bean .   | 145        |
| 2:10:2- Modulus of fineness in kaki and strawberry.  | 147        |
| 2:10:3- Modulus of fineness in germinated faba bean and<br>fenugreek.  | 149        |
| 2:10:4- Comparison between coarse , medium and fine  | 151        |
| 2:11- Flow rate :  | 152        |
| 2:11:1- Flow rate of luncheon and sausage.   | 153        |
| 2:11:2- Flow rate of kaki and strawberry.  | 156        |
| 2:11:3- Flow rate of legume samples (lentil, green bean, g.<br>fenugreek, g. faba bean).                     | 158        |
| <b>Part III: Specified parameters measured in the tested samples .</b>                                       | <b>163</b> |
| 3:1- Pesticide residues in the investigated samples  | 163        |
| 3:1:1- Pesticide residues in luncheon and sausage  | 163        |
| 3:1:2- Pesticide residues in kaki and lentil   | 168        |
| 3:1:3- Pesticide residues in green bean and strawberry   | 171        |
| 3:1:4- Pesticide residues in germinated fenugreek and<br>germinated faba bean.                               | 175        |
| 3:2- Nitrite content of the tested samples .   | 180        |
| 3:2- Ascorbic acid oxidase enzyme (AAO) in the investigated  | 182        |



samples .

|  |     |
|--|-----|
| <b>Part IV: Functional groups presented in the tested samples .</b>                        | 189 |
| 4:1- Functional groups in luncheon and sausage samples.                                    | 191 |
| 4:2 - Functional groups in lentil and green bean samples.                                  | 196 |
| 4:3 - Functional groups in kaki and strawberry samples.                                    | 199 |
| 4:4 – Intensities and functional groups in germinated and ungerminated fenugreek samples.  | 202 |
| 4:5- Intensities and functional groups in the tested germinated and ungerminated Faba bean | 205 |
| <b>Part V : General discussion</b>   | 208 |
| Scientific bases used for formulating and manufacturing of the suggested functional foods. | 208 |
| 5:1- Technochemical properties   | 219 |
| 5:2- Ascorbic acid oxidase enzyme of the processed samples.                                | 222 |
| 5:3- Emulsion  | 226 |
| 5:4- Functional groups of the processing samples   | 228 |
| 5:4:1- Functional groups of <i>kishk</i> samples   | 228 |
| 5:4:2- Functional groups of <i>cake</i> samples  | 230 |
| 5:4:3- Functional groups of Faba bean paste ( <i>Bosara</i> ) samples                      | 233 |
| 5:5- Organoleptic test in the processed samples.   | 236 |
| 5:5:1- Organoleptic tests for <i>kishk</i> samples.  | 236 |
| 5:5:2- Organoleptic tests for <i>cake</i> samples.   | 240 |
| 5:5:3- Organoleptic tests for Faba bean paste ( <i>Bosara</i> )samples.                    | 247 |
| <b>Summary and Conclusion</b>  | 251 |
| <b>Guide line for recommendation</b>   | 267 |
| <b>References</b>  | 268 |
| الملخص العربي  | 1   |

## **List of TABLES**

| <b>No</b> |  | <b>Page</b> |
|-----------|--|-------------|
| A         | Suggested recipes for manufacturing a control and a suggested <i>kishk</i> samples.                      | 42          |
| B         | The <i>cake</i> samples were prepared in the form of control and suggested recipes as follows:           | 44          |
| C         | Suggested recipes for manufacturing a control and a suggested faba bean paste ( <i>Bosara</i> ) samples. | 46          |
| 1         | Moisture content of the fresh and dried samples under investigation.                                     | 57          |
| 2         | Identified Sugars in the investigated samples (values are given as area % of HPLC chromatogram.          | 59          |
| 2-A       | One-way anova of fructose; dextrose; maltose; maltotriose and polysaccharides in the tested samples.     | 59          |
| 3         | Ether extract and Protein content of the investigated samples ( g /100g dry).                            | 65          |
| 4         | Ash and crude fiber contents of the investigated samples.  | 69          |
| 5         | Minerals content in the tested samples.  | 72          |
| 5-A       | Analysis of variance within the identified minerals and the tested samples.                              | 75          |
| 6         | Heavy metals content in the investigated samples .   | 79          |
| 6 - A     | Statistical pattern in terms of comparison between the investigated samples.                             | 81          |
| 7         | The ideal protein concept and the available energy of identified amino acids in the tested samples.      | 84          |
| 7 - A     | Analysis of variance of both ideal protein concept and amino acids of the tested samples.                | 85          |
| 8         | Essential amino acids (g/100g dry matter of protein) of luncheon and sausage of the tested samples.      | 87          |
| 9         | Protein efficiency ratio and protein quality of luncheon and sausage samples.                            | 88          |
| 10        | Uncharged and charged side chains amino acids of the luncheon and sausage samples                        | 90          |

|      |  |     |
|------|--|-----|
| 11   | Identified saturated and mono unsaturated fatty acids of the oil content extracted from edible portion of the tested luncheon and sausage samples. | 94  |
| 12   | Identified poly unsaturated fatty acids (polyenoic) in the oil content extracted from the edible portion of tested samples .                       | 97  |
| 13   | Level of vitamin C in the investigated samples. (mg / 100 g).  | 99  |
| 14   | Water absorbtion ( g H <sub>2</sub> O/2g sample ) of the investigated green bean and lentil samples.   | 103 |
| 15   | Water absorbtion ( g H <sub>2</sub> O/2g sample ) of the germinated faba bean and fenugreek samples.   | 104 |
| 16   | Water absorbtion (g H <sub>2</sub> O/2g sample ) of kaki and strawberry samples .  | 105 |
| 17   | Sorption isotherm (gH <sub>2</sub> O/2g sample) of the tested luncheon and sausage samples .   | 107 |
| 18   | Sorption isotherm (gH <sub>2</sub> O/2g sample) of the tested lentil and green bean samples under investigation.                                   | 109 |
| 19   | Sorption isotherm (gH <sub>2</sub> O/2g sample) of the tested kaki and strawberry samples under investigation.                                     | 110 |
| 20   | Sorption isotherm (gH <sub>2</sub> O/2g sample) of the tested germinated fenugreek and faba bean samples.  | 112 |
| 21   | Wetability in terms of (min/0.1 g sample) of the samples under investigation.  | 115 |
| 21-A | ANOVA: sausage; lentil; green bean; strawberry; germinated fenugreek and germinated faba bean.   | 116 |
| 22   | Water activity for the tested samples .  | 118 |
| 23   | Measurements of bulk densities of the investigated samples (g/ ml)   | 121 |
| 24   | Foaming capacity and foam stability cm/ml of the tested samples .  | 123 |
| 24-A | Principal component analysis and regression analysis of the foaming properties of the tested samples within 180 min.                               | 125 |

|      |  |     |
|------|--|-----|
| 25   | Emulsion pattern of the tested samples measured (over 100% of transmittance level ).   | 130 |
| 26   | Least gelation concentration (%) of the investigated samples.  | 135 |
| 27   | Rehydration pattern of the investigated luncheon and sausage samples.  | 136 |
| 27-A | Regression analysis based on multiplicative model for the relation between rehydration and time in meat products.                          | 137 |
| 27-B | Regression analysis based on multiplicative model for the relation between rehydration and time.   | 138 |
| 28   | Rehydration pattern of the investigated kaki and strawberry samples.   | 139 |
| 28-A | Regression Analysis based on multiplicative model for the relation between rehydration and time in fruit and vegetable.                    | 140 |
| 29   | Rehydration Pattern for the legume samples under investigation .   | 142 |
| 29-A | Regression Analysis based on multiplicative model for the relation between rehydration and time in legumes samples.                        | 143 |
| 29-B | Regression analysis based on multiplicative model for the relation between rehydration and time in the tested germinated fenugreek sample. | 144 |
| 30   | Modulus of fineness and modulus of uniformity of legume samples under investigation.   | 146 |
| 31   | Modulus of fineness and modulus of uniformity of kaki and strawberry samples.  | 148 |
| 32   | Modulus of fineness and modulus of uniformity of the tested germinated samples.  | 150 |
| 33   | Comparison between modulus of fineness and modulus of uniformity of all of the tested samples.   | 151 |
| 34   | Flow rate of different concentration of luncheon and sausage samples under investigation.  | 154 |

|      |   |     |
|------|---|-----|
| 34-A | Rgression analysis between $\text{KH}_2\text{PO}_4$ concentration and time in sec..   | 155 |
| 35   | Non Newtonian fluidity of strawberry and kaki samples.  | 157 |
| 36   | Flow rates ,specific and reduced viscosities of the tested legumes samples .  | 160 |
| 36-A | Regression Analysis: CON% versus time of legumes samples in sec.  | 161 |
| 37   | Pesticide residues identified in the luncheon sample.   | 164 |
| 38   | Pesticide residues identified in the sausage sample.  | 166 |
| 39   | Pesticide residues identified in the lentil sample.   | 169 |
| 40   | Pesticide residues identified in the green bean sample.   | 172 |
| 41   | Pesticide residues identified in the strawberry sample .  | 173 |
| 42   | Pesticide residues identified in the germinated fenugreek sample.   | 175 |
| 43   | Pesticide residues identified in the germinated Faba bean sample .  | 177 |
| 44   | Nitrite content (mg/ kg sample) analyzed within the investigated samples.   | 181 |
| 45   | Activity of ascorbic acid oxidase enzyme (AAO) based on vitamin C content of the investigated samples   | 183 |
| 45-A | Statistical analysis of ascorbic acid oxidase enzyme in terms of linear regression of the residual amount of ascorbic acid in the tested samples. | 184 |
| 46   | Intensities and functional groups identified in the tested luncheon and sausage samples.  | 194 |
| 47   | Intensities and functional groups identified in the tested lentil and green bean samples.   | 197 |
| 48   | Intensities and functional groups identified in kaki and strawberry samples under investigation .   | 200 |
| 49   | Intensities and functional groups identified in germinated and ungerminated fenugreek samples.  | 203 |
| 50   | Intensities and Functional groups identified in germinated and ungerminated broad bean samples .  | 206 |

|      |  |     |
|------|--|-----|
| 51   | Accumulated major constituents analyzed in the                   | 210 |
|      | investigated food samples.                                       |     |
| 52   | Accumulated functional parameters applied in the                 | 215 |
|      | investigated food samples.                                       |     |
| 53   | Improvement of functional groups as a function of the            | 218 |
|      | application of the suggested recipes.                            |     |
| 54   | Improvement level in selected parameters in the tested           | 220 |
|      | samples.   |     |
| 55   | Activity of ascorbic acid oxidase based on the                   | 223 |
|      | concentration of vitamin C content of the investigated           |     |
|      | samples.   |     |
| 55-A | Regression analysis of ascorbic acid oxidase based on the        | 224 |
|      | residual amounts of ascorbic acid in the tested samples.         |     |
| 56   | Emulsion for the processing samples .                            | 226 |
| 57   | Comparison between the intensity and functional groups of        | 228 |
|      | the tested <i>kishk</i> samples.                                 |     |
| 58   | Comparison between the intensitiy and functional groups of       | 231 |
|      | the tested <i>cake</i> samples.                                  |     |
| 59   | Comparison between the intensitiy and functional groups of       | 234 |
|      | the tested Faba bean paste ( <i>Bosara</i> ) .                   |     |
| 60   | Organoleptic evaluation of the investigated kishk sample.        | 237 |
| 60-A | ANOVA: Colour; Appearance; Flavour; After Taste;                 | 239 |
|      | Aroma; Overall Acceptabilty and Structure in <i>kishk</i> sample |     |
|      | .  |     |
| 61   | Organoleptic evaluation of the <i>cake</i> sample under          | 241 |
|      | investigation.   |     |
| 61-A | ANOVA: Colour; Appearance; Flavour; After Taste;                 | 242 |
|      | Aroma; Overall Acceptability and Height in <i>cake</i> sample.   |     |
| 62   | Organoleptic evaluation of the Faba bean paste ( <i>Bosara</i> ) | 248 |
|      | sample.  |     |
| 62-A | Correlations between Colour; Appearance; Flavour; After          | 249 |
|      | Taste; Aroma; Overall acceptability and structure in faba        |     |
|      | bean paste ( <i>Bosara</i> ) sample.                             |     |

## **List of FIGURS**

| <b>No</b> |   | <b>Page</b> |
|-----------|---|-------------|
| 1         | Moisture content of the fresh and dried samples under test  | 58          |
| 2         | HPLC profiles of sugars identified in the tested meat , fruit and vegetable samples.                          | 60          |
| 3         | HPLC profiles of sugars identified in the tested legumes.   | 61          |
| 4         | Ether extract and protein content for the tested samples  | 66          |
| 5         | Ash and crude fiber of the tested samples.  | 69          |
| 6         | Mineral levels in the tested samples in terms of their No.of folds of the responded highest level.            | 74          |
| 6-A       | Distribution of the tested minerals within the tested samples based on principal components analysis          | 76          |
| 7         | Heavy metals content (mg/100g dry sample) in the tested samples.  | 80          |
| 8         | Comparision between the available energy (K.cal/mol) of the identified amino acids within the tested samples. | 86          |
| 9         | Uncharged and charged side chains amino acids of luncheon and sausage.  | 91          |
| 10        | Identified saturated and mono unsaturated fatty acids of the tested samples .                                 | 95          |
| 11        | Identified poly unsaturated fatty acids (poly enoic in the oil extracted from luncheon.                       | 98          |
| 12        | Level of vitamin C in the tested samples  | 100         |
| 13        | Average of sorption isotherm of luncheon and  | 108         |