



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ





شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الالكتروني والميكرو فيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
على هذه الأفلام قد أعدت دون أية تغييرات



يجب أن

تُحفظ هذه الأفلام بعيدا عن الغبار

في درجة حرارة من 15-25 مئوية ورطوبة نسبية من 20-40%

To be Kept away from Dust in Dry Cool place of
15-25- c and relative humidity 20-40%



شبكة المعلومات الجامعية

التوثيق الالكتروني والميكروفيلم

Recent surgical techniques in management of Pilonidal sinus

Essay

*Submitted for partial fulfillment of Master degree in General
Surgery*

By

Amira Yehia Mohammed Abu Taleb
M.B.B.Ch.

Prof. Sayed Rashad **Supervisors**

Prof. Dr. Sayed Mohamed Rashad

Professor of General Surgery
Ain Shams University

Prof. Dr. Amr Ali Salah Mahmoud


Professor of Plastic and Reconstructive Surgery
Ain Shams University

Dr. Ayman Mohamed Shaker

Lecturer of General Surgery
Ain Shams University

Faculty of Medicine
Ain shams University

(2005)



بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

وَقُلْ رَبِّ زِدْنِي عِلْمًا

صدق الله العظيم

سورة طه : آية (١١٤)

Acknowledgement

First and foremost to "Allah"

I would like to express my sincere gratitude to Prof. Dr. Sayed Mohamed Rashad, Prof. of General Surgery, Faculty of Medicine, Ain Shams University, for his fatherly guidance, continuous help and support through this work.

To Prof. Dr. Amr Ali Salah Mahmoud, Professor of Plastic and Reconstructive Surgery, Faculty of Medicine, Ain Shams University, Thanks for guidance, patience and effort you have exerted.

To Dr. Ayman Mohamed Shaker, Lecturer of General Surgery Faculty of Medicine, Ain Shams University, thanks for his kind supervision, encouragement and great effort.

Last but not least, to my family, friends and colleagues.

List of Figures

Fig.	Title	Page
1	Pilonidal sinus-Primary opening and secondary track	13
2	Stages of pilonidal disease from top to bottom	14
3	Excision of midline pits	57
4	Lateral drainage of abscess cavity	58
5	Suturing of subcutaneous flap to the skin bridge and subcuticular closure of midline wound	59
6	Cleft lift procedure, the buttocks are pushed together before surgery	61
7	The ellipse of the skin is removed	61
8	A: The area of skin to be mobilized as a full thickness skin flap. B: The left flap will be removed and the right flap will provide coverage	62
9	Following total excision of the midline sinus and its lateral tracks the incision made at 45 degree to the midline	64

- | | | |
|----|--|----|
| 10 | The tow triangular flaps with their apices at A and B are mobilized to include skin and fat down to the level of the sacral fascia. Further mobilization of the other tissues may be required at the same level. The triangular flaps are transposed at the skin sutured | 64 |
| 11 | Local rectangular rotation flap operation. | 67 |
| 12 | A: Skin markings for construction of the rhomboid flap. Incisions are made along all solid lines and the shaded area is excised.
B: Rhomboid flap being transposed | 68 |
| 13 | Left: a biconcave incision started and centered 2 cm to side of the midline cleft and shirting the cleft just sufficiently to excise the primary opening (•) and the whole track. | 76 |
| 14 | On the right is cross- section of the center of the wound indicated on the left with a dashed line. | 77 |
| 15 | The deepest suture are placed before any is tied | 78 |

List of Contents

Introduction	1
Aim of the work	4
Review of Literature	
Pathology of pilonidal sinus	5
Presentation of pilonidal sinus	17
Management of pilonidal sinus	25
Surgical	37
Non surgical	90
Summary	95
References	97
Arabic summary	



Introduction

