Assessment of Pulmonary Hypertension in Critically III Pediatric Patients with Acute Respiratory Distress Syndrome (ARDS)

Thesis

Submitted for Partial fulfillment of Master Degree in **Pediatrics**

By **Ahmed Gamal Ibrahim Yousif**MB.B.Ch. Dec,

Under Supervision of

Prof. Dr. Hanan Mohamed Ibrahim

Professor of Pediatrics Faculty of Medicine – Ain Shams University

Dr. Nevin Mohamed Mamdouh

Assistant Professor of Pediatrics Faculty of Medicine – Ain Shams University

Dr. Marwa Moustapha Attia AL-Fahham

Lecturer of Pediatrics Faculty of Medicine – Ain Shams University

> Faculty of Medicine Ain Shams University

تقييم ارتفاع ضغط الدم الرئوى لدى الحالات الحرجة من الأطفال المصابين بمتلازمة الضائقة التنفسية الحادة

رسالة توطئة للحصول على درجة الماجستير في طب الأطفال

مقدمة من الطبيب/ أحمد جمال إبر اهيم يوسف بكالوريوس الطب والجراحة؛ ديسمبر، ٢٠٠٧

تحت إشراف أ.د. حنان محمد إبراهيم أستاذ طب الأطفال كلية الطب – جامعة عين شمس

د/ نيفين محمد ممدوح أستاذ مساعد طب الأطفال كلية الطب – جامعة عين شمس د/مروة مصطفى عطية الفحام مدرس طب الأطفال كلية الطب جامعة عين شمس

> كلية الطب جامعة عين شمس ٢٠١٣



سورة البقرة الآية: ٣٢



First of all, great thanks to GOD who enabled us to complete this work, hoping to provide a useful guide to the scope of ARDS and pulmonary hypertension.

No words can express my deepest appreciation and profound respect to **Prof. Dr. Hanan Mohamed Ibrahim,** Professor of Pediatrics, Ain Shams University, for her continuous guidance, support through the work. She has generously devoted much of her time and her effort for planning and supervision of this study. It was great honor to work under her supervision.

Also, my profound gratitude to **Dr. Nevin Mohamed Mamdouh**, Assistant of Pediatrics, Ain Shams University, for her kind supervision and support to complete this work...

Also, my profound gratitude to **Dr. Marwa Moustapha Attia AL-Fahham,** Lecturer of Pediatrics, Ain Shams University for her kind supervision and support.

Also a special thanks to all staff members of PICU Unit- Ain Shams University, for their continuous great support.

Also I don't forget my family, the best helper for me, their full support, prayers and wishes were a great motive to accomplish this work. My deepest gratitude to them and thanks will never appreciate what I owe them.

Lastly special thanks to my wife who supported me a lot.



Contents

Title	Page	
List of Abbreviation		I
List of Figures		IV
List of Tables		VII
Introduction		
Aim of the Work,		
Review of Literature:		
Chapter (): Acute Respiratory D (ARDS)	•	
Chapter (): Treatment of ARDS		
Chapter (): Pulmonary Hypertension	ı	
Patients and Methods		
Results	•••••	
Discussion		
Summary and Conclusion		
Recommendations		
References		
Arabic Summary		

List of Abbreviations

Abb.	Meaning
°-HTT	Serotonin transporter gene
AECC	The American-European Consensus Committee
ALI	Acute lung injury
ALK \	Activin-receptor-like kinase gene
AP	Acute pancreatitis
APAH	Associated Pulmonary arterial hypertension
ARDS	Acute respiratory distress syndrome
BALF	Bronchalveolar lavage fluid
BMPR [*]	Bone morphogenic protein receptor-
BMPR-Y	Bone morphogenetic receptor protein gene
BOOP	Bronchiolitis obliterans with organizing pneumonia
BPN	Bronchopneumonia
CDC	Center for Disease Control
cGMP	Cyclic guanosine monophosphate
CPS	Carbamyl-phosphate synthetase gene
CS	Cigarette smoke
CXR	Chest X-ray
DAD	Diffuse alveolar damage
DAHS	Diffuse alveolar hemorrhage syndrom
DIC	Disseminated intravascular coagulation
DVT	Deep venous thrombosis
ECG	Examination, electrocardiogram
ec-NOS	Nitric oxide synthase gene
EF	Ejection fraction
EGF	Epidermal growth factor
ET-1	Endothelin-
FIO	Fractional concentration of oxygen in inspired air
FS	Fractional shortening
GER	Gastroesophageal reflux
HFOV	High frequency oscillatory ventilation
HFV	High-frequency Ventilation
HP	Hypersensitivity pneumonitis
ICU	Intensive care unite

Abb.	Meaning
IGF	Insulin like growth factor
IL-1	Interleukin-
IL-\β	IL- beta
ILD	Interstitial lung disease
INOS	Inducible nitric oxide synthesis
IPAH	Idiopathic Pulmonary arterial hypertension
MODS	Multiorgan dysfunction syndrome
MRSA	Methicillin-resistant S. aureus
NO	Nitric Oxide
NPPV	Non-invasive positive pressure ventilation
OLC	Open lung concept
PA pressure	Pulmonary arterial pressure
PAAT	Pulmonary artery acceleration time
PaO	Partial arterial pressure of oxygen
PAP	Pulmonary alveolar proteinosis
PASMC	Pulmonary artery smooth muscle cells
PASP	Pulmonary artery systolic pressure
PCH	Pulmonary capillary haemangiomatosis
PDE	Phosphodiesterase enzyme
PDE-°	PDE- isoenzymes.
PDGF	Platelet derived growth factor
PH	Pulmonary hypertension
PIP	Peak inspiratory pressure
PPHN	Persistent pulmonary hypertension of the newborn
Ppm	Part per million
PVOD	Pulmonary veno-occlusive disease
PVR	Pulmonary vascular resistance
PVRi	Pulmonary vascular resistance index
Qp	Transpulmonary flow
RAP	Right atrial pressure
RVSP	Right Ventricular systolic pressure
SD	Standar deviation.
SIMV-PC	Synchronized intermittent Mandatory ventilation –
	pressure controle
SIRS	Systemic onset inflammatory response

Abb.	Meaning
SL-selectin	Soluble L-selectin
SVR	Systemic vascular resistance
TNF-α	Tumour necrosis factor
TRALI	Transfusion-related acute lung injury
TRV	Tricuspid regurgitation velocity
TTE	Trans thoracic Echocardiography
TVIRVOT	Right ventricular outflow tract time-velocity integral
V	Velocity
V/Q	Ventilation/perfusion
VILI	Ventilation induced lung injury
VRE	Vancomycin-resistant enterococci
WHO	World Health Organization
Δp	Transpulmonary pressure gradient

List of Figures

Figure	Title	Page
Fig. (1)	Light Microscopy and Electron Microscopy	
	during the Acute Phase (Panels A and D) and	
	the Fibrosing-Alveolitis Phase (Panels B, C,	
	and E) of Acute Lung Injury and the Acute	
	Respiratory Distress Syndrome	
Fig. (7)	An anteroposterior chest radiograph from a	
	patient with ARDS associated with gram-	
	negative sepsis who was receiving	
	mechanical ventilation	
Fig. (*)	High-resolution computed tomography scan	
	of the chest, slice : ground-glass infiltrate	
	and septal thickening predominantly in the	
	upper thirds, as well as bilateral pleural	
	effusion (a typical finding)	
Fig. (1)	Pulmonary arterial hypertension: potential	
, ,	pathogenetic and pathobiological mechanisms	
Fig. (•)	Electrocardiogram demonstrating the changes	
_, ()	in a patient with pulmonary hypertension.	
Fig. (٦)	(B) Apical chamber view. Common	
- , (.)	echocardiographic findings in PAH	
Fig. (Y)	Determination of pulmonary artery pressure	
T	using Doppler echocardiography.	
Fig. (^)	Images showing peak tricuspid regurgitant	
	velocity (TRV) and right ventricular outflow	
	time-velocity integral (TVIRVOT) in a	
	patient with normal pulmonary vascular	
	resistance (PVR).	

Figure	Title	Page
Fig. (٩)	Images showing TRV and TVIRVOT in a	
T ' (,)	patient with elevated PVR.	
Fig. (1.)		
Fig. (۱۱)		
	ultrasound, Horten, Norway) and its probe	
	(*S).	
Fig. (17)	A: D echocardiography showing dilated	
	right atrium (apical chamber view) in a	
	patient with ARDS. B: Color flow Doppler	
	showing severe tricuspid regurgitation in	
	patient with ARDS (apical chamber view).	
Fig. (۱۳)	CW Doppler showing increased TRV and	
	elevated RVSP in a patient with ARDS	
	(apical chamber view).	
Fig. (\ \ \ \)	RVOT/TVI detected by Pulsed wave Doppler	
J	at the level of RVOT (short axis parasternal	
	view).	
Fig. (10)	,	
- -8. ()	by pulsed wave Doppler at the level of	
	pulmonary valve (short axis parasternal	
T' (4)	view).	
Fig. (١٦)	Gender distribution in the study population	
Fig. (۱۷)	Percentage of indication for ICU admission	
	among the studied population	
Fig. (١٨)	Percentage of studied population as regard the	
	etiology of ARDS	
Fig (19)	Box plot for ARDS score in the study group.	

Figure	Title	Page
	Box represents interquartile range. Horizontal	
	line across box represents median. Error bars	
	represent thand th percentiles	
Fig. (۲۰)	Box plot for pH at diagnosis and on	
	echocardiography. Box represents	
	interquartile range. Horizontal line across box	
	represents median. Error bars represent th	
	and th percentiles and marker lying beyond	
	these limits represents an outlier	
Fig. (۲۱)	Initial ventilation mode in the study	
	population	
Fig. (۲۲)	Shows that of studied patients were	
	ventilated on SIMV-PC during Echocardiography	
Fig. (۲۳)	Box plot showing PVR in controls and	
± -8· \ /	patients with ARDS	
Fig. (7 £)	Box plot showing RVSP in controls and	
	patients with ARDS	
Fig. (۲0)	Prevalence of high RVSP in controls and	
8- (patients with ARDS	
Fig. (۲٦)	Prevalence of high PVR in controls and	
8.	patients with ARDS	
Fig. (۲۷)	Scatter plot for correlation between PIP on	
6. ()	echocardiography and PVR	
Fig. (۲۸)	Kaplan-Meier Survival curves defined by	
-	etiology of ARDS. Vertical dashes along	
	survival curves represent censored data.	

List of Tables

Table	Title	Page
Tab. (1)	Parameters used in the detection of the Lung	
	injury score	
Tab. (7)	Modified table showing common Causes	
	and predisposing factors of acute respiratory	
	distress syndrome	
Tab. (٣)	Summary of drugs used for treatment of	
	pulmonary hypertension	
Tab. (1)	Modified table showing revised WHO	
	classification of PH:	
Tab. (•)	Strength of Association	
Tab. (٦)	Descriptive data of the studied population	
Tab. (Y)	Percentage of indication for ICU admission	
	among the studied population	
Tab. (^)	Percentage of studied population as regards	
	the etiology of ARDS	
Tab. (٩)	Distribution of ARDS score	
Tab. (۱۰)	ABG parameters of studied population at	
	time of diagnosis of ARDS	
Tab. (۱۱)	ABG parameters of studied population at	
	time of echocardiography	
Tab. (۱۲)	Description of Initial ventilator mode	
	among studied population	
Tab. (۱۳)	Description of ventilator mode among	
	studied population on echocardiography	
Tab. (۱٤)	Initial ventilator setting among studied	
	population	

Table	Title	Page
Tab. (۱٥)	Ventilator setting on echocardiography	
	among studied population	
Tab. (۱٦)	Left ventricular systolic functions	
	dimentions among studied population	
Tab. (۱۷)	Echocardiographic parameters assessing	
	pulmonary hypertension and pulmonary	
	vascular resistance among studied	
	population	
Tab. (۱۸)	Comparison between cases and control as	
	regards PVR and RVSP	
Tab. (۱۹)	Comparison between pulmonary systolic	
	hypertension and PVR. In patients and	
	control	
Tab. (۲۰)	Comparison of left ventricular systolic	
	functions dimentions indices in survivors	
	and non-survivors	
Tab. (۲۱)	Comparison of various echocardiographic	
	parameters indices assessing pulmonary	
	hypertension and pulmonary vascular	
	resistance in survivors and non-survivors	
Tab. (۲۲)	Comparison of left ventricular systolic	
	functions dimentions indices in SIMV-PC	
	mode and HFOV mode	
Tab. (۲۳)	Comparison of various echocardiographic	
	parameters indices assessing pulmonary	
	hypertension and pulmonary vascular	
	resistance in SIMV-PC mode and HFOV	
	mode	

Table	Title	Page
Tab. (۲٤)	Correlation among left ventricular systolic	
	functions dimentions indices and ABG	
	indices at diagnosis	
Tab. (۲٥)	Correlation among various	
	echocardiographic parameters indices	
	assessing pulmonary hypertension and	
	pulmonary vascular resistance among	
	studied population and ABG indices at	
	diagnosis	
Tab. (۲٦)	Correlation among left ventricular systolic	
	functions dimentions indices and ABG	
	indices at echocardiographic examination	
Tab. (۲۷)	Correlation among various	
	echocardiographic parameters indices	
	assessing pulmonary hypertension and	
	pulmonary vascular resistance among	
	studied population and ABG indices at	
	diechocardiography Completion among left ventricular systelia	
Tab. (۲۸)	Correlation among left ventricular systolic functions dimentions indices and initial	
	ventilator setting	
Tab. (۲۹)	Correlation among various	
1ab. (' ')	echocardiographic parameters indices	
	assessing pulmonary hypertension and	
	pulmonary vascular resistance among	
	studied population and initial ventilator	
	setting	

Table	Title	Page
Tab. ("\)	Correlation among left ventricular systolic	
	functions dimentions indices and ventilator	
	setting during echocardiography	
Tab. (٣١)	Correlation among various	
	echocardiographic parameters indices	
	assessing pulmonary hypertension and	
	pulmonary vascular resistance among	
	studied population and ventilator setting	
	during echocardiographic examination	
Tab. (٣٢)	Distribution of the studied cases as regards	
	outcome measures	
Tab. (٣٣)	Survival analysis according to etiology of	
	ARDS	
Tab. (٣٤)	Comparison of survival curves by etiology	
	of ARDS (Log-rank test)	
Tab. (٣0)	ARDS Score and PaO /FiO among	
	survivors and Non-Survivors	