



Development of sensitive HPLC method with fluorescence detection for determination of memantine and its application to pharmacokinetic study

A thesis Presented by

Mohamed Gamaleldin Hassan Oraby

B. Pharm. Sci., Al-Azhar University Assiut branch, 1999

For partial fulfillment of

**The degree of PhD in Pharmaceutical Sciences
(Pharmaceutical Analytical Chemistry)**

Under supervision of

Prof. Dr. Horria A. Mohamed

Department of Pharmaceutical Analytical Chemistry
Faculty of Pharmacy
Assiut University

Prof. Dr. Hanaa M. Abd El-Wadoud

Department of Pharmaceutical Analytical Chemistry
Faculty of Pharmacy
Assiut University

Prof. Dr. Kenichiro Nakashima

Department of Clinical Pharmacy
School of Pharmaceutical Sciences
Nagasaki University, Japan

Faculty of Pharmacy
Assiut University
2013

يَسْمِ اللّٰهُ الرَّحْمٰنَ الرَّحِيْمَ

In the Name of Allâh, the Most Beneficent, the Most Merciful.

نَرْفَعُ دَرَجَاتٍ مِّنْ نَّشَأٍ وَفَوْقَ
كُلِّ ذِي عِلْمٍ عَلِيمٌ

We raise to degrees whom We please, but over all those
endowed with knowledge is the All-Knowing (Allâh).

صَدَقَ اللّٰهُ الْعَظِيْمُ

76 verse from



Acknowledgement

First and foremost, all praise and glory to the Almighty ALLAH, the most kind and merciful to whom I am always indebted. Thee do I worship and Thee aid I seek.

My sincere thanks and gratitude go to Professor **Horria Abdel-Maged Mohamed**, for her kind support, stimulating suggestions, enthusiastic guidance and encouragement helped me in all the time of the research and writing this thesis.

My sincere appreciation to Professor **Hanaa Mohamed Abdel-Wadood**, for her sincere help, guidance, kind support, valuable scientific comments and supervision throughout this work.

I would like to express my sincere gratitude and deep appreciation to Professor **Kenichiro Nakashima**, for his encouragement, valuable suggestions and support in the course of this research.

I wish to express my warm thanks for Associate Professor **Mitsuhiro Wada**, for his constant support and helpful suggestions and for his great effort during the research period.

I sincerely thank Assistant Professor **Rie Ikeda**, for her valuable, detailed and constructive comments that contributed to this thesis.

I wish also to express my thanks and appreciation to all members in the division of pharmacoinformatics, department of clinical pharmacy, course of pharmaceutical sciences, graduate school of biomedical sciences, Nagasaki University.

I wish also to express my thanks and appreciation to all members of the department of pharmaceutical analytical chemistry, faculty of pharmacy, Al- Azhar University, Assiut branch and to all members of the department of pharmaceutical analytical chemistry, faculty of pharmacy, Assiut University and my deepest gratitude goes to my mother, ants, brothers and my sister for their love, continuous encouragement and support throughout my life.

I extend heartfelt thanks to my wife for her understanding, comforting, support and, above all, bringing me the most impressive scene, I have ever met, on the birth of my daughter and my sons.

Mohamed Gamaleldin Hassan Oraby

List of contents

	Page
Contents	i
List of publications	v
List of abbreviations	vi
List of tables	lx
List of figures	xi
Summary	1
Introduction	6
A. General Introduction	7
A.1. Adamantane	8
A.2. Alzheimer's disease	9
A.2.1. Treatment of Alzheimer's disease	10
A.2.2. Memantine	12
a. Neuropharmacological basis for MT use in the treatment of Alzheimer's disease	12
b. Pharmacology of memantine	15
c. Pharmacokinetics of memantine	18
d. Dosage and administration	19
B. Analytical Review	21
B.1. Spectrometric methods	22
B.2. Separation methods	24
Scope of investigation	30

Chapter I: Determination of memantine by HPLC with fluorescence detection	
using 4 - (4,5 - Diphenyl - 1 <i>H</i> - imidazol - 2 - yl)benzoyl	
chloride (DIB-Cl) as labeling reagent	33
I.1. Introduction	34
I.2. Experimental	37
I.2.1. Instruments	38
I.2.2. Chemicals and reagents	39
I.2.3. Preparation of reagents	40
I.2.4. Labeling with DIB-Cl	41
I.2.5. Optimization of chromatographic conditions	41
I.2.6. Preparation of synthesized DIB-MT derivative	42
I.3. Results and discussion	43
I.3.1. Mechanism of reaction of DIB-Cl with MT	44
I.3.2. Optimization of the labeling conditions	46
I.3.3. Identification of synthesized DIB-MT	52
I.3.5. Chromatographic separation	54
I.3.6. Method validation	60
i. Linearity range	60
ii. Limit of detection and limit of quantitation	60
iii. Method precision	62
iv. Robustness	64
Chapter II: Interaction study of methazolamide, donepezil and galantamine on	
the pharmacokinetics of memantine in rat plasma.	66
II.1. Introduction	67
II.2. Experimental	70

II.2.1. Instruments	71
II.2.2. Chemicals and reagents	71
II.2.3. Preparation of reagents	71
II.2.4. Preparation of MT, MZA, DP and GAL for injection to rat	71
II.2.5. Animal treatment	72
II.2.6. Pharmacokinetic of MT in rat plasma	72
II.2.7. Labeling with DIB-Cl	73
II.2.8. Clean-up with solid phase extraction	73
II.3. Results and discussion	75
II.3.1. Chromatographic separation	76
II.3.3. Solid phase extraction conditions	76
II.3.4. Method validation	79
i. Linearity range	79
ii. Limit of detection and limit of quantitation	81
iii. Method precision	84
iv. Robustness	86
II.3.5. Application of the method for the pharmacokinetic study of memantine	88
II.3.5.1. Pharmacokinetic study after co-administration of MT without/with methazolamide	88
II.3.5.2. Pharmacokinetic study after co-administration of MT without/with acetylcholinesterase inhibitors	94
a. Pharmacokinetic study of memantine co-administered with donepezil.	95
b. Pharmacokinetic study of memantine co-administered with galantamine.	98

Conclusion	102
References	105
Summary in Arabic	114

List of publications

- 1- **Mohamed G. Hassan**, Kamla M. Emara, Horria A. Mohamed, Hanaa M. Abdel-Wadood, Rie Ikeda, Mitsuhiro Wada, Naotaka Kuroda and Kenichiro Nakashima: Determination of memantine in rat plasma by HPLC-fluorescence method and its application to study of the pharmacokinetic interaction between memantine and methazolamide. **Biomedical Chromatography**; **26** (2012), 214-219.
- 2- **Mohamed G. Hassan**, Rie Ikeda, Mitsuhiro Wada, Naotaka Kuroda, Hanaa M. Abdel-Wadood, Horria A. Mohamed and Kenichiro Nakashima: Interaction study of acetylcholinesterase inhibitors on pharmacokinetics of memantine in rat plasma by HPLC-fluorescence method. **Biomedical Chromatography**; **27** (2013), 1685-1689.

List of abbreviations

AChE	Acetylcholinestrerase
AD	Alzheimer's disease
ASC	Anthraquinone sulfonyl chloride
AUC	Area under curve for concentration versus time
CAI	Carbonic anhydrase inhibitor
CL	Clearance
C_{max}	The peak concentration
Conc.	Concentration
CYP450	Cytochrome P450 isoenzymes
DIB-Cl	4-(4,5-diphenyl-1 <i>H</i> -imidazol-2-yl)benzoyl chloride
DMF	Dimethylformamide
DP	Donepezil
EDTA	Ethylenediamine tetra-acetic acid
EMA	European Medicines Agency
FAB-MS	Fast atom bombardment mass spectrometer
FDA	Food and Drug Administration
FL	Fluorescence
FLX	Flouxetine
Fmol	Femtomole
Fmoc	9-fluorenylmethyl chloroformate
GAL	Galantamine
HPLC	High Performance Liquid Chromatography
Inj.	Injection
<i>i.p.</i>	Intraperitoneal
IS	Internal standard

<i>i.v.</i>	Intravenous
IUPAC	International Union of Pure and Applied Chemistry
LIF	Laser induced fluorescence
MEKC	Micellar electrokinetic chromatography
MRT	Mean resident time
MT	Memantine (1-amino-3,5-dimethyladamantane)
MZA	Methazolamide
NAC	(2-naphthoxy)acetyl chloride.
NBD-Cl	4-Chloro-7-nitro-2,1,3-benzoxadiazole
nAChRs	Nicotinic acetylcholine receptors
NMDA	<i>N</i> -methyl-D- aspartate.
OPA	<i>o</i> -phthaldialdehyde
RFI	Relative fluorescence intensity
RIV	Rivastigmine
SPE	Solid phase extraction
THF	Tetrahydrofuran
E	Molar absorbitivity

List of tables

Table		Page
1	Reported HPLC methods for the determination of memantine	26
2	Effect of buffer type on the RFI of MT due to labeling with DIB-Cl (1.5 mM).	47
3	Elemental analysis of DIB-MT	52
4	UV-, FL-spectral data and FL quantum yield of DIB-MT in several solvents	55
5	Quantitative parameters of the proposed method for standard MT	61
6	Method precision for standard MT using the proposed HPLC method	63
7	Robustness of the proposed method for standard MT	65
8	Quantitative parameters of the proposed method for rat plasma spiked with MT	80
9	A comparison of the proposed method with other reported methods	82
10	Method precision for rat plasma spiked with MT	85
11	Robustness of the proposed method for rat plasma spiked with MT	87
12	Pharmacokinetic parameters of MT (2.5 mg/kg) without/with MZA (5.0 mg/kg) in plasma following a single <i>i.p.</i> administration to rats	92
13	Pharmacokinetic parameters of MT (2.5 mg/kg)	

	without/with DP (5.0 mg/kg) in plasma following a single <i>i.p.</i> administration to rats	97
14	Pharmacokinetic parameters of MT (2.5 mg/kg) without/with GAL (3.0 mg/kg) in plasma following a single <i>i.p.</i> administration to rats	99

List of figures

Figure		Page
1	Schematic representation of the NMDA receptor complex	13
2	HPLC-FL system for determination of MT	39
3	Labeling reaction of MT (100 ng/mL) with DIB-Cl (1.5 mM)	45
4	Effect of DIB-Cl concentration on the peak height of reaction product with (■) 50 ng/ml of FLX and (Δ) 100 ng/ml of MT	46
5	Effect of carbonate buffer pH on the peak height of the reaction product of DIB-Cl with (■) 50 ng/ml of FLX and (Δ) 100 ng/ml of MT	48
6	Effect of carbonate buffer concentration on the peak height of the reaction product of DIB-Cl with (■) 50 ng/ml of FLX and (Δ) 100 ng/ml of MT	49
7	Effect of heating temperature on the peak height of the reaction product of 100 ng/ml of MT with DIB-Cl (1.5 mM) at (□) 25°C, (■) 45°C, (▲) 60°C and (x) 80°C	51
8	Stability time of the derivatives; 50 ng/ml FLX (■) and (▲) 100 ng/ml of MT (n=3).	51
9	Fast atom bombardment mass spectrometry for DIB-MT	53
10	Effect of acetonitrile ratio on peak height (A) and on retention time (B) of 50 ng/mL of IS (■) and 100 ng/mL of MT (▲)	57
11	Chromatograms of reagent blank (A) and that for standards FLX, 50 ng/mL and MT, 100 ng/mL (B)	58
12	Chemical structure of FLX (IS)	59