







شبكة المعلومــات الجامعية التوثيق الالكتروني والميكروفيلم



جامعة عين شمس

التوثيق الالكتروني والميكروفيلم



نقسم بللله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأفلام قد اعدت دون آية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

40-20 في درجة حرارة من 15-20 منوية ورطوبة نسبية من

To be kept away from dust in dry cool place of 15 – 25c and relative humidity 20-40 %









B914-5

LOCALIZATION OF THE KAPPA AND LAMBDA LIGHT CHAINS IN REACTIVE AND NEOPLASTIC LESIONS OF THE MUCOSA ASSOCIATED LYMPHOID TISSUE (MALT)

Thesis
Submitted for the Partial Fulfillment
of the Requirement of the M.D. Degree in
Pathology

By Walid Mohammed Sharaf M.B. B.Ch., M.Sc

Supervisors

Prof. Dr. Effat Ramadan. El Sherif

> Prof. of Pathology Cairo University

Prof. Dr.

Nashwa Shafik Aggour

Prof. of Pathology National Research Center Prof. Dr.

Ali Ahmed El Hindawy

Prof. of Pathology Cairo University

FACULTY OF MEDICINE CAIRO UNIVERSITY 2000



AKNOWELDEGMENT

I would like to express my deepest and cordial appreciation to **Prof. Dr. Effat El Sherif**, Professor of Pathology, Cairo University, for her constant support and encouragement and whose guidance has shown me my way in this work. She gave me a lot of her experience, and new ideas.

I wish also, to acknowledge my heartful and deepest gratitude to *Prof. Dr. Nashwa Aggour*, Professor and Head of Pathology Department, National Research Center, for her faithful supervision, guidance and encouragement. I am also thankful for her support and interest throughout all stages of the work.


I am greatly indebted to *Prof. Dr. Aly El Hindawy*, Professor of Pathology, Cairo University, who gave me a lot of his pointfull ideas and stimulating suggestions concerning this work. He supplied me with a lot of his own experience, assistance and encouragement. The credit of bringing this work to light goes to him.

I would like to thank *Prof. Dr. Wafaa Abdel Aal*, Professor of Pathology, National Research Center and all members of the Pathology Department at the National Research Center, for their encouragement and sincere support.



TABLE OF CONTENTS

INTRODUCTION	1
AIM OF THE WORK	4
MUCOSA-ASSOCIATED LYMPHOID TISSUE (MALT) CONCEPT	5
EXTRANODAL LYMPHOMA	9
GENERAL FEATURES SITES OF ORIGIN CLASSIFICATION OF LYMPHOMA	10
GASTRITIS AND ITS CLINICAL IMPLICATIONS	21
NORMAL ANATOMY AND HISTOLOGY OF THE STOMACHGASTRITISTYPES OF GASTRITIS AND GASTROPATHY	23
GASTROINTESTINAL LYMPHOMA AND MUCOSA ASSOCIATED LYMPHOID TISSUE (MALT)	33
IMMUNOPHENOTYPING IN MUCOSA-ASSOCIATED LYMPHOID TISSUE (MALT)	52
MATERIAL AND METHODS	67
I – Material II – Methods	
RESULTS	
DISCUSSION	103
SUMMARY	114
CONCLUSION AND RECOMMENDATIONS	116
REFERENCES	110



LIST OF FIGURES

Fig. 1: A normal immunoglobulin consists of a heavy chain and a light chain 5
Fig. 2: A Fab fragment is a piece of a particular antibody
Fig. 3: Immunophenotyping technique
Fig. 4: Immunoperoxidase light chain staining
Fig. 5: The equivalent study done by flow cytometry.
Fig. 6: A photomicrograph of a section of gastritis showing gastric crypts overlyin
gastric glands with mild inflammatory reaction. (H&E X 60)
Fig. 7: A photomicrograph of a section of antral mucosa. Showing atrophic change
with elongated gastric crypts, moderately edematous lamina propria and moderat
lymphoplasmacytic infiltrate. (H&E X 100)
Figs. 8: A photomicrograph of a section of chronic gastritis showing gastric crypts wit
mucous secreting lining and scattered curved bacilli (Helicobacter pylori) relate
to surface epithelium. (H&E X 400)9
Figs. 9: The same previous photomicrograph (H&E X 1000[oil immersion]
Fig. 10: A photomicrograph of a section of chronic gastritis showing gastric mucos
with incomplete intestinal metaplasia with moderate lymphocytic infiltrates in th
lamina propria. (H&E X 150)
Fig. 11: A photomicrograph of a section of chronic gastritis showing intestina
metaplasia, marked lymphocytic infiltration and lymphatic follicle. (H&E stain 2
60)99
Fig. 12: A photomicrograph of a section of chronic gastritis showing kappa positivity i
the lymphocytic infiltrates in the lamina propria (polyclonal), with no light chair
restriction. (Kappa X 150)
Fig. 13: A photomicrograph of a section of chronic gastritis showing lambda positivit
in the lymphocytic infiltrates in the lamina propria (polyclonal), with no light
chain restriction. (Lambda X 150)
Fig. 14: A photomicrograph of a section of previously diagnosed chronic gastritis wit
moderate lymphocytic infiltrates showing evidence of light chain restriction
diagnosed as MALT lymphoma after immunohistochemistry (positive kappa
(Kappa X 300)
lambda) (Lambda X 150)
Fig. 16: A photomicrograph of a section of erosive gastritis showing marked dens
lymphocytic infiltrates with superficial erosion. (H&E X 60)
Fig. 17: A photomicrograph of a section of erosive gastritis showing partial ulceration
moderate number of polymorphs infiltrating the epithelium with moderate dens
lymphocytic infiltration and partial effacement of glandular tissue. (H&EX 300
9:
Fig. 18: A photomicrograph of a section of a previously diagnosed erosive gastritis with
dense lymphocytic infiltrates showing light chain restriction, diagnosed as MAL
lymphoma after immunohistochemistry (positive lambda). (Lambda X 150) 90
, di

Fig. 19:	The same case as above showing evidence of light chain restriction (negative appa). (Kappa X 150)
Fig. 20:	A photomicrograph of a section of MALT lymphoma showing widely spaced ini partially devoid of mucous secreting activity with dense infiltration of the
Fig. 21: de	mina propria by lymphocytes. (H&E X 150)
Fig. 22:	: A photomicrograph of a section of MALT lymphoma showing light chain striction (positive lambda). (Lambda X 150)
Fig. 23:	: A photomicrograph of a section of MALT lymphoma showing light chain striction (negative kappa). (Kappa X 150)
ac	A photomicrograph of a section of MALT lymphoma showing few scattered sini, slightly distorted, significantly infiltrated by lymphocytes with dense mphocytic infiltration in the surrounding lamina propria. (H&E stain X 150) 99
in	A photomicrograph of a section of MALT lymphoma showing marked filtration by CCL cells showing marked aggression against glandular pithelium with well formed lympho-epithelial lesions. (H&E X 300)
ге	: A photomicrograph of a section of MALT lymphoma showing light chain striction (positive lambda). (Lambda X 300)
re	: A photomicrograph of a section of MALT lymphoma showing light chain striction (negative kappa). (Kappa X 300)
de de 15	A Photomicrograph of a section of gastric MALT lymphoma showing marked estruction of glandular element with infiltration by CCL cells together with ense lymphocytic infiltrates with well formed lymphatic follicles. (H&E stain X 50)
wi (H	: A photomicrograph of a section of gastric MALT lymphoma showing areas ith dense lymphocytic infiltration with scattered well formed lymphoid follicles. 101
ly	: A photomicrograph of a section of MALT lymphoma showing dense emphocytic infiltration with light chain restriction (positive kappa). (Kappa X 50)
Fig 31:	: A photomicrograph of a section of MALT lymphoma showing dense rmphocytic infiltration with light chain restriction (negative lambda). (Lambda 150)

.

r

.

LIST OF TABLES AND GRAPHS

Table 1 : Revised European - American Lymphoma Classification
Table 2: WHO Classification of lymphoid tumors
Table 3: Principal types and main associations of gastritis and gastropathy
Table 4 : Gastrointestinal lymphoma staging
Table 5 : Classification of Primary gastrointestinal lymphoma
Table 6: The Male/Female ratio and their Mean Age
Table 7: Common symptoms in MALT lymphoma and gastritis groups
Table 8: Types and number of lesion with correlation to Sex
Table 9: Helicobacter pylori organism relevance in H&E results according to sex and
total number of patients
Graph 1: Helicobacter pylori organism relevance according to sex and to total number
of patients
Table 10: Intestinal metaplasia relevance in H&E results according to sex and total
number of patients
Graph 2: Intestinal metaplasia relevance according to sex and total number of patients
Table 11: The Neutrophilic activity and the density of the lymphoid population
relevance according to the number of patients in MALT lymphoma and gastritis
groups
Table 12: Number of patients in each group showing H. pylori organism, Intestinal
metaplasia, Neutrophilic activity and Intense density of lymphoid population 83
Graph 3: Number of patients in each group showing H. pylon organism, Intestinal
metaplasia, Neutrophilic activity and Intense density of lymphoid population 83
Table 13 : Kappa & Lambda immunostaining results
Graph 4 : The Kappa and Lambda light chains immunostaining results
Table 14: Types and number of lesion with correlation to Sex (Kappa & Lambda
immunostaining results)
Graph 5: Diagnosis with H&E vs. Kappa and Lambda Immunostaining

