



شبكة المعلومات الجامعية

شبكة المعلومات الجامعية
@ ASUNET



شبكة المعلومات الجامعية
@ ASUNET



شبكة المعلومات الجامعية



شبكة المعلومات الجامعية

التوثيق الالكتروني والميكرو فيلم

جامعة عين شمس

التوثيق الالكتروني والميكرو فيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
على هذه الأفلام قد أعدت دون أية تغيرات



يجب أن


تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15 – 20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of
15 – 25c and relative humidity 20-40 %



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



شبكة المعلومات الجامعية



بالرسالة صفحات

لم ترد بالأصل

ETIOLOGICAL RESEARCH OF NEPHROPATHIES EPIDEMIOLOGICAL APPROACH

Thesis
Submitted For Partial Fulfillment
Of The Master Degree In Internal Medicine

By

Mohamed Saleh Mokhtar
M. B., B. Ch.

Faculty Of Medicine – Cairo University

Under Supervision Of

Prof. Dr. Magdy Mohamed Soliman

Professor of internal medicine
Cairo University

Prof. Dr. Mohamed Hany Hafez

Professor of internal medicine
Cairo University

**Faculty Of Medicine
Cairo University
2000**

B
✓Y01

باللغة العربية :- بمش فیداحس مع الحبيب أمرا هذا الكس

باللغة الانجليزية :-

١ - الأستاذ الدكتور / محمد بن عبد الله بن سليمان
٢ - الأستاذ الدكتور / محمد بن عبد الله بن سليمان
٣ - الأستاذ الدكتور / محمد بن عبد الله بن سليمان

واستهل الباحث بحرث بلود الرسالة والنتائج التي توصل اليها وكذلك الاسس العلمية التي قام عليها البحث هناك بعدية مع الاذراع التي تؤمن اني انشغل الكون الخرسيم، معصم صندور
فما ينسجى الكلى نصير والمصير الخضر اذ ان لمعصية حامد تؤخر على افعلى وديان ا أجزاء جسم
لزيادة أسباب النشل اللؤلؤ الزمرد، فحقنا بإجراء هذا البحث على عينه مع كاي ج. م. في سكون
وعشر معصم بالنشل اللؤلؤ الزمرد، معا حسد على لست صفا و الدومت من مرالز شرفله لعل
الكل على رفته هذا البحث حاد لنا انه قد حله على حله بغير أسباب النشل اللؤلؤ الخرسيم، لسم، على
و د هجنا انه لسم لسم بالنشل اللؤلؤ الخرسيم على لرسيم هجنا :-

- ١- وثائق لبيات العدل المزمع (١٩٩٢٪)
٢- إيفاد قسوس البصر (١٩٥٥٪)
٣- إيفاد قسوس البصر المزمع (١٩٥١٪)
٤- ضوابط العدل المزمع (١٩٨٩٪)
٥- العدل البصر (١٩٨٨٪)
٦- الفصل العدل المزمع (١٩٨٧٪)
٧- مناقشة السادة أعضاء اللجنة الحكم كل على حده في جميع أبواب وتبويب الرسالة ونتائجها العلمية ومصادرها العلمية.
- ٧- تبليغ العدل (١٩٨٤٪)
٨- العدل العدل المزمع (١٩٨٤٪)
٩- العدل (١٩٨٩٪)
١٠- العدل العدل المزمع (١٩٨٤٪)
١١- العدل العدل المزمع (١٩٨٤٪)
١٢- العدل العدل المزمع (١٩٨٤٪)
١٣- العدل العدل المزمع (١٩٨٤٪)
١٤- العدل العدل المزمع (١٩٨٤٪)
١٥- العدل العدل المزمع (١٩٨٤٪)
١٦- العدل العدل المزمع (١٩٨٤٪)

وقد رت اللجنة : " قبول الرسالة "

استاذ دكتور

کے لیے جو زر

Acknowledgment

* I dedicate this work to the soul of our late Professor, Prof. El-Husseney El-Ghonamy who guided me in the plan & strategy this study.

* I would like to express my deepest gratitude and appreciation to my **Prof.Dr. Magdy Mohamed Soliman** Professor of internal medicine, Cairo University, for his continuous guidance, expert advice and valuable suggestions that greatly enriched this work.

* I'm deeply grateful to the kindness of **Dr. Mohamed Hany Hafez** Professor of internal medicine, Cairo University, for his keen support, sincere help and valuable efforts throughout this study.

Last but not least, I would like to thank all the physicians as well as the nurses and the patients who agreed willingly to participate in that study.

M. Saleh
2000

Abstract

There are many disorders which lead to CRF, these disorders may be specific for the kidney itself or systemic disorders affecting many systems including the kidneys. In a trial to find the common leading causes of CRF in Egypt, this study was done, which included 710 patients with CRF on regular hemodialysis from various dialysis centers in Cairo & Giza. Further more we tried to find the relation between causes of CRF and age, sex and residence. So we found that the leading causes of CRF were respectively: 1. Chronic G.N (22.2 %) 2. Obstructive uropathy (15.5%) 3. Chronic pyelonephritis (15.1 %) 4. Hypertensive nephrosclerosis (13.9 %) 5. Diabetic nephropathy (10.8 %) 6. Unknown causes (9.7 %)

7. Polycystic kidney (3.7 %) 8. Drug nephropathy (3.2%) 9. Gouty nephritis (2.9 %) 10. Lupus nephritis (1.4 %) 11. Toxemia of pregnancy (1.5 %) 12. Primary amyloidosis (0.6 %).

- As regard sex distribution, this study included 710 patients 461 (64.9%) male and 249 (35.1%) females.

- Also we found that the median age of ESRD patients under dialysis treatment ranges between 30-60 years.

Finally the relation between patients and residence was (77.02%) of the patients were living in urban areas and the other (22.98%) were living in rural areas.

Key words

Chronic renal failure - Diabetic nephropathy - Hypertensive nephrosclerosis - Chronic G.N

Table of Contents

Introduction and Aim of the work	
❖ Introduction	1
❖ Aim of the work	2
Review of Literature	
❖ Chronic Renal Failure.....	6
❖ Epidemiology of Chronic Renal Failure	9
❖ Etiology of Chronic Renal Failure in Egypt	22
❖ Etiology of Chronic Renal Failure in other parts in the world	28
Patients and Methods	57
Results	58
Discussion	87
Summary	91
Conclusion and Recommendations	92
References	93
Arabic Summary	

List of Table

Table 1	Classification of CRF	7
Table 2	Results of the study done at 1987 to determine the etiology of CRF in Egypt	24
Table 3	Results of the study done at 1991 to CRF in Egypt	25
Table 4	Results of the study done at 1991 to CRF in Egypt	26
Table 5	Results of the study done at 1996 to CRF in Egypt	27
Table 6	Frequency of patients according to cause of CRF at 1998	29
Table 7	Incidence and prevalence of treated ESRD patients in USA, Canada, Europe and Several Western European Countries (1990-1991)	32
Table 8	The acceptance rates for ESRD therapy in various countries, data from 1992, with some data from 1991 and in some instance updated to 1993	34
Table 9	Etiology of CRF in USA	37
Table 10	Etiology of CRF in Australia	40
Table 11	Etiology of CRF in Europe	42
Table 12	Etiology of CRF in Spain, Poland, Bulgaria and Finland	43
Table 13	Etiology of CRF in Romania	44
Table 14	Etiology of CRF in Republic of Slovenia	45
Table 15	Etiology of CRF in Japan	46
Table 16	Etiology of CRF in China	47
Table 17	Etiology of CRF in South Korea	48
Table 18	Etiology of CRF in India	49
Table 19	Etiology of CRF in Saudi Arabia	51

Table 20	Etiology of CRF in Kuwait	53
Table 21	Etiology of CRF in Sudan	54
Table 22	Etiology of CRF in Nigeria	55
Table 23	Etiology of CRF in South Africa	56

Table of Results

Table 1	Distribution of cases by gender	63
Table 2	Distribution of cases by age	65
Table 3	Geographical Distribution of cases	67
Table 4	Distribution of cases by gender and past history of dysuria	69
Table 5	Distribution of cases by gender and past history of Renal Colic	71
Table 6	Distribution of cases by gender and past history of fever of possible renal origin	73
Table 7	Distribution of cases by gender and past history of schistosomiasis (Urinary)	75
Table 8	Distribution of cases by gender and past history of Urinary tract Infection (proved by lab)	77
Table 9	Distribution of cases by gender and past history Hypertension	79
Table 10	Distribution of cases by gender and past history of Diabetes mellitus	81
Table 11	Distribution of cases by gender and past history of gout before ESRD	83
Table 12	Distribution of cases by gender and Ultrasonic finding	85
Table 13	Distribution of cases according to original disease and etiology of C.R.F.	87
Table 14	Distribution of cases by renal biopsy results	92

List of Figures

Fig 1	Annual report in Egypt, 1997	30
Fig 2	Annual report in Egypt, 1998	31
Fig 3	Age of start of renal replacement therapy in different countries	35
Fig 4	Incidence rates for newly treated ESRD patients in the USA from 1984 to 1992 (solid line) forecast to the year 2000 (dotted line)	38
Fig 5	Incidence rate for newly treated ESRD patients in Europe	41

Fig of Results

Fig 1	Distribution of cases by gender	64
Fig 2	Distribution of cases by age	66
Fig 3	Geographical Distribution of cases	68
Fig 4	Distribution of cases by gender and past history of dysuria	70
Fig 5	Distribution of cases by gender and past history of Renal Colic	72
Fig 6	Distribution of cases by gender and past history of fever of possible renal origin	74
Fig 7	Distribution of cases by gender and past history of schistosomiasis (Urinary)	76
Fig 8	Distribution of cases by gender and past history of Urinary tract Infection (proved by lab)	78
Fig 9	Distribution of cases by gender and past history Hypertension	80
Fig 10	Distribution of cases by gender and past history of Diabetes mellitus	82
Fig 11	Distribution of cases by gender and past history of gout before ESRD	84
Fig 12	Distribution of cases by gender and Ultrasonic finding	86
Fig 13	Distribution of cases according to original disease and etiology of C.R.F. (In Male)	88
Fig 14	Distribution of cases according to original disease and etiology of C.R.F. (In Female)	89
Fig 15	Distribution of cases according to original disease and etiology of C.R.F. (In Total Cases No.)	90
Fig 16	Distribution of cases according to original disease and etiology of C.R.F. (In Total Cases %)	91