



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ



شبكة المعلومات الجامعية
@ ASUNET



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الالكتروني والميكرو فيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأفلام قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيدا عن الغبار

في درجة حرارة من ١٥-٢٥ مئوية ورطوبة نسبية من ٢٠-٤٠%

To be Kept away from Dust in Dry Cool place of
15-25- c and relative humidity 20-40%

بعض الوثائق الأصلية تالفة

بالرسالة صفحات لم ترد بالاصل

SEROEPIDEMIOLOGY OF HBV AND HCV

٢٤٧٨
٢٥٩

AMONG MEDICAL STUDENTS

CAIRO UNIVERSITY

616,3623

Thesis

Submitted for Fulfillment of the Master
Degree in Tropical Medicine

BY

Zeinab Zakaria Mohamed
(M.B.B.Ch.)

Under supervision of

DR. SOHEIR ZAKARIA MOHAMED EISSA

Professor of Tropical Medicine
Faculty of Medicine
Cairo University

DR. HANAN ABD EL HALEEM MARZOUK

Assistant Professor of Tropical Medicine
Faculty of Medicine
Cairo University

DR. OLFAT GAMIL SHAKER

Professor of Medical Biochemistry
Faculty of Medicine
Cairo University

Cairo University
2005

اجتماع لجنة الحكم على الرسالة المقدمة من
الطبيب / زيت رزاق محمد صالح الرز
توطئة للحصول على درجة الماجستير / الدكتوراه
في طب المناظير الجراحية

Seroepidemiology of HBV and HCV among
medical students Cairo University

«مَنْ كَانَ فِي شَيْءٍ مِنَ الْإِسْلَامِ

للرسالة المذكورة أعلاه على النحو التالي :-

١. انذار السيد محمد عيسى
٢. ١.٠٠ هياكلية السيد محمد عيسى
٣. ١.٠٠ هياكلية السيد محمد عيسى

ممتحن داخلي

ممتحن خارجي

ط. محمد عيسى

قرار اللجنة :

~~6/12/20~~

Curves of ρ

W. R. S. P.

455

ABSTRACT

The aim of this study was to determine the seroprevalence of hepatitis B virus and hepatitis C virus among medical students. The results of this study revealed that the overall infection rate for hepatitis B was (14%), with the carrier rate was (1.4%). Eighty nine (25.4%) gave history of full dose HBV vaccine and (82.8%) of the non vaccinated group had no markers for hepatitis B and are at risk of infection. Anti-HCV was found in (2.3%). Exposure to blood/blood products, mucocutaneous exposure and contact to jaundiced patients were the most common risk factors in our studied group. The overall frequency of anti-HAV IgG was (98.3%).

Key Words: Hepatitis A virus - Hepatitis B virus - Hepatitis C virus-
Health care workers.

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

قَالُوا سُبْحَانَكَ لَا عِلْمَ لَنَا إِلَّا مَا عَلَّمْتَنَا

إِنَّكَ أَنْتَ الْعَلِيمُ الْحَكِيمُ

صدق الله العظيم

سورة البقرة

آية (٣٢)

100
100

Acknowledgement

I would like to start by thanking "God" for his help during this work as a little part of his generous help through out my life.

I wish to express my sincere gratitude to Dr. Sohir Zakaria Mohamed Eissa, Professor of Tropical Medicine, Faculty of Medicine, Cairo University. She patiently gave me much of her time, experience, knowledge, directions, supervision and support that can not be expressed by words. From her, I learnt the way of scientific research.

Very great appreciation and thanks are for Dr. Hanan Abdel Haleem Marzouk, Assistant Professor of Tropical Medicine, Faculty of Medicine, Cairo University, for her continuous advice all through this work.

I am so grateful to Dr. Olfat Gamil Shaker Professor of Medical Biochemistry, Faculty of Medicine, Cairo University who gave me much of her support.

Words stand short to express my deep appreciation for the staff members of Tropical Medicine Department, Cairo University for their continuous support.

I am much grateful to Dr. Wafaa El Akel for her great support.

My deep thanks to all my colleagues, the residents of Tropical Medicine Department, Cairo University, for their great support and help.

Thanks for my family who was of great help, gave me lots of guidance and support.

65.