



Transcranial US and Color Doppler studies in Neonatal Hypoxic-Ischemic Brain Insults

**Protocol of an Essay
Submitted for Partial Fulfillment
of the Master Degree in Radiodiagnosis**

Presented by

**Mahmoud Raafat Yehia Hassan
M.B.B.CH
Faculty of Medicine - Ain Shams University**

Supervisors

**Prof.Dr. Hossam Abd-El-Kader Morsy
Professor of Radiodiagnosis
Faculty of Medicine
Ain Shams University**

**Dr. Togan Taha
Lecturer of Radiodiagnosis
Faculty of Medicine
Ain Shams University**

**Ain Shams University
2014**

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

{ قالوا سبحانك لا علم لنا إلا ما علمتنا

انك أنت العليم الحكيم }

صَدَقَ اللَّهُ الْعَظِيمُ

List of Figures

Fig.No.	Title	Pages
1.1	Coronal Plane–Frontal Lobes. This anatomic plane is just anterior to the frontal horns of the lateral ventricles. Normal periventricular echogenicity is evident centrally in each frontal lobe (straight arrow). The curved arrow indicates the midline sagittal fissure. o, orbit	9
1.2	Coronal Plane–Frontal Horns. Images demonstrate the full field of view (A) and a coned down view (B) of the frontal horns (shortest arrow) in two different infants.	10
1.3	Coronal Plane–Third Ventricles. The echogenic choroid plexus (large arrow) is visible lying along the roof of the third ventricle just inferior to the cavum septum pellucidum. This landmark indicates the plane of section is posterior to the foramen of Monro. Echogenic choroid plexus may be seen normally overlying the caudate nucleus in the lateral ventricles in this plane of section. The third ventricle is usually slit-like and is not discretely visualized in coronal plane. Also seen on this image are a portion of the brainstem (b), the cerebellar hemisphere (h), the cisterna magna (small arrow), and the temporal lobes (t)	12
1.4	Coronal Plane–Lateral Ventricles. A. Posteriorly angled coronal view shows the prominent glomus (arrow) of the choroid plexus in the atrium of the lateral ventricles. When this scan was obtained, the infant was lying with the left side of his head down. The choroid plexus lies dependently against the left lateral wall of the ventricle. B. Image angled further posteriorly shows the occipital horns of the lateral ventricles (small arrow). The size of the occipital horns is commonly asymmetric. Note the prominent but symmetric echogenicity of the periventricular white matter tracts (large arrow).	13
1.5	Coronal Plane–Periventricular White Matter. With the transducer held in coronal orientation at the anterior fontanel, this image is obtained by angling the transducer far posteriorly approaching an axial plane. The prominent echogenic periventricular white matter (arrow) is demonstrated superior to the lateral ventricles. The normal echogenicity is symmetric. Asymmetry of the periventricular echogenicity suggests the early hemorrhagic stage of periventricular leukomalacia	14

Fig.No	Title	Pages
1.6	Sagittal Plane–Midline. Images of a premature (A) and a full-term (B) newborn infant demonstrate the important normal anatomic landmarks. The midline cystic cavum septum pellucidum (long arrow) and cavum vergae are prominent in the premature infant but are smaller and limited to only the cavum septum pellucidum in the full-term infant. Note the increased prominence of the gyri and sulci in the full-term infant compared to the relatively featureless brain of the premature. The corpus callosum (fat arrow) forms a curving hypoechoic band just above the cavum septum pellucidum. The choroid plexus (short curved arrow) in the roof of the third ventricle forms an echogenic band just below the cavum septum pellucidum. The vermis (v) of the cerebellum is a round echogenic mass in the posterior fossa. The fourth ventricle (tiny arrow) is seen as a triangular lucency indenting the vermis. Anterior to the vermis is the hypoechoic brainstem (b). The pons (p) is echogenic compared to the remainder of the brainstem. A normal fluid-filled cisterna magna (long curved arrow) should always be present. Absence of the cisterna magna is indicative of a Chiari defect	15
1.7	Sagittal Plane–Lateral Ventricle. Laterally angled, sagittally oriented image demonstrates the rounded mass of the thalamus (T) and the tongue-shaped structure of the caudate nucleus (c). The slight indentation between these two structures is the caudothalamic groove (straight arrow). The caudothalamic groove is the sagittal plane marker of the location of the foramen of Monro. Echogenic mass replacing the tip of the tongue of the caudate nucleus anterior to the caudothalamic groove is indicative of germinal matrix hemorrhage. The echogenic glomus of the choroid plexus (curved arrow) is seen in the atrium of the lateral ventricle. Echogenic choroid plexus extends anteriorly and inferiorly into the temporal horn.	17
1.8	Sagittal Plane–Periventricular White Matter. Sagittally oriented view angled further laterally demonstrates the echogenic periventricular white matter (arrow). This view is utilized to detect periventricular leukomalacia. The echogenicity of the right side should be compared to the echogenicity of the left side.	19
1.9	Axial Plane–Third Ventricle. An axial plane view obtained by imaging through the thin squamous portion of the temporal bone is optimal for demonstrating the third ventricle (straight arrow). The normal third ventricle is a slit-like structure between the two walnut-shaped thalami (T). The brain-cranium interface (curved arrow) should be noted to detect extra-axial fluid collections	20

Fig.No.	Title	Pages
1.10	Axial Plane–Suprasellar Cistern. A. Axial plane view obtained more inferiorly demonstrates the suprasellar cistern as a 5-pointed-star-shaped echogenic structure (short arrow) surrounding the hypothalamus (long arrow). Posterior to the suprasellar cistern is the heart-shaped structure of the cerebral peduncles (p). The echogenic spot (curved arrow) at the apex of the “heart” is the cerebral aqueduct (of Sylvius). B. With color Doppler (shown here in gray scale), the circle of Willis is clearly visible in the suprasellar cistern	21
1.11	Coronal image of a 24 week gestation preterm infant. Note the lack of cerebral sulci. Only the Sylvian Fissures are clearly visible.	22
1.12	Sagittal images of the (a) right and (b) left lateral ventricles demonstrating asymmetry in the choroid plexus (arrows) with the right being larger and more lobulated than the left. Both are normal	23
1.13	Coronal image in a premature infant (note the lack of sulci) demonstrating hyperechoic pseudolesions in the periventricular white matter consistent with normal “periventricular halo”. This should not be confused with the more echogenic, less homogeneous findings of periventricular leukomalacia.	24
1.14	Calcar avis. US image obtained through the posterior fontanelle shows a calcar avis (®) with a central echogenic sulcus (®) as well as an occipital horn (®) and a normal echogenic choroids glomus G1® 1	25
1.15	Lobular choroid plexus. (a) Sagittal US image of the lateral ventricle through the anterior fontanelle shows a large germinal matrix hemorrhage (GMH[®]) with possible intraventricular extension, (b) US image obtained through the posterior fontanelle demonstrates a normal-appearing trigone, occipital horn, and choroid glomus (G[®]) of the right lateral ventricle. (c) US image obtained through the posterior fontanelle shows a lobular left choroid plexus and raises suspicion for intraventricular hemorrhage (?), especially in view of the asymmetry of the lesion relative to the right lateral ventricle. (d) Pulsed color Doppler US image obtained through the posterior fontanelle demonstrates arterial signal arising from the lobular left choroid plexus, a finding that helps exclude DVT. These findings remained stable on multiple follow-up examinations.	26

Fig.No.	Title	Pages
1.16	Normal foramen of Magendie in a 27-week-old infant.(a) Sagittal midline US image obtained through the anterior fontanelle shows a normal fourth ventricle (®), inferior vermis (®), and cisterna magna (®). (b) US image obtained through the mastoid fontanelle demonstrates a normal –appearing cisterna magna (®) and cerebellar hemispheres (®). (c) more caudally angled US image obtained through the mastoid fontanelle shows the normal fourth ventricle (®) and inferior vermis (®). (d) US image obtained through the mastoid fontanelle with further caudal angulation shows a thin communication between the fourth ventricle and the cisterna magna (foramen of Magendie)	28
1.17	Atlas of neuroanatomy and neurophysiology 2002	31
1.18	Normal color flow Doppler sonogram via the anterior fontanelle in a term neonate. (A) Sagittal midline view with infant facing the left demonstrates the ACA coursing over the corpus callosum. Cursor is placed on the pericallosal artery demonstrating a normal RI for a term infant measuring 0.73. (B) Coronal view with cursor placed on the peripheral left MCA demonstrating a normal RI of 0.66.	34
1.19	Normal TCD color Doppler image obtained from the left temporal approach. Labeled vessels include middle cerebral artery (MCA), posterior cerebral artery (PCA), and anterior cerebral artery (ACA). <i>Red</i> and <i>blue</i> indicate arbitrary designation of flow toward and away from the transducer, respectively. <i>Lt</i> left, <i>Rt</i> right	35
2.1	The effect of ischemia on cerebral perfusion, Berger et al 2000	40
2.2	Diagram illustrates the cascade of events contributing to brain injury in hypoxia-ischemia	44
3.1	Drawing illustrates the neonatal skull (lateral view) with the major sutures as well as the posterior (®) and anterior (®) fontanelles. A US transducer is positioned over the mastoid fontanelle (Reprinted, with permission, from reference2)	48
3.2	. van Wezel-Meijler 2007	49
3.3	G. van Wezel-Meijler 2007	50
3.4	Posterior fontelle imaging (a) US image obtained the posterior fontelle with the transducer positioned directly over the fissure (®) demonstrates the body of the right lateral ventricle (®) as well as the occipital horn (®) and a choroid glomus (G[®]). (b) US image obtained with the the transducer positioned slightly cephalad to the calcarine fissure (®) provides an excellent view of the body of the right lateral ventricle (®) as well as the occipital (®) and temporal (®) horns and the choroids glomus (G[®]). Because the patient was supine, the posterior fontanelle was not accessible.	51

Fig.No.	Title	Pages
3.5	G. van Wezel-Meijler 2007	52
3.6	Mastoid fontelle imaging.(a)US image of the midbrain obtained through the mastoid fontanelle shows an echogenic line denoting a collapsed third ventricle and aqueduct (®) as well as the cerebral peduncle (®),thalamus (®), and quadrigeminal plate cistern (Q[®]), which blends into the upper cerebellum. The perimesencephalic cistern (®) and tentorium (®)are also a seen.(b) US image of the poserior fossa obtained through the mastoid fontanelle depicts the fourth ventriclw (®), posterior vermis (®), cisterma manga (®), and cerbellar hemisphere (CH [®]).	53
3.7	Normal coronal gray-scale images of 7-day-old boy with seizures. A , Sonogram through frontal lobes (F) shows orbits (O) and hyper echoic falx cerebri (arrow) located within interhemispheric fissure. B , Sonogram through frontal horns shows corpus callosum, seen as hypo echoic midline structure outlined by echogenic superior and inferior borders (arrow). Frontal horns (F), cavum septum pellucidum (asterisk), globus pallidus (G), putamen (P), caudate nucleus (C), temporal lobes (T), and Sylvian fissure (dotted line) are seen. C , Sonogram at level of cerebral peduncles identifies hyper echoic choroid plexus along roof of third ventricle (straight arrow) and floor of lateral ventricles (curved arrows). Also noted are left cerebral peduncle (CP), thalamus (T), cerebellar hemispheres (Cb), and cisterna magna (arrowhead). D , Sonogram at level of quadrigeminal plate (Q) illustrates temporal lobes (TL), cerebellar hemispheres (Cb), thalamus (T), hippocampus (H), and third ventricle (3). E , Sonogram obtained through hyper echoic choroid plexus (asterisks) within lateral ventricles. Note less echogenic adjacent periventricular white matter (arrows). Periventricular white matter halo should normally be less echogenic than adjacent choroid plexus. F , Sonogram through cerebral convexities reveals normal layers of cortex, which should be seen throughout brain. Well-defined hyper echoic pia (curved arrow) on surface of cortex overlies hypo echoic cortical gray matter (straight arrow), which overlies slightly hyper echoic white matter (arrowhead)	55
3.8	Normal sagittal gray-scale images of 7-day-old boy with seizures (same patient as Fig. 3). A , Sonogram through midline shows corpus callosum (straight arrows), hypo echoic structure bound by echogenic superior and inferior borders, cingulate gyrus (arrowheads), cavum septi pellucidi (asterisk), occipital lobe (O), third ventricle (3) with choroid plexus in its roof (curved arrow), and fourth ventricle (4). Also noted are hypo echoic midbrain (M); Pons (P), which has hyper echoic ventral and hypo echoic dorsal regions; and cerebellar vermis (V). B , Parasagittal sonogram through lateral ventricles shows caudothalamic groove (arrow), separating caudate nucleus (C) from thalamus (T), and choroid plexus along posterior margin of thalamus (dotted line). Note hyper echoic focus adjacent to lateral ventricle trigone that was not seen on coronal view (not shown). This periventricular pseudo lesion (arrowhead) is common artifact on this view. C , Normal peripheral sagittal gray-scale sonogram through Sylvian sulcus (dotted line) reveals temporal lobe (T), frontal lobe (F), and parietal lobe (P). Note normal hypo echoic gray matter (arrow) between superficial hyper echoic pia and slightly hyper echoic white matter.	56

Fig.No.	Title	Pages
3.9	<p>Normal color Doppler Sonography of 8-day-old premature girl being screened for intracranial hemorrhage.</p> <p>A, Sonogram obtained through circle of Willis illustrates right (Rt) and left (Lt) internal carotid arteries (ICA), right and left middle cerebral arteries (MCA), and anterior cerebral (ACA) arteries. Distinguishing between right versus left ACA is often difficult, but usually unnecessary. Thus, they are labeled together.</p> <p>B, Spectral Doppler tracing of circle of Willis obtained via anterior fontanel shows normal arterial flow pattern. in MCA. Note continuous flow above baseline, including rapid systolic upstroke (<i>arrow</i>) followed by gradual decline in flow during diastole (<i>arrowhead</i>). Normal peak systolic velocity (PSV), end-diastolic velocity (EDV), and resistive index (RI) are indicated.</p> <p>C, Sagittal Doppler sonogram of venous system shows patent deep venous system, including vein of Galen (VOG) (<i>arrow</i>). Pericallosal branch of ACA (<i>arrowhead</i>) is noted.</p> <p>D, Sonogram of superficial venous system, including sagittal sinus (<i>arrow</i>), is shown. Observe normal venous spectrum showing continuous undulating venous flow. Incidentally noted is reversal of color map</p>	57
3.10	<p>Normal gray-scale posterior fossa sonogram obtained via mastoid fontanel in 3-day-old infant, on extracorporeal membrane oxygenation therapy screened for intracranial hemorrhage. Cerebellar hemispheres (Cb), fourth ventricle (4), cistern magna (asterisk), quadrigeminal plate (Q), vermis (V), and temporal lobes (T) are shown</p>	59
3.11	<p>Posterior fossa hemorrhages in 2-week-old boy born at 27 weeks' gestational age. A and B, Coronal sonogram via anterior fontanel shows area of hypoechogenicity in right cerebellum (<i>arrow</i>), concerning for hemorrhage. Note developing post hemorrhagic hydrocephalus. When evaluated via mastoid fontanel (B), right cerebellar hemorrhage (<i>arrow</i>) is confirmed</p>	59
3.12	<p>Normal linear Sonographic image in coronal plane in 6-week-old girl with macrocephaly. Color. Doppler linear image of superficial cortex reveals prominence of subarachnoid space. Normal vessels (<i>arrows</i>) are present. (</p>	60
3.13	<p>Hypoxic ischemic brain injury in 2-day-old boy with seizures.</p> <p>A, Coronal sonogram reveals bilateral increased thalamic echogenicity (<i>arrows</i>). MRI (not shown) confirmed restricted diffusion in dorsal putamina and ventrolateral thalami. B, Coronal power Doppler image using linear-array transducer reveals hyperemic flow in lenticulostriate and thalamoperforate arteries of basal ganglia and thalami</p>	62

Fig.No.	Title	Pages
3.14	Right middle cerebral artery stroke in 1-week-old boy. Coronal sonogram shows loss of normal cortex in right cerebrum (<i>arrows</i>). Compare with normal cortex on left	63
3.15	Periventricular leukomalacia in 6 week-old premature girl. Coronal sonogram identifies bilateral periventricular hyper echogenicity and subtle periventricular cystic encephalomalacia (<i>arrows</i>). MRI (not shown) confirmed diffuse abnormal signal in white matter and some periventricular hemorrhage.	63
3.16	Low resistive index (RI) in 2-day-old infant with hypoxic ischemic brain injury. Spectral Doppler tracing of middle cerebral artery (MCA) obtained via temporal fontanel reveals elevated diastolic flow (<i>arrow</i>) and low RI of 0.53 (normal RI, approximately 0.6–0.9). Gray-scale images (not shown) revealed diffuse increased echogenicity suggesting cerebral edema. PSV = peak systolic velocity, EDV = end diastolic velocity.	66
3.17	Elevated resistive index (RI) in 8-day-old preterm infant due to patent ductus arteriosus. Spectral Doppler tracing obtained from middle cerebral artery (MCA) shows elevated RI of 1.34 (normal preterm RI, 0.77 ± 0.15). Note normal rapid systolic upstroke followed by abnormally rapid decline with reversal of flow during diastole (<i>arrow</i>). PSV = peak systolic velocity, EDV = end-diastolic velocity	67
4.1	12 h old girl born at 39 weeks of gestation with a history of perinatal asphyxia after total abruption placenta and the respective Apgar scores 1-2-3 at the age of 1-5-10 minutes with severe HIE who survived with cerebral paralysis, tetraplegia, seizures and bulbar paralysis. (Ilves et al., 2009a). a) Coronal US image shows ill-defined structures including basal ganglia, and slitlike ventricles at the age of 12 h b) Magnified sagittal view using linear transducer shows the accentuation of the grey –white matter differentiation with prominent sulci at the age of 12 h c) Left parasagittal view shows hyperaemia in the region of basal ganglia with power Doppler at the age of 12 h d) Colour Doppler evaluation shows a low cerebral blood flow velocity with normal resistance in the right anterior cerebral artery at the age of 12 h e) Coronal US image shows still ill-defined structures with general increase in echogenicity including basal ganglia, and still slitlike ventricles at the age of 3 days f) Colour Doppler evaluation shows a high cerebral blood flow velocity with low resistance in the left medial cerebral artery at the age of 3 days g) MRI FLAIR image with pathological signal intensity bilaterally in basal ganglia at the age of 8 days h) Coronal US with moderately enlarged side ventricles and ill-defined basal ganglia at the age of 1 month i) CT image with atrophy and increased attenuation in basal ganglia at the age of 3 years.	72

Fig.No.	Title	Pages
4.2	12 h old girl born at 39 weeks of gestation with a history of perinatal asphyxia after total abruption placenta and the respective Apgar scores 1-2-3 at the age of 1-5-10 minutes with severe HIE who survived with cerebral paralysis, tetraplegia, seizures and bulbar paralysis. (Ilves et al., 2009a). a) Coronar US image shows ill-defined structures including basal ganglia, and slitlike ventricles at the age of 12 h b) Magnified sagittal view using linear transducer shows the accentuation of the grey –white matter differentiation with prominent sulci at the age of 12 h c) Left parasagittal view shows hyperaemia in the region of basal ganglia with power Doppler at the age of 12 h ,d) Colour Doppler evaluation shows a low cerebral blood flow velocity with normal resistance in the right anterior cerebral artery at the age of 12 h ,e) Coronar US image shows still ill-defined structureswith general increase in echogenicty including basal ganglia, and still slitlike ventricles at the age of 3 days, f) Colour Doppler evaluation shows a high cerebral blood flow velocitywith low resistance in the left medial cerebral artery at the age of 3 days ,g) MRI FLAIR image with pathological signal intensity bilaterally in basal ganglia at the age of 8 days h) Coronar US with moderately enlarged side ventricles and ill-defined basal ganglia at the age of 1 month, i) CT image with atrophy and increased attenuation in basal ganglia at the age of 3 years	77
4.3.a	Sagital view at lateral ventricle level echogenic material in the thalamo-caudate groove; b) Detail the irregular echogenic deposit is located between the caude nucleus (anterior) and thalamus (posterior).(Diagnostic:small subependumal grade I haemorrhage	83
Fig.3.a	Coronal view at the frontal horns level echogenic material on both supependimal regions, more abundant on the left, b) Sagittal view: oval shaped echogenic deposit in the left thalamo-caudate groove, apparently protruding into the lateral ventricle. Diagnostic: subependimal grade I haemorrhage	83
Fig.3.a)	Coronal view at the frontal horns level echogenic material on both supependimal regions,more abundant on the left; b)sagittal view :oval shaped echogenic deposit in the left thalamo-caudate groove, apparently protruding into the lateral ventricle. Diagnostic: subependimal grade I haemorrhage.	83
4.4	Control view: the thalamic and caudate grooves are filled with hyperechoic material, more abundant on the right side; b) Coronal view at the lateral ventricles body: asymmetric left ventricles: minimal right ventricle dilation.Diagnostic: bilateral subependimal haemorrhage with right interaventricular haemorrhage and minimal ventricular distension-grade H haemorrhage.	84

Fig.No.	Title	Pages
4.5	Coronal view at ventricular body level: both ventricles are distended; on the left side an echogenic dot extends to the frontal horn; b) sagittal view demonstrate oval shaped irregular clot on the left ventricle's body, distinct by echogenicity and location from the choroids plexus which can be seen posterior. Diagnostic: grade III Intraventricular haemorrhage	86
4.6	Grade III intraventricular haemorrhage outcome: ultrasound examination at 5 and 8 weeks postnatal. a) Coronal view: dilated frontal & temporal horns, mild IIIrd ventricle distension; b) Asymmetrical frontal and temporal horns distension. Diagnostic: intraventricular grade III haemorrhage with secondary hydrocephalus	86
4.7	Full term newborn with severe birth hypoxia- examination on day 5. a) Coronal view: both lateral ventricles filled by an echogenic structure, consistent with bilateral intraventricular clots: Hyperechoic periventricular parenchyma, suggestive for venous infarctions; b) Sagittal view at lateral ventricle level: The hyperechoic clot extends to almost entire ventricle. Diagnostic: grade IV intraventricular haemorrhage	88
4.8	The same case as in fig.7 reexamined at 18 days of age. A) sagittal view: pronounced distension of the left lateral ventricle with persistent hyperechoic images inside. B. Coronal view bilateral ventriculomegaly, intraventricular clot on a slow resolution process. Diagnostic grade IV intraventricular haemorrhage post hemorrhagic hydrocephalus	88
4.9	A3 days old newborn, 28 weeks gestational age weight 1900 gr with severe perinatal hypoxia, Apgar score 3/1 min. Head ultrasound on day 7. a) coronal view: both frontal and temporal horns are distended by an echogenic irregular, inhomogeneous structure inside the ventricles. Dilated third ventricle; b) Sagittal view to secondary ventricle level: voluminous clot with irregular borders leading to secondary ventricular distension. Diagnostic grade IV intraventricular haemorrhage. secondary hydrocephalus	88
4.10	Coronal view: distended lateral ventricle with porencephalic cyst formation. Diagnostic: grade IV intraventricular haemorrhage. Post hemorrhagic hydrocephalus, porencephalic cysts	89
4.11	Preterm newborn examined for periventricular hyperechogenicities observed at 48 h of age a) sagittal view day 3 periventricular hyperechogenicities superior to the echogenicity of choroids plexus; b) Examination on day 10: small cyst in the periventricular area, c) coronal view cyst on the external angle left frontal horn, d) sagittal view multiple periventricular cystic cavities. Diagnostic: grade II PVL	95

Fig.No.	Title	Pages
4.12	a)and b) Coronal view: in various cerebral regions-multiple extensive cystic cavities on the fronto occipital regions; c) and d) Sagittal views: multiple cysts with various shapes and dimensions located on both peniventricular area. Diagnostic hypoxic ischemic encephalopathy; grade III PVL	95
4.13	Preterm newborn-36 weeks gestrional age . Apgar score2/lmin 4/5 min examined in the 14h days of life a) and b) coronal view numerous cysts of different shapes and dimensions extending periventricular bilaterally. Marked extraaxial spaces distension with anechoic content, c) and d) sagittal view peniventricular cysts with fronto-paneto-occipital extension. Enlarged subarachnoid space .Note the normal dimensions of the lateral ventricles .Diagnostic: hypoxic encephalopathy ; grade IV PVL	95
5.1	Doppler curve of pericallosal artery: PI – pulsatility index, RI – resistive index, PSV – peak systolic blood flow velocity, EDV – end-diastolic blood flow velocity, MnV – mean blood flow velocity, FlowT – flow time	105
5.2	High RI values in the first day of life, with typical diastolic notch on the spectral curve, followed by low RI characteristic for luxurious perfusion, after 24 hours in the reperfusion period	119
5.3	Serial changes in the mean BFV (TAV) in the BA during the first 6 months of age. Data are represented as mean \pm SD. * $P < .05$, HIE 0 versus HIE 3; ** $P < .0005$, HIE 0 versus HIE 3; § $P < .05$, HIE 1 + 2 versus HIE 3; §§ $P < .0005$, HIE 1 + 2 versus HIE 3; # $P < .05$, HIE 0 versus HIE 1 + 2	120
5.4	Transcranial duplex Doppler ultrasonography demonstrating a low resistance pattern of the middle cerebral artery (MCA) in an asphyxiated infant.	123
5.5	Transcranial suplex Doppler ultrasonography demonstrating a low resistance pattern of the anterior cerebral artery (ACA) in an asphyxiated infant.	124

List of Table

Fig.No.	Title	Pages
4.1	Peri and intraventricular hemorrhage PVH-IVH classification is detailed in	81
5.1	The reference values of Doppler parameters of A1 and A3 segment of anterior cerebral artery during the first year of life	108
5.2	The reference values of Doppler parameters of M1 and M3 segment of right middle cerebral artery during the first year of life	109
5.3	The reference values of Doppler parameters of M1 and M3 segment of left middle cerebral artery during the first year of life	109
5.4	The reference values of resistive index of Doppler curve in the vertical segment of pericallosal artery before the genu corporiscallosi in preterm newborns	109
5.5	Extracranial and intracranial factors that influence Doppler parameters of the cerebral circulation	114
5.6	Cerebral hemodynamic parameters in studied groups in the first 72 h of life	118
5.7	Comparison of daily values of the mean, minimum and maximum RI values in the anterior cerebral (ACA) and middle cerebral artery (MCA) between the study groups	125

Contents

	pages
<u>List of Tables</u>	<u>I</u>
<u>List of Figures</u>	<u>XI</u>
<u>Introduction</u>	<u>1</u>
<u>Aim of Work</u>	<u>5</u>
<u>Sonographic anatomy of infant brain.</u>	<u>6</u>
<u>Pathophysiology of Hypoxic Ischemic Encephalopathy in Neonates.</u>	<u>36</u>
<u>Technique of Transcranial Doppler ultrasound in Neonates.</u>	
<u>Findings of Transcranial Doppler ultrasound in Hypoxic-Ischemic Encephalopathy in Neonates.</u>	
<u>Illustrative Cases</u>	
<u>Summary & Conclusion.</u>	
<u>References.</u>	
<u>Arabic summary.</u>	

References

1. Allison JW, Faddis LA, Kinder DL, Roberson PK, Glasier CM, Seibert JJ. Intracranial resistive index (RI) values in normal term infants during the first day of life. *Pediatr Radiol* 2000; 30:618– 620.
2. Anca IA. Ecografia transfontanelara la nou-nascuti si sugari– abordarea practica. Bucuresti 2007, Editura Medicala,
3. Arabin B , Saling E . Economy circuit of fetal circulation illustrated on the basis of personal quantitative Doppler blood flow parameters]. . *Geburtshilfe Perinatol.* (1987)
4. Arie Franco ,Kristopher Neal Lewis Neonatal cranial ultrasound: current perspectives 2013
5. Babcock, DS. & Ball, W. Jr. Postasphyxial encephalopathy in full-term infants: US diagnosis. *Radiology* 1983; 148: 417–423.
6. Babcock D. The normal, absent, and abnormal corpus callosum: Sonographic findings. *Radiology* 1984;151:449-453
7. Barkovich AJ, ed. *Pediatric neuroimaging*. 3rd ed. New York, NY: Lippincott Williams & Wilkins, 2000; 162–208.