

**MULTI PLANER DETECTOR CT OF EMERGENCY
BILIARY PATHOLOGICAL CONDITION**

EssaY

Submitted for partial fulfillment of Master Degree

In Radiodiagnosis

By

ENAS ELSAID MOHAMED MOHAMED

M.B., B.Ch

Under The Supervision Of

Prof. Dr. MOHAMED ABD AZIZ

Professor of Radiodiagnosis

Faculty of Medicine, Ain Shams University

Dr. REMON ZAHER

Lecturer of Radiodiagnosis

Faculty of Medicine, Ain Shams University

Faculty of Medicine

Ain Shams University

2016



سورة البقرة الآية: ٣٢



Acknowledgments

First and foremost, I feel always indebted to Allah the Most Beneficent and Merciful.

*I wish to express my deepest gratitude and thanks to **Prof. Dr. Mohamed Abd Aziz**, Professor of Radiodiagnosis, Faculty of Medicine, Ain Shams University, for his constructive criticism, unlimited help and giving me the privilege to work under his supervision.*

*My most sincere gratitude is also extended to **Dr. Remon Faris**, Lecturer of Radiodiagnosis, Faculty of Medicine, Ain Shams University, for his enthusiastic help, continuous supervision, guidance and support throughout this work.*

*Last but not least, I can't forget to thank all members of my Family, especially my **Parents**, for pushing me forward in every step in the journey of my life.*

Candidate

 **Enas Elsaid Mohammed**

List of Contents

<i>Subject</i>	<i>Page No.</i>
List of Abbreviations	i
List of Figures	ii
Introduction	16
Aim of the Work.....	3
Chapter (1): Anatomy	4
Chapter (2): Definition and Pathology of Emergency Biliary Pathological Condition.....	27
Chapter (3): Technique of multi detector CT	42
Chapter (4): Role of Multi Detector CT in Diagnosis of Emergency biliary Pathological Condition.....	53
Summary and Conclusion	82
References	85
Arabic Summary	—

List of Abbreviation

CT	CT Computed tomography
MDCT	Multi detector computed tomography
IHBDS	Intra hepatic bile ducts
CHD	Common hepatic duct
CBD	Common bile duct
GB	Gall bladder
MinIP	Minimum intensity projection
MPR	Multiplanar reformation
VR	Volume rendering
PVP	Portal venous phase
HAP	Hepatic arterial phase
CPR	Curved planar reformation
PACS	Picture archiving and communication
CDL	Choledochlithiasis
CC	Cholangiocarcinoma
ERCP	Endoscopic retrograde cholangiopancreatography
MRCP	Magnetic resonance cholangiopancreatography
US	Ultrasound
PTC	Percutaneous trans hepatic cholangiography

List of Figures

<i>Figure No.</i>	<i>Title</i>	<i>Page No.</i>
Figure (1):	over all arrangement of the intrahepatic and extrahepatic biliary tree.....	5
Figure (2):	Segmental liver anatomy based on couinaud classification (segment1-8) (RHV right hepatic vein, LHV: left hepatic vein, MHV: middle hepatic vein, IVC: inferior vena cava.....	6
Figure (3):	Interior of the gallbladder and its cystic duct.....	9
Figure (4):	Drawing illustrates the normal biliary tract. The cystic duct (arrow) connects the gall bladder to the extra hepatic duct	10
Figure (5):	The extra hepatic biliary tract and four portions of the common bile duct	12
Figure (6):	Variations of the pancreatic ducts (a) both ducts open into the duodenum (b)the accessory duct ends blindly in the duodenal wall (c) the main duct is smaller than the accessory duct and they are not connected	13
Figure (7):	Segment VII of the biliary duct originate from the right hepatic duct while segment II is located in the left hepatic lobe.....	14

Figure (8):	Normal intrahepatic duct. Contrast enhanced computed tomography shows non –dilated left intra hepatic bile duct.....	15
Figure (9):	CT scan shows normal common hepatic duct at the confluence of the right hepatic duct with the left hepatic duct	17
Figure (10):	MDCT shows the normal gallbladder located at anterior – inferior edge of the liver. It is divided into three parts: neck, body and infundibulum	18
Figure (11):	Contrast enhanced CT scan reveals normal appearance of gall bladder in coronal plane	19
Figure (12):	Phrygian cap. CT image of the gallbladder fundus shows normal variant Phrygian cap	21
Figure (13):	Normal cystic duct anatomy. Axial CT scan shows the normal cystic duct extending from the gallbladder	22
Figure (14):	Schematic drawing of cystic duct variation.....	23
Figure (15):	Parallel course of the cystic duct. ERCP image obtained in a 57-year-old man shows the long, parallel course of the normal cystic duct	24
Figure (16):	Coronal intravenous contrast-enhanced reformatted image from CT data shows the CBD which is normal caliber and have an imperceptible wall	25

Figure (17):	MDCT shows normal pancreatic duct merges with the duodenum together with the common bile duct.....	26
Figure (18):	Cholesterol stone.....	29
Figure (19):	CT oblique coronal reformation (A) axial images should be complemented by oblique coronal reformation parallel to both hepatic duct (B) the oblique coronal reformatted image displays a longitudinal view of both hepatic ducts and extra hepatic duct together	47
Figure (20):	CT curved planar reformation (A) CPR are obtained by interactively placing a cursor on a stack of axial sections along a course of a specific anatomic structure (B) CPR through main pancreatic duct throughout entire course as well as portions of the pancreas	48
Figure (21):	Minimum intensity projection.....	50
Figure (22):	30 years old woman with acute calculous cholecystitis. Contrast-enhanced CT shows a distended gallbladder (arrowheads) with a slightly thickened wall and subtle regional fat-stranding (asterix). There is an impacted obstructing stone in the neck of the gallbladder.....	54
Figure (23):	43-year-old woman with acute calculous cholecystitis. Contrast-enhanced CT shows a distended gallbladder with a slightly thickened wall and pericholycystic fluid collection.....	55

Figure (24):	Axial nonenhanced CT image in a 70-year-old woman shows intraluminal air in the gallbladder and pericholecystic fluid	55
Figure (25):	Contrast-enhanced CT abdomen confirms a gallbladder distended with gas and fluid, and intramural gas	56
Figure (26):	CT of abdomen axial cut shows Air fluid level in gall bladder	56
Figure (27):	Distended gallbladder with wall thickening (full arrows), pericholecystic fluid (dotted arrow) and surrounding fat stranding (dashed arrows) are signs of an acute inflammatory process.....	57
Figure (28):	Axial contrast-enhanced CT image in a 55-year-old woman shows irregular thickened gallbladder mucosa with discontinuity and sloughed membranes . A common bile duct stent is in place for biliary decompression from known pancreatic cancer	58
Figure (29):	Coronal contrast-enhanced CT image in a 70-year-old woman shows a hydropic gallbladder with dependent stones and sludge , focal disruption of the gall bladder wall , and multiple loculated pericholecystic fluid collections	58
Figure (30):	CT scan upper abdomen showing a large haematoma in the gallbladder fossa, which distends the gallbladder wall causing mass effect	59

- Figure (31):** Contrast CT scan through the gall bladder show streak of contrast escaping into gall bladder lumen and hyper dense contents within gall bladder 59
- Figure (32):** CT demonstrates acute cholecystitis with gallbladder perforation and pericholecystic fluid. Yellow arrows indicate gallbladder wall defects..... 60
- Figure (33):** In CT images of a patient diagnosed with acute emphysematous cholecystitis with gallbladder perforation, yellow arrows indicate free intraluminal air in the gallbladder. Red arrows point to free air and biloma over pericholecystic and perihepatic regions 61
- Figure (34):** Axial contrast-enhanced CT image shows a gas-distended gallbladder lumen with multiple foci of air in the gallbladder wall (arrowheads). Air and fluid in the porta hepatis (black arrow) and a rim-enhancing air-containing abscess (white arrow) anterior to the left lobe of the liver indicate perforation..... 61
- Figure (35):** Axial contrast-enhanced CT image at a more caudal level shows circumferential gallbladder wall thickening and pericholecystic inflammatory changes with reactive mural thickening of the ascending and transverse colon. An additional rim-enhancing abscess with surrounding inflammatory change is also seen..... 62

- Figure (36):** Two axial images from a contrast-enhanced CT of the abdomen show air in the gall bladder and air in the common bile duct representing pneumobilia, the gallstone in the small bowel lumen and dilated and fluid-filled loops of small bowel from small bowel obstruction all of the components of a gallstone ileus 63
- Figure (37):** CT confirms the findings of gall stone ileus: four axial images from a contrast-enhanced CT show gallstone in the small bowel lumen and dilated and fluid-filled loops of small bowel from small bowel obstruction..... 63
- Figure (38):** Mirizzi syndrome in an 80-year-old woman with right upper quadrant pain (a) Axial nonenhanced CT image shows intrahepatic biliary ductal dilatation 64
- Figure (39):** Coronal nonenhanced CT image shows several hyper-attenuating gallstones and an impacted stone in the gallbladder neck. Dilatation of the common hepatic duct is seen proximal to the stone. The common bile duct distal to the stone is normal in caliber 64
- Figure (40):** Computed tomographic scan showing a calcified stone causing bile duct obstruction with the proximal end of the stent just below the stone. Residual contrast from the ERCP is present in the gallbladder..... 65

- Figure (41):** Variable CT appearance of gall stones a) large gall stone with calcified peripheral rim(black arrow) with completely collapsed gall bladder (white arrow).b)gall stones appear as layer of calcified microlith with high attenuation foci layering dependently in gall bladder from small, calcified stones (arrow). C) multiple small, rim- like calcified stones in gallbladder (arrow) d)multiple gas containing stones are identified as low attenuation foci within gall bladder 66
- Figure (42):** Distal common bile duct stone. a) Noncontract scan demonstrating high attenuation stone in distal common bile duct .b) following contrast enhancement stone appears to be of soft tissue attenuation relative to pancreas, with high attenuation peripheral rim in distal common bile duct..... 67
- Figure (43):** Axial contrast-enhanced arterial phase CT image in a 62-year-old woman shows intrahepatic biliary ductal dilatation and wedge-shaped hepatic parenchymal enhancement..... 68
- Figure (44):** This is the computed tomography (CT) of a 45-year-old male with enhanced CT abdomen show necrosis of the head and body of the pancreas. The gallstone in the gallbladder is also noted..... 69
- Figure (45):** Acute biliary pancreatitis with a thickened pancreas and an effusion

	around the pancreatic tail and around spleen.....	69
Figure (46):	CT scan demonstrating a single large liver abscess	71
Figure (47):	CT scan of the abdomen with contrast showing a large, loculated liver abscess measuring 10 cm	71
Figure (48):	Contrast-enhanced computed tomography scan of the abdomen showing a huge gas-forming abscess with an air-fluid level in the right lobe of the liver.....	71
Figure (49):	Post contrast CT image demonstrates avidly contrast-enhancing polypoid gall bladder lesion from renal cell carcinoma.....	72
Figure (50):	Carcinoma of pancreatic head (a) peri-ampullary tumor with associated with dilatation of common bile duct and pancreatic duct (double duct sign, arrow head) (b) tumor of pancreatic head with local lymphaginitis in the mesenteric fat and a small lymph node.....	73
Figure (51):	intra hepatic mass forming cholangio carcinoma (a-c)ct scan during arterial, portal, delayed phase show hypo attenuating mass with early peripheral enhancement and gradual and centeripetal progression in right lobe of the liver. Satellite nodule or intra hepatic metastasis noted	74

Figure (52):	Common bile duct cholangiocarcinoma in a 72-year-old man	74
Figure (53):	Abdominal CT showed a 5x8x7cm lobulated fluid collection	75
Figure (54):	Coronal contrast-enhanced CT image shows a large amount of ascites (*) scattered throughout the abdomen. Cholecystectomy clips are seen in the gallbladder fossa.....	76
Figure (55):	Image from technetium 99m–mebrofenin cholescintigraphy shows intense radiotracer activity in the expected position of the gallbladder fossa with activity tracking along the right hepatic lobe and superiorly toward the dome, findings compatible with an active biliary leak. Radiotracer activity is also seen in the small bowel, indicating patency of the common bile duct.....	76
Figure (56):	Migrated cholecystostomy tube in a 74-year-old man with acute cholecystitis (a) Coronal contrast-enhanced CT image shows a loculated air-containing perihepatic fluid collection (*). A cholecystostomy tube (arrow) is seen in the hydropic gallbladder.....	77
Figure (57):	Coronal contrast-enhanced CT image with wide window width and high level settings shows the radiopaque marker of the cholecystostomy tube outside the gallbladder and in the peritoneum. The radiopaque marker denotes the position	

of the most proximal drainage side hole as outside the gallbladder, allowing bile spillage into the peritoneal cavity 77

Figure (58): Occluded biliary stent in a 74-year-old woman with pancreatic cancer (a) Coronal contrast-enhanced CT image shows a malfunctioning common bile duct stent with intra- and extrahepatic biliary ductal dilatation and a hydropic gallbladder (*). No pneumobilia is seen 78

Figure 59): Coronal contrast-enhanced CT image obtained 6 weeks earlier shows a normal-caliber biliary system, a normal gallbladder, and pneumobilia, findings suggestive of patency of the common bile duct stent 79

Figure (60): Postcholecystectomy clip migration in a 57-year-old man (a) Axial contrast-enhanced CT image shows a cholecystectomy clip in the gallbladder fossa and dilatation of the commonbile duct.... 79

Figure (61): Early hepatic artery thrombosis in an 18-month-old girl 3 weeks after liver transplantation (a) Axial contrast-enhanced CT image shows marked branched biliary ductal dilatation (*) adjacent to the portal vein branches in the hepatic hilum, a finding compatible with biliary necrosis 80

Figure (62): Coronal arterial phase maximum intensity projection CT image shows absence of the proper hepatic artery (circle) and lack of hepatic arterial