

Tranexmic Acid and reducing blood loss after Vaginal Delivery: A randomized controlled trial

Thesis

Submitted for Partial Fulfillment of Master Degree
In Obstetrics and Gynecology

By

Basma Yehia Saeed Yehia

M.B.B.Ch

*Faculty of Medicine- Alexandria University (2009)
Resident of Obstetrics and Gynaecology
Al Ameryia Hospital - Alexandria*

Under Supervision of

Dr. Nashwa El-Said Hassan

*Assistant Professor of Obstetrics and Gynaecology
Faculty of Medicine - Ain Shams University*

Dr. Mohammed Mahmoud Abd El-Aleem

*Lecturer in Obstetrics and Gynaecology
Faculty of Medicine – Ain Shams University*

**Faculty of Medicine
Ain Shams University
2017**

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

قالوا

سبحانك لا علم لنا
إلا ما علمتنا إنك أنت
العليم الحكيم

صدق الله العظيم

سورة البقرة الآية: ٣٢



*Above and before all, I would like to kneel thank to **Allah** the all Mighty, the most Merciful for the support, guidance and mercy He grants me throughout my life.*

*I want to express my deepest gratitude to **Dr. Nashwa El-Said Hassan**, Assistant Professor of Obstetrics and Gynaecology, Faculty of Medicine, Ain Shams University, for her positive attitude and support.*

*I would like to thank to **Dr. Mohammed Mahmoud Abd El-Aleem**, Lecturer of Obstetrics and Gynaecology, Faculty of Medicine, Ain Shams University, for his guidance and suggestions which were of great value to me.*

Finally my truthful affection and love to My Family, specially my husband, who their and will always be, by my side all my life.

Basma Yehia



CONTENTS

Subject	Page No.
List of Abbreviations	i
List of Tables	iii
List of Figures	iv
Protocol	
Introduction	1
Aim of the Work	3
- Chapter (1): Primary Postpartum Hemorrhage	4
- Chapter (2): Tranexamic Acid	34
Patients and Methods	48
Results	59
Discussion	78
Summary	85
Conclusion and Recommendations	90
References	91
Arabic Summary	v

List of Abbreviations

Abb.	Full term
ACOG	American College of Obstetricians Gynaecologists
AMTSL	Active Management of Third Stage of Labor
BMI	Body Mass Index
CC	Cubic Centimeter
CCT	Controlled Cord Traction
CRASH	Clinical Randomisation of an Antifibrinolytic in Significant Haemorrhage
HB	Hemoglobin
HCT	Hematocrite
HELLP	Haemolysis, Elevated liver enzyme levels, and Low platelets level
ICU	Intensive Care Unit
IM	Intramuscular
IU	International Unit
IV	Intravenous
Max	Maximum
Mg	Milligram
Min	Minute
ml	Milliliter
MRSA	Methicillin resistant staph aureous
PPH	Post Partum Hemorrhage
RCOG	Royal College for Obstetricians and Gynaecologists
RCT	Randomized Controlled Trial

Abb.	Full term
SD	Standard Deviation
TRAAP	TRAnexmic Acid for Prevention of Postpartum Hemorrhage
TXA	Tranexamic Acid
UNICEF	United Nations Children`s Emergency Fund
WHO	World health organization

List of Tables

Tables No.	Title	Page No.
Table (1):	Risk factors for PPH	14
Table (2):	Medical Management of Postpartum Hemorrhage	26
Table (3):	Comparison between the two studied groups according to demographic data	59
Table (4):	Comparison between the two studied groups according to obstetric history	60
Table (5):	Comparison between the two studied groups according to amount of blood loss	60
Table (6):	Comparison between the two studied groups according to hemoglobin Hb	63
Table (7):	Comparison between the two studied groups according to Hct	67
Table (8):	Comparison between the two groups according to pulse rate	70
Table (9):	Comparison between the different periods according to blood pressure	71
Table (10):	Comparison between the two studied groups according to ecbolics needed to control the bleeding	73
Table (11):	Comparison between the two studied groups according to the need for blood transfusion	75

List of Figures

Figure No.	Title	Page No.
Fig. (1):	Brandt-Andrews maneuver for cord traction	11
Fig. (2):	Haemostatic brace sutures (B lynch sutures)	27
Fig. (3):	Bilateral uterine artery ligation	29
Fig. (4):	Algorithm for management of postpartum hemorrhage	33
Fig. (5):	Structural formula of tranexamic acid	34
Fig. (6):	Comparison between the two studied groups according to the blood loss during and after delivery	61
Fig. (7):	Comparison between the two studied groups according to total blood loss	61
Fig. (8):	Comparison between the two studied groups according to HB before and after delivery	64
Fig. (9):	Comparison between the two studied groups according to % of change of HB	64
Fig. (10):	Comparison between the two studied groups according to hematocrit Hct before and after delivery	68
Fig. (11):	Comparison between the two studied groups according to % of change of hematocrit Hct	68
Fig. (12):	Comparison between the two studied groups according to pulse rate	70

<i>Figure No.</i>	<i>Title</i>	<i>Page No.</i>
Fig. (13):	Comparison between the studied groups according to blood pressure	72
Fig. (14):	Comparison between the two studied groups according to ecbolics needed to control the bleeding	74
Fig. (15):	Comparison between the two studied groups according to the amount of ecbolic needed to control the bleeding	74
Fig. (16):	Comparison between the two studied groups according to the need for blood transfusion	76

Protocol



Introduction



Aim of the Work



Review of Literature



Primary Postpartum Hemorrhage



Tranexamic Acid

