

127, 17 27, 17 (20) 77, 17 (20









جامعة عين شمس

التوثيق الالكتروني والميكروفيلم



نقسم بللله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأفلام قد اعدت دون آية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15-20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of 15 – 25c and relative humidity 20-40 %



ثبكة المعلومات الجامعية





Information Netw. " Shams Children Sha شبكة المعلومات الجامعية @ ASUNET بالرسالة صفحات لم ترد بالأص

A STUDY OF SOME DETERMINANTS OF PREMENSTRUAL SYNDROME (PMS) AMONG A GROUP OF TEACHERS IN ALEXANDRIA

Thesis

Submitted to the High Institute of Public Health
University of Alexandria
In Partial Fulfillment of The Requirements of the
Degree of Master in Public Health

(Maternal and Child Health)

By

Randa Amir Wasfy Fahmy (M.B.B.Ch.)

> High Institute of Public Health University of Alexandria

Bac. E

SUPERVISORS

Dr. Nihad Ibrahim Dabbous

Assistant Professor of MCH

Family Health Department

High Institute of Public Health

Alexandria University

N/Abbou &

Hedhat Ath

Dr. Medhat Salah Eldin Attia

Assistant Drofessor of Mental Health Family Health Department Kigh Institute of Dublic Kealth Alexandria University

Dr. Kholoud Yehia Tayel 16 bud Tay


Assistant Professor of School Health Family Kealth Department Righ Institute of Lublic Kealth Alexandria University

Dr. Mona Hassan Ahmed

Lecturer of Biostatistics Biostatistics Department \ Kigh Institute of Lublic Kealth Alexandria University

Dedicated

To My Great Mum and Dad
To My Lovable Kusband
To my beloved daughters





First and foremost thanks to God.

I wish to express my deepest gratitude and appreciation to **Dr. Nihad Dabbous**, Assistant Professor of Family Kealth, HIPH, Alexandria University, who offered me generous assistance, support and supervision and for her continuous advice and help.

I am most grateful to **Dr. Medhat Attia**, Assistant Lrofessor of Family Health, HIPH, Alexandria University, for his precise and valuable supervision and suggestions, continuous support and constructive criticism.

My sincere thanks to **Dr. Kholoud Tayel**, Assistant Drofessor of Family Health, HIPH, Alexandria University, for her generous cooperation all through the study and her effort in revising this work.

I would like to offer my thanks to **Dr. Mona Hassan**,

Lecturer of Biostatistics, HIPH, Alexandria University,

for her effort in performing the statistical design and

analysis of this work and for her valuable guidance.

My warm appreciation to personnel of the Statistical Department in the HIPH who offered help in the formulation of the results of this research.

Last but not least, I owe special thanks and gratitude to all who helped me by time, effort or spirit in fulfilling and performance of this work.

Contents D

Chapter		Page
I.	Introduction	1
II.	Aim of The Work	40
III.	Material and Methods	41
IV.	Results	55
V.	Discussion	103
VI.	Conclusion and Recommendations	127
VII.	Summary	130
VIII.	References	135
·	Appendices	
	Arabic Summary	i

List of Tables

Table		Page
(I)	Distribution of the study sample according to their personal characteristics.	56
(II)	Distribution of the study sample according to menstrual history, obstetric history and feminine role acceptance score.	59
(III)	Distribution of the study sample according to the severity of certain selected symptoms on MDQ and different menstrual cycle phases.	62
(IV)	Summary statistics of the MDQ seven scales at the premenstrual phase.	64
(V)	Distribution of the study sample on the seven MDQ scales according to their scores.	67
(VI)	Intercorrelations of the seven MDQ scales across the three cycle phases.	70
(VII)	Intercorrelations of MDQ premenstrual symptoms scales.	72
(VIII)	Distribution of the high scorers on the MDQ seven scales for the premenstrual symptoms according to their age.	75
(IX)	Distribution of the high scorers on the MDQ seven scales for the premenstrual symptoms according to their consumption of caffeine-containing beverages/day.	77

Table		Page
(X)	Distribution of the high scorers on the MDQ seven scales for the premenstrual symptoms according to their age at menarche.	; 79 -
(XI)	Distribution of the high scorers on the MDQ seven scales for the premenstrual symptoms according to regularity of menstrual cycle.	81
(XII)	Distribution of the high scorers on the MDQ seven scales for the premenstrual symptoms according to the length of menstrual cycle in days.	85 i
(XIII)	Distribution of the high scorers on the MDQ seven scales for the premenstrual symptoms according to the number of days of bleeding.	88
(XIV)	Distribution of the high scorers on the MDQ seven scales for the premenstrual symptoms according to their predictability of next menses.	90
(XV)	Distribution of the high scorers on the MDQ seven scales for the premenstrual symptoms according to parity.	93
(XVI)	Distribution of the high scorers on the MDQ seven scales for the premenstrual symptoms according to method of contraception used.	96
(XVII)	Correlation between ferminine role acceptance and MDQ scales.	98
(XVIII)	Results of Stepwise Logistic Regression Analysis of factors affecting MDQ scales.	101

i

List of Figures

Figure		Page
(1)	Average T score of the MDQ seven scales at the premenstrual phase.	65
(2)	Distribution of the high scorers of the study sample on the seven MDQ scales.	68
(3)	Distribution of the high scorers on the MDQ seven scales for the premenstrual symptoms according to regularity of menstrual cycle.	82
(4)	Distribution of the high scorers on the MDQ seven scales for the premenstrual symptoms according to the length of menstrual cycle in days.	86
(5)	Distribution of the high scorers on the MDQ seven scales for the premenstrual symptoms according to their predictability of next menses.	91

INTRODUCTION