

Role of different ultrasound approaches in diagnosis of female pelvic floor dysfunction

A Thesis Submitted for Partial Fulfilment
Of M.D. degree in
Radiodiagnosis

By
Eman Abdel Salam Ahmed Hasan
MSc (Radiodiagnosis)

Supervised by

Professor Doctor Mamdouh Ahmed Ghoneim
Professor of Radiodiagnosis
Faculty of Medicine
Ain Shams University

**Professor Doctor Motaz Mohamed Samy EL-
Beblawy**
Professor of Radiodiagnosis
Faculty of Medicine
Ain Shams University

Doctor Omar Mohammad Atef Sharaf
Lecturer of Radiodiagnosis
Faculty of Medicine
Ain Shams University

Faculty of Medicine
Ain Shams University

2011

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

"قَالُوا سُبْحَانَكَ لَا عِلْمَ لَنَا إِلَّا مَا عَلَّمْتَنَا

إِنَّكَ أَنْتَ الْعَلِيمُ الْحَكِيمُ"

صدق الله العظيم

سورة البقرة - آية ٣٢

Acknowledgment

First, thanks are all due to God for blessing this work until it has reached its end as a part of his generous help throughout my life.

I wish to express my thanks and profound gratitude to Professor Dr. Mamdouh Ahmed Ghoneim, professor of Radiodiagnosis, Ain Shams University. Whom I am deeply indebted for encouraging me to develop this work, and for the valuable supervision and continuous help, he has given me since I started this work,

Also, I am deeply indebted to Professor Dr. Motaz Mohamed Samy EL-Beblawy, Assistant Professor of Radiodiagnosis, Ain Shams University, for his valuable instructions, inspiring guidance and support throughout this work,

My sincere thanks and utmost appreciation are humbly presented to Dr. Omar Mohammad Atef Sharaf, Assistant Professor of Radiodiagnosis, Ain Shams University, for his supervision and assistance.

Also, my deepest gratitude to Dr. Mohamed Samir Eid Sweed, lecturer of Obstetrics and Gynecology, Ain Shams University, for his help with the clinical examination and recruitment of the patients.

Finally, I would like to express my deepest gratitude to my whole family who offered me support, advice and motivation.

Eman Abdel Salam.

CONTENTS

INTRODUCTION:	1
AIM OF THE WORK:	3
REVIEW OF LITERATURE:	4
PATIENTS AND METHODS:	74
RESULTS:	78
ILLUSTRATIVE CASES:	88
DISCUSSION:	128
SUMMARY AND CONCLUSION:	135
REFERENCES:	137
ARABIC SUMMARY	

List of figures

Figure	Title	Page
Fig. 1	Anatomy of female pelvic floor.	4
Fig. 2	Schematic diagrams of the pelvic floor, specifically the levator ani musculature and its fascial condensations.	5
Fig. 3	Pelvic view of the levator ani demonstrating its components: puborectalis, pubococcygeus, iliococcygeus, and coccygeus.	10
Fig. 4	Ultrasound image of a healthy bladder before (A) and after (B) a pelvic-floor muscle (PFM) contraction.	37
Fig. 5	Transabdominal ultrasound image of a paravaginal defect (PVD).	37
Fig. 6	Ultrasonographic section at the level of mid-urethra (U) of a woman with normal support.	40
Fig. 7	(A) Scanning at a higher plane. The obturator internus (Ob) muscle is visible laterally covered by the hyperechogenic fascia (obturator fascia) (Ob-f). (B) Iliococcygeus muscle (IC) originating from the obturator fascia (Ob-f). The vaginal wall (V) loses the H shape appearance and becomes more flattened.	41
Fig. 8	The anterior vaginal sulci (V) are visualised laterally to the proximal urethra (U) giving a characteristic 'H shape' appearance.	42

List of figures (ii)

Fig. 9	Anal endosonography. Normal anatomy of the anal sphincter and puborectalis muscle.	43
Fig. 10	Perineal sonography obtained with the introital approach using an endovaginal probe at 6.5 MHz (Sagittal view).	45
Fig. 11	An axial view of the four-ring target like urethra is seen.	46
Fig.12	Axial view of the anal canal by introital ultrasonography.	46
Fig.13	A, Transducer placement on perineum and B, schematic representation of imaging in midsagittal.	48
Fig. 14	Midsagittal view by transperineal sonography.	49
Fig. 15	Measurements according to Schaer et al., by rectangular coordinate system.	51
Fig. 16	Determination of bladder neck mobility.	52
Fig. 17	Measurements according to Creighton et al.	52
Fig. 18	Using 5-7 MHZ endocavitary probe. The posterior urethrovesical angle is defined as the angle between the urethral axis and the bladder base as it proceeds to the trigone.	53
Fig. 19	The urethropelvic angle is defined as the angle formed by the convergence between the line that passes through the pubis and the urethral axis.	53
Fig. 20	Urethral rotation angle.	54

List of figures (iii)

Fig. 21	Measurement of the height of bladder neck.	55
Fig. 22	Measurement of bladder wall thickness in a patient with symptoms of an overactive bladder and detrusor instability.	56
Fig. 23	The inferior margin of the symphysis pubis serves as a line of reference against which the maximal descent of bladder, uterus, cul de sac and rectal ampulla on Valsalva maneuver can be measured.	57
Fig. 24	Measurement of internal (c) and external anal sphincter (d).	58
Fig. 25	A typical finding in a patient with stress incontinence and anterior vaginal wall descent with postero-inferior rotation of the urethra, opening of the retrovesical angle and funneling of the proximal urethra.	60
Fig. 26	Ultrasonographic appearance of bladder neck position during rest and strain showing increased proximal pubourethral distance from 9 mm at rest to 21 mm at strain.	61
Fig. 27	Ultrasonographic appearance of urethral diameter. It measures 8 mm indicating intrinsic sphincter deficiency.	61
Fig. 28	Gartner duct cyst (arrow) close to bladder neck.	62
Fig. 29	Transitional cell carcinoma of the bladder (arrow) as seen on parasagittal translabial ultrasound.	63

Fig. 30	Urethral diverticulum (arrow) herniating downwards and clinically simulating a cystourethrocele as shown at rest (a) and on Valsalva maneuver (b). The neck of the diverticulum is seen close to the bladder neck.	63
Fig. 31	Typical translabial ultrasound images (midsagittal plane) in a patient with a symptomatic, single compartment prolapse in the form of cystocele.	64
Fig. 32	The two main types of cystocele as imaged on maximal Valsalva in the midsagittal plane.	65
Fig. 33	Paravaginal defect.	66
Fig. 34	First-degree uterine descent in a patient after Burch colposuspension.	67
Fig. 35	Rectocele on translabial ultrasonography.	67
Fig. 36	The distinction between a ‘true rectocele’ and perineal hypermobility.	68
Fig. 37	Recto-enterocele after Burch colposuspension at rest (a) and on Valsalva maneuver (b).	69
Fig. 38	Anterior external sphincter defect by endoanal ultrasonography.	70
Fig. 39	Internal anal sphincter defect (ISD) by endoanal Ultrasonography.	70
Fig. 40	Defect of the internal and external sphincter complex.	71
Fig. 41	Synthetic implants such as tension-free vaginal tape (TVT) or suprapubic arc (SPARC) tape are easily visualized as highly echogenic structures posterior to the urethra.	73

List of Tables

Table	Content	Page
Table 1	Distribution of age and body mass index (BMI) among study group and control.	78
Table 2	Parity among the study group.	79
Table 3	Mode of delivery among the study group.	79
Table 4	Menstrual status among the study group.	79
Table 5	Types of urinary incontinence (UI) among the study group.	80
Table 6	Difference in ultrasonographic measurements between women with urinary incontinence (UI) among study group and control group.	81
Table 7	Difference in ultrasonographic measurements between women with SUI among study group and control group.	82
Table 8	Difference in ultrasonographic measurements between women with urge incontinence (uI) among study group and control group.	83
Table 9	Difference in ultrasonographic measurements between cases with SUI and uI.	84
Table10	Comparison of ultrasonographic measurements during rest and straining in cases with SUI.	85

List of tables (ii)

Table11	Comparison of urethral diameter among the subgroups of SUI with and without ISD.	85
Table12	Distribution of cases according to clinical examination.	86
Table13	Cases of cystocele with paravaginal defects (PVD).	86
Table14	Cases of rectocele with perineal hypermobility and recto-vaginal septal defect.	86

INTRODUCTION

AIM OF THE WORK

REVIEW OF LITERATURE