EFFECT OF SOME AGRICULTURAL PRACTICES ON GROWTH, YIELD AND ACTIVE CONSTITUENTS OF MEXICAN MARIGOLD (*TAGETES LUCIDA*) PLANT

By

RASHA FOUAD ISMAIL AHMED

B.Sc. Agric. Sc. (Horticulture), Ain Shams University, 2008

A thesis submitted in partial fulfillment of the requirements for the degree of

MASTER OF SCIENCE

in

Agricultural Science (Ornamental Plants)

Department of Horticulture Faculty of Agriculture Ain Shams University

EFFECT OF SOME AGRICULTURAL PRACTICES ON GROWTH, YIELD AND ACTIVE CONSTITUENTS OF MEXICAN MARIGOLD (*TAGETES LUCIDA*) PLANT

By

RASHA FOUAD ISMAIL AHMED

B.Sc. Agric. Sc. (Horticulture), Ain Shams University, 2008

Under the supervision of:

Dr. Awaad Mohamed Kandeel

Prof. Emeritus of Ornamental and Medicinal & Aromatic Plants, Dept.of Horticulture, Faculty of Agriculture, Ain Shams University. (Principal Supervisor).

Dr. Ayman Kamal Ibrahim

Assistant Prof. of Ornamental and Medicinal & Aromatic Plants, Dept.of Horticulture, Faculty of Agriculture, Ain Shams University.

Dr. Elsayed Abouelfotowh Omer

Research Prof. Emeritus of Medicinal and Aromatic Plants, Dept. of Medicinal& Aromatic Plants Research, National Research Centre.

Approval Sheet

EFFECT OF SOME AGRICULTURAL PRACTICES ON GROWTH, YIELD AND ACTIVE CONSTITUENTS OF MEXICAN MARIGOLD (*TAGETES LUCIDA*) PLANT

By

RASHA FOUAD ISMAIL AHMED

B.Sc. Agric. Sc. (Horticulture), Ain Shams University, 2008

This Thesis for M.Sc. Degree has been approved by:		
Dr. Ahmed Atef Sadek Radwan		
Head of Research Emeritus of Medicinal and Aromatic Plants,		
Horticulture Research Institute, Agricultural Research Centre.		
Dr. Sohair El-Sayed Mohamed Hassan		
Prof. Emeritus of Ornamental and Medicinal & Aromatic Plants,		
Faculty of Agriculture, Ain Shams University.		
Dr. Awaad Mohamed Kandeel		
Prof. Emeritus of Ornamental and Medicinal & Aromatic Plants,		
Faculty of Agriculture, Ain Shams University.		
•		

Date of examination: 14/5/2013.

ACKNOWLEDGEMENT

First of all, I wish that I could thank Allah, the Almighty, the most beneficent, most merciful for all countless graces and overcoming all the obstacles in front of me to achieve goals and make them possible.

My heart is full of thanks and gratefulness to Spirit of Late *Prof.Dr.Mahmoud Elsayed Hashem*, Professor of Ornamental and Medicinal & Aromatic Plants, Faculty of Agriculture, Ain Shams University, I pray to Allah to grant him mercy, forgiveness and continue living in paradise Insha'Allah.

My deepest thanks and gratitude are extended to *Prof.Dr.Awaad Mohamed Kandeel*, Emeritus Professor of Ornamental and Medicinal & Aromatic Plants, Faculty of Agriculture, Ain Shams University, for his continuous supervision, valuable guidance, indispensable advice, beneficial instructions, moral encouragement, worthy support during the courses and the curriculum of this work as well as his suave help in the performance of the thesis.

Sincere appreciation and special gratefulness are due to *Prof.Dr.Elsayed Abouelfotowh Omer*, Emeritus Professor of Medicinal and Aromatic Plants, National Research Centre for suggesting the subject of this work, his constant supervision, everlasting encouragement, constructive criticism, inspiring guidance, providing all necessary facilities required for the experimental work as well as his fruitful help in writing and performing the thesis.

I wish to express my indebtedness and profound gratitude to *Dr.Ayman Kamal Ibrahim*, Assistant Professor of Ornamental and Medicinal & Aromatic Plants, Faculty of Agriculture, Ain Shams University, for his supervision, constructive encouragement during courses and his pleasant help in performing the thesis.

Special grateful and thanks to *Dr.Saber Fayez Hendawy*, Associate Professor of Medicinal and Aromatic Plants, National Research Centre, for

his everlasting support, continuous help and great effort in the performance of the agricultural experiment.

Many and great thanks are due to *Prof.Dr.Fayez Nazir Zaki*, *Prof.Dr.Mohamed Fared Abd-ElFataah*, *Prof.Dr.Mahfouz Abd-Elgawad* and *Dr.Azza Nour El-Deen* for their help and valuable effort in the biological activity experiments.

Thanks are expressed to SEKEM Company for allowing me to cultivate in its farm. Also deep thanks to all staff members of Medicinal and Aromatic Plants Research Department, National Research Centre for their help, encouragement and co-operation during this work.

Finally, I dedicate this work to the most precious people in my life; to spirit of my grandfather and grandmother, to my dear parents who are the cause of my happiness and all beautiful things in my life, to my beloved sisters (*Heba and Hend*) and my dear brother (*Abd El-Rahman*) for their valuable cheer, help and support as well as to my sweethearts (*Alaa, Areeg and Ahmed*).

Special dedication is given to my dear *Prof.Dr.Saeed El-Shazly* for supporting and encouraging me.

CONTENTS

LIST OF TABLES	
LIST OF FIGURES	
INTRODUCTION	
REVIEW OF LITERATURE	
MATERIALS AND METHODS	
RESULTS AND DISCUSSION	
1. Irrigation intervals experiment	
1.1. Growth and yield charachters.	
1.1.1. Plant height	
1.1.2. Plant diameter	
1.1.3. Number of branches / plant	
1.1.4. Fresh and dry weights of herb	
1.1.5. Fresh and dry yield of herb	
1.2. Chemical constituents	
1.2.1. Essential oil percentage	
1.2.2. Essential oil content and oil yield	
1.2.3. The main constituents of the essential oil	
1.3. Chemical composition.	
1.3.1. Total flavonoids content, total coumarins content and	
photosynthesis pigments content	
1.3.2. Proline content	
2. Planting date and plant spacing experiment	
2.1. Growth and yield characters	
2.1.1. Plant height	
2.1.2. Plant diameter	
2.1.3. Number of branch / plant	
2.1.4. Fresh and dry weights of herb.	
2.1.5. Fresh and dry yield	
2.2. Chemical constituents.	

2.2.1. Essential oil percentage	10
2.2.2. Essential oil content and yield	10
2.2.3 The main constituents of the essential oil	1
2.3. Chemical composition	13
2.3.1. Total flavonoids content	1.
2.3.2. Total coumarins content	1.
3. Biolgical activity	1.
3.1. Antioxidant activity	1.
3.2. Antimicrobial activity	1.
3.3. Insectcidal activity	1.
3.4. Nematocidal activity	14
SUMMARY AND CONCLUSION	14
REFERENCES	1:
ARABIC SUMMARY	

LIST OF TABLES

Table (1-a)	Physical and chemical properties of the experimental soil of the SEKM Farm during 2009 and 2011 seasons	33
Table (1-b)	Monthly average of metrological data of the experimental farm of SEKEM, Bilbas during 2009 and 2011 seasons	34
Table (1-c)	The chemical analysis of the used compost during 2009 and 2011 seasons.	35
Table (2-a)	Physical and chemical properties of the experimental soil of Nubaria farm during 2009 and 2010 seasons	37
Table (2-b)	Monthly average of metrological data of the Experimental farm of Nubaria farm during 2009 and 2010 seasons	38
Table (2-c)	The chemical analysis of the used compost during 2009 and 2010 seasons	39
Table (3)	Effect of irrigation intervals on plant height of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	47
Table (4)	Effect of irrigation intervals on plant diameter (cm) of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	50
Table (5)	Effect of irrigation intervals on number of branches / plant of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	52
Table (6)	Effect of irrigation intervals on herb fresh and dry weight (g / plant) of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	54
Table (7)	Effect of irrigation intervals on herb fresh and dry yield (Ton / Feddan) of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	58

Table (8)	Effect of irrigation intervals on essential oil percentage (%) of <i>Tagetes</i>	
	lucida plants during 2009 and 2011 seasons	63
Table (9)	Effect of irrigation intervals on essential oil content (ml / plant) and yield (L / Feddan) of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons.	66
Table (10)	The main constituents of the essential oil of <i>Tagetes lucida</i> as affected by irrigation intervals in two cuts during 2009 season	70
Table (11)	Effect of irrigation intervals on total flavonoids content (%), coumarins content (mg / 100 g) and photosynthesis pigments (chlorophyll a, b and carotenoids) content of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons.	77
Table (12)	Effect of irrigation intervals on proline content (mg / g) of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	80
Table (13)	Effect of planting date and plant spacing on plant height (cm) of <i>Tagetes lucida</i> plants during 2009 and 2010 seasons	83
Table (14)	Effect of planting date and plant spacing on plant diameter (cm) of <i>Tagetes lucida</i> plants during 2009 and 2010 seasons	87
Table (15)	Effect of planting date and plant spacing on number of branches / plant of <i>Tagetes lucida</i> plants during 2009 and 2010 seasons	91
Table (16)	Effect of planting date and plant spacing on herb fresh and dry weight (g / plant) of <i>Tagetes lucida</i> plants during 2009 and 2010 seasons	94
Table (17)	Effect of planting date and plant spacing on herb fresh and dry yield (Ton / Feddan) of <i>Tagetes lucida</i> plants during 2009 and 2010 seasons	100
		_ 0

Table (18)	Effect of planting date and plant spacing on essential oil percentage	
(_0)	(%) of <i>Tagetes lucida</i> plants during 2009 and 2010 seasons	105
Table (19)	Effect of planting date and plant spacing on essential oil content (ml / plant) and yield (L / Feddan) of <i>Tagetes lucida</i> plants during 2009 and 2010 seasons.	108
Table (20)	Effect of planting date and plant spacing on the main constituents of the essential oil of <i>Tagetes lucida</i> in the 1 st cut during 2009 season	114
Table (21)	Effect of planting date and plant spacing on the main constituents of the essential oil of <i>Tagetes lucida</i> in the 2 nd cut during 2009 season	115
Table (22)	Effect of planting date and plant spacing on the main constituents of the essential oil of <i>Tagetes lucida</i> in the 3 rd cut during 2009 season.	116
Table (23)	Methyl chaviol and monoterpens in the essential oil of <i>T. lucida</i> as affected by planting dates and plant spacing in three cuts during 2009 season.	129
Table (24)	Total oxygenated and non oxygenated compounds of <i>T. lucida</i> essential oil as affected by planting dates and plant spacing in three cuts of during 2009 season	130
Table (25)	Effect of planting date and plant spacing on total flavonoids content (%) and coumarins content (mg / 100g) of <i>Tagetes lucida</i> plants during 2009 and 2010 seasons	131
Table (26)	Antioxidant activity of the different concentrations and IC_{50} of T . <i>lucida</i> herb extract using DPPH method	137
Table (27)	Inhibition zone (diameter (cm)) of the essential oil of <i>T. lucida</i> for some microorganism	138
Table (28)	Effect of <i>Tagetes lucida</i> extract on the average no. of aphid (<i>Aphis brassicaceae</i>) on cabbage aphids before and after treatment	139

Table (29)	Effect of herb and root extract from Mexican marigold (<i>Tagetes lucida</i>) on percentage motility of soil stages of phytonematodes and their recovery in distilled water	141
Table (30)	Effect of herb and root extract from Mexican marigold (<i>Tagetes lucida</i>) on development to hatching of <i>Meloidogyne incognita</i> eggs.	142

VII

LIST OF FIGURES

Fig. (1)	Effect of irrigation intervals on plant height (cm) of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	48
Fig. (2)	Effect of irrigation intervals on plant diameter (cm) of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	51
Fig. (3)	Effect of irrigation intervals on number of branches / plant of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	53
Fig. (4)	Effect of irrigation intervals on herb fresh weight (g / plant) of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	55
Fig. (5)	Effect of irrigation intervals on herb dry weight (g / plant) of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	56
Fig. (6)	Effect of irrigation intervals on herb fresh yield (Ton / Feddan) of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	59
Fig. (7)	Effect of irrigation intervals on herb dry yield (Ton / Feddan) of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	60
Fig. (8)	Effect of irrigation intervals on essential oil percentage (%) of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	64
Fig. (9)	Effect of irrigation intervals on essential oil content (ml / plant) of <i>Tagetes lucida</i> plants during 2009 and 2011 season	67
Fig. (10)	Effect of irrigation intervals on essential oil yield (L / Feddan) of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	68
Fig. (11)	GL Chromatogram of the essential oil of <i>T. lucida</i> plants irrigated with 2 days intervals in the 1 st cut during 2009 season	71

VIII

Fig. (12)	GL Chromatogram of the essential oil of <i>T. lucida</i> plants irrigated with 4 days intervals in the 1 st cut during 2009 season	72
Fig. (13)	GL Chromatogram of the essential oil of <i>T. lucida</i> plants irrigated with 6 days intervals in the 1 st cut during 2009 season	73
Fig. (14)	GL Chromatogram of the essential oil of <i>T. lucida</i> plants irrigated with 2 days intervals in the 2 nd cut during 2009 season	74
Fig. (15)	GL Chromatogram of the essential oil of T . lucida plants irrigated with 4 days intervals in the 2^{nd} cut during 2009 season	75
Fig. (16)	GL Chromatogram of the essential oil of <i>T. lucida</i> plants irrigated with 6 days intervals in the 2 nd cut during 2009 season	76
Fig. (17)	Effect of irrigation intervals on total flavonoids content (%) and coumarins content (mg / 100 g) of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	78
Fig. (18)	Effect of irrigation intervals on photosynthesis pigments (chlorophyll a, b and carotenoids) of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	7 9
Fig. (19)	Effect of irrigation intervals on proline content (mg / g) of <i>Tagetes lucida</i> plants during 2009 and 2011 seasons	81
Fig. (20)	Effect of planting date and plant spacing on plant height (cm) of <i>Tagetes lucida</i> plants during 2009 and 2010 seasons	84
Fig. (21)	Effect of planting date and plant spacing on plant diameter (cm) of <i>Tagetes lucida</i> plants during 2009 and 2010 seasons	88
Fig. (22)	Effect of planting date and plant spacing on number of branches / plant of <i>Tagetes lucida</i> plants during 2009 and 2010	92
Fig. (23)	Effect of planting date and plant spacing on herb fresh weight (g / plant) of <i>Tagetes lucida</i> plants during 2009 and 2010 seasons	95

Fig. (24)	plant) of <i>Tagetes lucida</i> plants during 2009 and 2010 seasons	96
Fig. (25)	Effect of planting date and plant spacing on herb fresh yield (Ton / Feddan) of <i>Tagetes lucida</i> plants during 2009 and 2010 seasons	101
Fig. (26)	Effect of planting data and plant anguing on both day yield (Ton /	102
Fig. (27)	Effect of planting date and plant spacing on essential oil percentage (%) of <i>Tagetes lucida</i> plants during 2009 and 2010 seasons.	106
Fig. (28)	(ml / plant) of Tagetes lucida plants during 2009 and 2010	109
Fig. (29)	Effect of planting date and plant spacing on essential oil yield (L / Feddan) of <i>Tagetes lucida</i> plant during 2009 and 2010 seasons.	110
Fig. (30)	GL Chromatogram of the essential oil of <i>T. lucida</i> plants cultivated in February at 60 cm plant spacing in the 1 st cut during 2009 season.	117
Fig. (31)	GL Chromatogram of the essential oil of <i>T. lucida</i> plants cultivated in April at 40 cm plant spacing in the 1 st cut during 2009 season	118
Fig. (32)	GL Chromatogram of the essential oil of <i>T. lucida</i> plants cultivated in February at 50 cm plant spacing in the 2 nd cut during 2009 season	119
Fig. (33)	GL Chromatogram of the essential oil of <i>T. lucida</i> plants cultivated in February at 60 cm plant spacing in the 2 nd cut during 2009 season.	120

Fig. (34)	GL Chromatogram of the essential oil of <i>T. lucida</i> plants cultivated in April at 40 cm plant spacing in the 2 nd cut during 2009 season
Fig. (35)	GL Chromatogram of the essential oil of <i>T. lucida</i> plants cultivated in February at 40 cm plant spacing in the 3 rd cut during 2009 season.
Fig. (36)	GL Chromatogram of the essential oil of <i>T. lucida</i> plants cultivated in February at 50 cm plant spacing in the 3 rd cut during 2009 season.
Fig. (37)	GL Chromatogram of the essential oil of <i>T. lucida</i> plants cultivated in February at 60 cm plant spacing in the 3 rd cut during 2009 season.
Fig. (38)	GL Chromatogram of the essential oil of <i>T. lucida</i> plants cultivated in March at 60 cm plant spacing in the 3 rd cut during 2009 season
Fig. (39)	GL Chromatogram of the essential oil of <i>T. lucida</i> plants cultivated in April at 40 cm plant spacing in the 3 rd cut during 2009 season.
Fig. (40)	GL Chromatogram of the essential oil of <i>T. lucida</i> plants cultivated in April at 50 cm pant spacing in the 3 rd cut during 2009 season
Fig. (41)	Effect of planting date and plant spacing on total flavonoids content (%) of <i>Tagetes lucida</i> dry herb during 2009 and 2010 seasons.
Fig. (42)	Effect of planting date and plant spacing on coumarins content (mg / 100 g) of <i>Tagetes lucida</i> dry herb during 2009 and 2010 seasons.
Fig. (43)	Effect of <i>Tagetes lucida</i> extract on the average no. of aphid (<i>Aphis brassicaceae</i>) on cabbage aphids before and after treatment.