

**Comparing Frontalis Sling Operation;
Supralash Stab Incision versus the Eyelid
Crease incision for the Treatment of Ptosis**

Thesis

*Submitted for Partial Fulfillment of Master
Degree in Ophthalmology*

By

Noha Ahmed Abdul-Khaliq

M.B.B.Ch

Faculty of Medicine, Ain Shams University

Under Supervision of

Prof. Dr. Saad Mohamed Rashad

Professor of Ophthalmology

Faculty of Medicine, Ain Shams University

Dr. Ahmed Mohamed Shafik

Lecturer of Ophthalmology

Faculty of Medicine, Ain Shams University

Dr. Mahmoud Ahmed EL-samkary

Lecturer of Ophthalmology

Faculty of Medicine, Ain Shams University

Ophthalmology Department

Faculty of Medicine - Ain Shams University

Cairo, Egypt

2018

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

قالوا

سببنا انك لا تعلم لنا
إلا ما علمتنا إنك أنت
العليم العظيم

صدق الله العظيم

سورة البقرة الآية: ٣٢

Acknowledgment

*First and foremost, I feel always indebted to **ALLAH**, the Most Kind and Most Merciful.*

*I'd like to express my respectful thanks and profound gratitude to **Prof. Dr. Saad Mohamed Rashad**, Professor of Ophthalmology - Faculty of Medicine- Ain Shams University for his keen guidance, kind supervision, valuable advice and continuous encouragement, which made possible the completion of this work.*

*I am also delighted to express my deepest gratitude and thanks to **Dr. Ahmed Mohamed Shafik**, Lecturer of Ophthalmology, Faculty of Medicine, Ain Shams University, for his kind care, continuous supervision, valuable instructions, constant help and great assistance throughout this work.*

*I am deeply thankful to **Dr. Mahmoud Ahmed El-samkary**, Lecturer of Ophthalmology, Faculty of Medicine, Ain Shams University, for his great help, active participation and guidance.*

I would like to express my hearty thanks to all my family for their support till this work was completed.

Last but not least my sincere thanks and appreciation to all patients participated in this study.

Noha Abdul-Khaliq



Dedication

To

MY DEAR PARENTS

*Who gave me too much
And received too little*

List of Contents

Title	Page No.
List of Tables	i
List of Figures	ii
List of Abbreviations	vii
Introduction	1
Aim of the Work.....	3
Review of Literature	
☞ Anatomy of the Eyelid.....	4
☞ Ptosis	14
☞ Diagnosis and Clinical Evaluation	21
☞ Surgical Correction of Ptosis	26
Patients and Methods	38
Results	55
Discussion	70
Conclusion.....	79
Summary.....	80
References	82
Arabic Summary	

List of Tables

Table No.	Title	Page No.
Table (1):	Demographic characteristics among the studied groups.....	55
Table (2):	Comparison between the closed group and the open group regarding pre-operative head posture	57
Table (3):	Comparison between the closed group and the open group regarding pre-op. MRD1	58
Table (4):	Comparison between the closed group and the open group regarding pre-op. Levator function.....	59
Table (5):	Comparison between the closed group & the open group regarding pre-op. eyelid crease.....	60
Table (6):	Comparison between pre and post-operative MRD1 in both groups	61
Table (7):	Comparison between the closed group and the open group regarding post-op. MRD1 grading.....	62
Table (8):	Comparison between the closed group and the open group regarding post-op. lid symmetry.....	64
Table (9):	Comparison between the closed group and the open group regarding post-op. lid contour.....	66
Table (10):	Comparison between the closed group and the open group regarding post-op. lid crease	68
Table (11):	Comparison between the closed group and the open group regarding post-op. lagophthalmos.....	69

List of Figures

Fig. No.	Title	Page No.
Figure (1):	Oblique cross-section of the orbit and adnxea	4
Figure (2):	Surface anatomy of the eyelid.....	5
Figure (3):	Anterior and posterior lamellae of the eyelid.....	6
Figure (4):	Orbicularis oculi muscle	7
Figure (5):	Major retractors of the upper eyelid.....	9
Figure (6):	Orbital septum and tarsi.....	11
Figure (7):	Blood supply to the eyelids	12
Figure (8):	Motor innervation of the lid	13
Figure (9):	Marcus Gunn Jaw-Winking syndrome	15
Figure (10):	Blepharophimosis-ptosis-epicanthus inversus syndrome.....	16
Figure (11):	Bilateral involutional ptosis.....	17
Figure (12):	Brow ptosis	18
Figure (13):	Right third nerve palsy.....	18
Figure (14):	Myasthenia gravis	19
Figure (15):	The interpalpebral fissure height	22
Figure (16):	MRD1 measurement	23
Figure (17):	Levator function measurement	24
Figure (18):	Skin crease measurement.....	24
Figure (19):	Good Bell's phenomenon	25

List of Figures (cont...)

Fig. No.	Title	Page No.
Figure (20):	The Herring law	25
Figure (21):	Algorithm for choice of Surgical Management for a Ptosis Patient.....	27
Figure (22):	Gore-Tex suture material	29
Figure (23):	Fox single pentagon.....	31
Figure (24):	Crawford procedure.....	32
Figure (25):	Mehta's modification of Crawford's technique	33
Figure (26):	Single triangle	33
Figure (27):	Friedenwald and Guyton procedure	34
Figure (28):	Open technique	36
Figure (29):	Left eye of a patient with severe ptosis.	40
Figure (30):	Left eye of a patient with poor levator function.	40
Figure (31):	Marking the eyelid skin.	41
Figure (32):	Insertion of PTFE through the wright needle from central to medial incision.....	42
Figure (33):	Insertion of PTFE through the wright needle from lateral to central incision.	43
Figure (34):	Good eyelid contour.....	43
Figure (35):	A mattress suture.....	44
Figure (36):	Insertion of PTFE from medial eyelid to medial eyebrow incision.	44

List of Figures (cont...)

Fig. No.	Title	Page No.
Figure (37):	Insertion of PTFE from central eyelid to medial eyebrow incision to complete a triangle.....	45
Figure (38):	Double triangle of PTFE.	45
Figure (39):	Adjustment of eyelid height and PTFE Tying.....	46
Figure (40):	Closure of eyebrow incisions.	46
Figure (41):	Marking the crease and the eyebrow skin.	47
Figure (42):	Dissection of orbicularis and levator muscles till the root of lashes.....	47
Figure (43):	Suturing PTFE to the lower third of the tarsus.....	48
Figure (44):	Wright needle passed from medial eyebrow incision to grab the PTFE suture.	48
Figure (45):	The lateral end of PTFE is passed to the lateral eyebrow incision.	49
Figure (46):	Good eyelid contour and height.....	49
Figure (47):	Tying and burying the PTFE in the forehead pocket.	50
Figure (48):	Closure of eyelid incision by interrupted sutures.....	50
Figure (49):	Sex distribution among the two studied groups.	56
Figure (50):	Patient with a head posture.....	57
Figure (51):	Comparison between (closed group) and (open group) regarding pre-op. MRD1.	58

List of Figures (cont...)

Fig. No.	Title	Page No.
Figure (52):	Patient with OD sever ptosis and pre-op. MRD1 = 0mm.....	58
Figure (53):	Comparison between the closed group and the open group regarding pre-op. Levator function.	59
Figure (54):	Patient with severe ptosis & absent crease OU.	60
Figure (55):	Comparison between pre & post-operative MRD1 in the two groups.	61
Figure (56):	Comparison between the closed group and the open group regarding post-op. MRD1 grading.	62
Figure (57):	Right eye of a patient in the closed group 3ms post-op.; graded as excellent.....	63
Figure (58):	Left eye of a patient in the open group 3ms post-op.; graded as excellent.	63
Figure (59):	{a} Left eye ptosis recurrence of a patient in the closed group and {b} left eye ptosis recurrence of a patient in the open group.....	63
Figure (60):	Comparison between the closed group and the open group regarding post-op. eyelid symmetry.....	64
Figure (61):	Comparison between the closed group & the open group regarding post-op. lid contour.....	66
Figure (62):	{a} Left eye of a patient in the closed group graded as excellent symmetry & contour {b}Left eye of a patient in the open group graded as excellent symmetry & contour.....	65

List of Figures (cont...)

Fig. No.	Title	Page No.
Figure (63):	A patient in the open group with right eye focal notching and scarring graded as poor contour.....	67
Figure (64):	Comparison between the closed group and the open group regarding post-op. eyelid symmetry.....	68

List of Abbreviations

<i>Abb.</i>	<i>Full term</i>
<i>&</i>	<i>and</i>
<i>CFEOM</i>	<i>Congenital Fibrosis of the Extraocular Muscles</i>
<i>CPEO</i>	<i>Chronic Progressive External Ophthalmoplegia</i>
<i>LPS</i>	<i>Levator Palpebrae Superioris</i>
<i>m</i>	<i>months</i>
<i>MD</i>	<i>Myotonic Dystrophy</i>
<i>mm</i>	<i>millimeter</i>
<i>MRD</i>	<i>Margin-Reflex Distance</i>
<i>Op.</i>	<i>Operative</i>
<i>OPMD</i>	<i>Oculopharyngeal Muscular Dystrophy</i>
<i>PTFE</i>	<i>Polytetrafluoroethylene</i>
<i>P-value</i>	<i>Probability Value</i>
<i>Sc</i>	<i>Subcutaneously</i>
<i>SD</i>	<i>Standard Deviation</i>
<i>Vs</i>	<i>Versus</i>
<i>X²</i>	<i>Chi-square</i>

ABSTRACT

This study aims at comparing the functional and cosmetic outcomes of correcting congenital ptosis associated with poor levator function via open versus closed approaches using PTFE (Gore-Tex) as a sling material.

Our results revealed that upper eyelid severe ptosis can be effectively managed utilizing sling operation either supraciliary incision (closed) or trans-crease (open) technique, with no significant difference in ptosis recurrence. However, the closed technique appeared to be superior to the open technique regarding the incidence of good post-op. MRD1, eyelid symmetry, and formation of aesthetic eyelid crease rendering the procedure cosmetically and clinically more promising.

Keywords: Congenital Fibrosis of the Extraocular Muscles - Levator Palpebrae Superioris - Myotonic Dystrophy

INTRODUCTION

Ptos, an abbreviation for the term blepharoptosis, refers to vertical narrowing of the palpebral fissure secondary to drooping of the upper eyelid to a lower than normal position. The normal adult upper lid margin is 0.5–2 mm below the superior corneal limbus and is highest just nasal to the pupil ⁽¹⁾.

Congenital ptosis can be isolated or associated with disease of one or more of the extraocular muscles and/or other systemic conditions. More severe forms may involve hypoplasia of the levator–palpebrae–superioris muscle or tendon with a minimal or absent eyelid crease. Congenital ptosis is considered a non-progressive condition; however, it is associated with the development of astigmatism, anisometropia, amblyopia, ocular torticollis, and strabismus. These sequelae of ptosis provide a compelling reason to pursue early surgical correction; for prevention of visual loss related to amblyopia. ⁽²⁾

Acquired ptosis can have a myogenic, neurogenic (peripheral or central lesion of the sympathetic nerves, or the oculomotor nerve), aponeurotic, mechanical, or traumatic cause. ⁽¹⁾ The large majorities are aponeurogenic and are associated with disinsertion or dehiscence of the levator aponeurosis from its distal insertion in the eyelid. ⁽³⁾

Frontalis suspension is indicated in patients with blepharoptosis and poor levator muscle function. “Poor”

function is defined as levator function of 4 mm or less.⁽⁴⁾ This surgery connects the eyelid to the brow with a sling material and utilizes the power of the frontalis muscle to elevate the poorly functioning eyelid.⁽⁵⁾

Cosmetic issues that are raised with standard frontalis suspension surgery include scarring in young children, unsatisfactory geometric tenting of the pretarsal and preseptal skin, and obliteration of the eyelid crease. These may be related to the choice of sling material and to the superficial location of the sling in the eyelid, so several surgical designs and materials have been proposed for frontalis suspension for treating ptosis with poor levator muscle function.⁽²⁾