

# **Relapsed Congenital Talipes Equinovarus**

An Essay Submitted for Partial Fulfillment of Master Degree  
In Orthopaedic Surgery

By

**Ahmed Mohamed Mohamed Mohamed Jaheen**  
M.B.B.Ch

Supervised by

**Professor Mohamed Reda Abdel-Wahab**

Professor of Orthopaedic Surgery  
Faculty of Medicine  
Ain Shams University

**Professor Tarek Hassan Abdelaziz**

Professor of Orthopaedic Surgery  
Faculty of Medicine  
Ain Shams University

Faculty of Medicine  
Ain Shams University  
2009

# **Acknowledgements**

I would like to express my endless gratitude and appreciation to

## **Prof. Mohamed Reda Abdel-wahab**

Professor of Orthopedic Surgery Faculty of Medicine Ain Shams University, for giving me the honor of working under his supervision, for his generous assistance, and providing me with a lot of encouragement and support

My deep thanks are to

## **Prof. Tarek Hassan Abdel-aziz**

Professor of Orthopedic Surgery Faculty of Medicine Ain Shams University, for his true concern, continuous advice during the whole period of the study, and encouragement which allowed completion of this study

And to

Ain Shams faculty of medicine and department of orthopedic surgery.

Children are (patients), not (customers),  
they require (care), not (management)

(G.A. Annas)

# Index

<b>List of Tables</b> .....	<b>I</b>
<b>List of Abbreviation</b> .....	<b>II</b>
<b>List of Figures</b> .....	<b>III</b>
<b>Introduction</b> .....	<b>1</b>
<b>Clubfoot Pathology</b> .....	<b>2</b>
Components of the deformity .....	2
Bony Deformities .....	5
Soft Tissue Contracture .....	10
Articular Malalignments .....	17
<b>Causes of relapse</b> .....	<b>21</b>
Iatrogenic causes .....	21
Congenital Causes .....	30
<b>Diagnosis</b> .....	<b>33</b>
Examination of the foot .....	33
Classification .....	39
<b>Radiological Evaluation</b> .....	<b>45</b>
Radiographs .....	46
MRI .....	54
<b>Conservative treatment</b> .....	<b>59</b>
Manipulation and casting .....	59
Bracing .....	64
Selective Botulinum Toxin Injection .....	75
<b>Operative Treatment</b> .....	<b>77</b>
Anterior Tibial Tendon Transfer .....	80
Soft Tissue Release .....	89
Bony Procedures .....	104
Lateral column shortening procedures .....	104

Double Tarsal Osteotomy .....	106
Os calcis osteotomy .....	107
Metatarsal Osteotomies .....	110
Lambrinudi Arthrodesis .....	112
lower leg Derotation Osteotomy .....	115
Triple Arthrodesis and Talectomy .....	117
<b>Ilizarov Management .....</b>	<b>121</b>
<b>Conclusion .....</b>	<b>127</b>
<b>Summary .....</b>	<b>130</b>
<b>References .....</b>	<b>133</b>
<b>Arabic Summary .....</b>	<b>148</b>

# List of Tables

<b>Table No.</b>	<b>Title</b>	<b>Page</b>
<b>Table 1:</b>	The rating system of Laaveg and Ponseti (1980)	43
<b>Table 2:</b>	Clinical details of patients treated with Ponseti method	63
<b>Table 3:</b>	Patients details treated with botulinium toxin A	76
<b>Table 4:</b>	Therapeutic strategy for resistant clubfoot.	77

## List of Abbreviation

AFO	Ankle foot orthosis
AP	Anteroposterior view
BTX-A	Botulinum toxin-A
CTEV	Congenital talipes equinovarus
DBB	Denis Brown Bar
FDL and FHL	Flexor digitorum longus and Flexor hallucis longus
HS	Hindfoot score
MR	Magnetic Resonance
MRI	Magnetic Resonance Imaging
MS	Midfoot score
SPLATT	Split anterior tendon transfer of tibialis anterior
TNA	Talar neck angle
TN	Talonavicular

## List of Figures

No. of figure	Title	Page
<b>Fig.1-1</b>	Bilateral clubfoot deformity in a 6-week-old infant.	3
<b>Fig.1-2</b>	Relapsed clubfoot of a 3-year-old boy.	5
<b>Fig.1-3</b>	Talus in clubfoot	6
<b>Fig.1-4</b>	The angle between the trochlea of the talus and the talar neck	6
<b>Fig.1-5</b>	3-D reconstruction of a CT scan of a clubfoot (9-years old boy).	9
<b>Fig.1-6</b>	Right clubfoot of 16-week-old fetus sectioned in the transverse plane through the calcaneocuboid joint.	9
<b>Fig.1-7</b>	A, B. Right clubfoot of a 16-week-old fetus, sectioned in the transverse plane through the talonavicular joint.	11
<b>Fig.1-8</b>	Sections in the frontal plane through the malleoli of the clubfoot.	13
<b>Fig.1-9</b>	A and B. In section posterior to those shown in Fig.1-8	13
<b>Fig.1-10</b>	In a section posterior to that shown in Fig.1-9	14
<b>Fig.1-11</b>	Right clubfoot of a 19-week-old fetus sectioned in the sagittal plane through the ankle and the posterior and middle subtalar joints.	15
<b>Fig.1-12</b>	A- Photomicrograph of a clubfoot specimen showing ligamentous cells haphazardly arranged. B- Photomicrograph of a control foot.	16



<b>Fig.1-13</b>	A- In the clubfoot, the anterior portion of the calcaneus lies beneath the head of the talus. B- Lateral displacement (abduction) of the anterior portion of the calcaneus	19
<b>Fig.1-14</b>	A and B: Normal right foot of a 30-week-old fetus (Fig, A) and right club-foot of a 19-week-old fetus (Fig, B)	20
<b>Fig.1-15</b>	A and B: The main errors in the correction of the clubfoot.	22
<b>Fig.1-16</b>	Kite's error.	23
<b>Fig.2-1</b>	Examination of the child's gait.	38
<b>Fig 2-2</b>	Classification of clubfoot according to Dimeglio	41
<b>Fig 2-3</b>	Classification of clubfoot according to Pirani .	42
<b>Fig.2-4</b>	The knee, leg, and foot are placed in a vertical position.	48
<b>Fig.2-5</b>	Roentgenographic evaluation of clubfoot. Talocalcaneal angle on anteroposterior view of normal left foot.	49
<b>Fig.2-6</b>	Roentgenographic evaluation of clubfoot. Anteroposterior view of right clubfoot with decrease in talocalcaneal angle.	50
<b>Fig.2-7</b>	Roentgenographic evaluation of clubfoot. Talocalcaneal and tibiocalcaneal angles on dorsiflexion lateral view of normal left foot.	50
<b>Fig.2-8</b>	Roentgenographic evaluation of clubfoot. Talocalcaneal angle of zero and negative tibiocalcaneal angle on dorsiflexion lateral view of right clubfoot.	51
<b>Fig.2-9</b>	Lateral view in talipes equinovarus demonstrates an abnormally elevated tibio-calcaneal angle. A normal	

	angle is 60-90°.	52
<b>Fig.2-10</b>	Anteroposterior talus–first metatarsal angle.	
	Diagram showing the long axis of the talar head	53
<b>Fig. 2-11</b>		
	Diagram showing four patterns of talonavicular alignment	55
<b>Fig.2-12</b>		
	MRI of the right normal foot of a 12-month-old boy. A lateral shift of the navicular,	57
<b>Fig.2-13</b>		
	MRI of the left side in an 11-month-old boy with bilateral club feet. A medial shift of the navicular	58
<b>Fig.2-14</b>		
	A child with relapsed clubfoot treated with Ponseti technique.	58
<b>Fig.3-1</b>		
	A five-year-old child with recurrent right clubfoot, depicting preoperative deformity.	60
<b>Fig.3-2</b>		
	Clinicoradiological deformity correction of the patient after treatment with manipulation	61
<b>Fig.3-3</b>		
	The brace consists of open toe high-top straight last shoes attached to a bar	63
<b>Fig.3-4</b>		
	Steenbeek Foot Abduction Brace	65
<b>Fig.3-5</b>		
	Markell Brace – United States	72
<b>Fig.3-6</b>		
	John Mitchel Brace – United States	73
<b>Fig.3-7</b>		
	Gottenburg Brace – Sweden	73
<b>Fig.3-8</b>		
	Lyon Brace – France	74
<b>Fig.3-9</b>		
	The first incision is on the medial side of the foot over	

<b>Fig. 3-10</b>	the insertion of the anterior tibial tendon	74
	The second incision is at the anterior distal tibia	82
<b>Fig.3-11</b>	The third incision is over the lateral cuneiform.	
<b>Fig. 3-12</b>	A drill hole is made in the lateral cuneiform in a dorsal-to-plantar direction.	83
<b>Fig.3-13</b>		84
	Using two Keith needles, the sutures are passed through the drill hole in the lateral cuneiform	84
<b>Fig.3-14</b>		
	The distal end of the tendon is passed subcutaneously	85
<b>Fig.3-15</b>	A. Turco incision. B. Carroll incision. C. Cincinnati incision.	
<b>Fig.3-16</b>		86
	Carroll two-incision technique.	
<b>Fig.3-17</b>	V-Y advancement flap for closure of Cincinnati incision.	95
<b>Fig.3-18</b>	Posterior portion of the Cincinnati incision.	96
<b>Fig.3-19</b>	Lateral portion of Cincinnati incision,	97
<b>Fig.3-20</b>	Medial portion of Cincinnati incision, superficial.	99
<b>Fig.3-21</b>	A. Foot of the prone patient positioned for AP and lateral intraoperative x-rays.	99
<b>Fig.3-22</b>		102
	Subtalar rotation of the foot.	
<b>Fig.3-23</b>	The Cincinnati incision used for standard clubfoot release.	103
<b>Fig.3-24</b>		104
	Lateral column shortening options for residual adduction deformity in clubfoot.	106
<b>Fig.3-25</b>		

	Double tarsal osteotomy for residual adductus deformity.	110
<b>Fig.3-26</b>	A, B: Principle of calcaneal osteotomy according to Dwyer	
<b>Fig. 3-27</b>	The lateral approach for osteotomy of the os calcis.	111
<b>Fig.3-28</b>	Preferred longitudinal incisions for metatarsal osteotomies.	112
<b>Fig.3-29</b>	Radiograph of internal fixation used for metatarsal osteotomies.	114
<b>Fig.3-30</b>	Lambrinudi operation for talipes equinus.	116
<b>Fig.3-31</b>	AP x-rays of the lower leg 4 weeks after supramalleolar tibial derotation osteotomy	116
<b>Fig.3-32</b>	Arthrodesis for persistent or untreated clubfoot.	119
<b>Fig.3-33:</b>	Talectomy. A, Anterolateral skin incision. B, Total talectomy.	120
<b>Fig.3-34</b>	Anteroposterior (a) medial (b) and lateral (c) views of the frame.	122
<b>Fig.3-35</b>	Correction of clubfoot deformity with the Ilizarov apparatus	124
<b>Fig.3-36</b>	Photograph showing the cushioned slipper secured by velcro straps.	128
<b>Fig.3-37</b>	The foot before operation (a), during treatment (b) and at follow-up at three years (c and d).	128
<b>Fig.3-38</b>		130
		130

# List of Tables

<b>Table No.</b>	<b>Title</b>	<b>Page</b>
<b>Table 1:</b>	The rating system of Laaveg and Ponseti (1980)	43
<b>Table 2:</b>	Clinical details of patients treated with Ponseti method	63
<b>Table 3:</b>	Patients details treated with botulinium toxin A	76
<b>Table 4:</b>	Therapeutic strategy for resistant clubfoot.	77

## List of Abbreviation

AFO	Ankle foot orthosis
AP	Anteroposterior view
BTX-A	Botulinum toxin-A
CTEV	Congenital talipes equinovarus
DBB	Denis Brown Bar
FDL and FHL	Flexor digitorum longus and Flexor hallucis longus
HS	Hindfoot score
MR	Magnetic Resonance
MRI	Magnetic Resonance Imaging
MS	Midfoot score
SPLATT	Split anterior tendon transfer of tibialis anterior
TNA	Talar neck angle
TN	Talonavicular

## List of Figures

No. of figure	Title	Page
<b>Fig.1-1</b>	Bilateral clubfoot deformity in a 6-week-old infant.	3
<b>Fig.1-2</b>	Relapsed clubfoot of a 3-year-old boy.	5
<b>Fig.1-3</b>	Talus in clubfoot	6
<b>Fig.1-4</b>	The angle between the trochlea of the talus and the talar neck	6
<b>Fig.1-5</b>	3-D reconstruction of a CT scan of a clubfoot (9-years old boy).	9
<b>Fig.1-6</b>	Right clubfoot of 16-week-old fetus sectioned in the transverse plane through the calcaneocuboid joint.	9
<b>Fig.1-7</b>	A, B. Right clubfoot of a 16-week-old fetus, sectioned in the transverse plane through the talonavicular joint.	11
<b>Fig.1-8</b>	Sections in the frontal plane through the malleoli of the clubfoot.	13
<b>Fig.1-9</b>	A and B. In section posterior to those shown in Fig.1-8	13
<b>Fig.1-10</b>	In a section posterior to that shown in Fig.1-9	14
<b>Fig.1-11</b>	Right clubfoot of a 19-week-old fetus sectioned in the sagittal plane through the ankle and the posterior and middle subtalar joints.	15
<b>Fig.1-12</b>	A- Photomicrograph of a clubfoot specimen showing ligamentous cells haphazardly arranged. B- Photomicrograph of a control foot.	16