

Management of Traumatic Intracerebral Hematoma

Thesis

Submitted for Complete Fulfillment of the Master Degree in

General Surgery

By

Mohammed Hussein Mohammed Ali

M.B.B.Ch

Senior Registrar of Neurosurgery
Faculty of Medicine, Cairo University

Supervised by

Prof.Dr. Faheem Aly Elbassiony

Professor of General Surgery
Faculty of Medicine, Cairo University

Prof.Dr. Ibrahim Mohmmmed Ibrahim

Professor of Neurosurgery
Faculty of Medicine, Cairo University

Dr. Mohammed Abd Almonam Elmasry

Lecturer of General Surgery
Faculty of Medicine, Cairo University

Faculty of Medicine
Cairo University

2012

Abstract

In the present study, the effect of estrogen on gastric acid secretion and peptic ulcer and whether it acts through alpha or beta estrogen receptors has been investigated in a total number of 90 female albino rats divided into five groups with 18 rats allocated in each group.

Results showed that gastric acid secretion and number of ulcers were significantly higher among ovariectomized group than in control group when estimated within the basal acid secretion, carbachol induced acid secretion and indomethacin induced ulcer sub groups ($P<0.05$).

Key Words:

Gastric Acid Secretion, Gastrointestinal barriers, Peptic ulcer, Estrogens

List of Contents

Acknowledgment	
List of Tables	I
List of Figures	III
List of Abbreviations	V
Introduction and Aim of the Work	1
Review of Literature	
Gastric Acid Secretion	2
Gastrointestinal barriers	12
Peptic ulcer	43
Estrogens	53
Materials and Methods	65
Results	69
Discussion	106
Recommendations	120
Summary and Conclusion	121
References	125
Raw Data	i
Arabic Summary	-

List of Tables

Table No.	Title	Page No.
1	Comparison between control group and ovariectomized group in terms of gastric acid secretion and number of ulcers in different studied sub-groups	72
2	Comparison between control group and ovariectomized with estrogen replacement group in terms of gastric acid secretion and number of ulcers in different studied sub-groups	76
3	Comparison between control group and ovariectomized with alpha estrogen receptor agonist (PPT) group in terms of gastric acid secretion and number of ulcers in different studied sub-groups	80
4	Comparison between control group and ovariectomized with beta estrogen receptor agonist (DPN) group in terms of gastric acid secretion and number of ulcers in different studied sub-groups	85
5	Comparison between ovariectomized group and ovariectomized with estrogen replacement group in terms of gastric acid secretion and number of ulcers in different studied sub-groups	88
6	Comparison between ovariectomized group and ovariectomized with alpha estrogen receptor agonist (PPT) group in terms of gastric acid secretion and number of ulcers in different studied sub-groups	92

7	Comparison between ovariectomized group and ovariectomized with beta estrogen receptor agonist (DPN) group in terms of gastric acid secretion and number of ulcers in different studied sub-groups	96
8	Comparison between ovariectomized with estrogen replacement group ovariectomized with alpha estrogen receptor agonist (PPT) group in terms of gastric acid secretion and number of ulcers in different studied sub-groups	100
9	Comparison between ovariectomized with estrogen replacement group and ovariectomized with beta estrogen receptor agonist (DPN) group in terms of gastric acid secretion and number of ulcers in different studied sub-groups	104

List of Figures

Figure No.	Title	Page No.
1-A	Comparison between control group and ovariectomized group in terms of gastric acid secretion in different studied sub-groups	73
1-B	Comparison between control group and ovariectomized group in terms of number of ulcers in different studied sub-groups	73
2-A	Comparison between control group and ovariectomized with estrogen replacement group in terms of gastric acid secretion in different studied sub-groups	77
2-B	Comparison between control group and ovariectomized with estrogen replacement group in terms of number of ulcers in different studied sub-groups	77
3-A	Comparison between control group and ovariectomized with alpha estrogen receptor agonist (PPT) group in terms of gastric acid secretion in different studied sub-groups	81
3-B	Comparison between control group and ovariectomized with alpha estrogen receptor agonist (PPT) group in terms of number of ulcers in different studied sub-groups	81
4-A	Comparison between control group and ovariectomized with beta estrogen receptor agonist (DPN) group in terms of gastric acid secretion in different studied sub-groups	85
4-B	Comparison between control group and ovariectomized with beta estrogen receptor agonist (DPN) group in terms of number of ulcers in different studied sub-groups	85
5-A	Comparison between ovariectomized group and ovariectomized with estrogen replacement group in terms of gastric acid secretion in different studied sub-groups	89
5-B	Comparison between ovariectomized group and ovariectomized with estrogen replacement group in terms of number of ulcers in different studied sub-groups	89

6-A	Comparison between ovariectomized group and ovariectomized with alpha estrogen receptor agonist (PPT) group in terms of gastric acid secretion in different studied sub-groups	93
6-B	Comparison between ovariectomized group and ovariectomized with alpha estrogen receptor agonist (PPT) group in terms of number of ulcers in different studied sub-groups	93
7-A	Comparison between ovariectomized group and ovariectomized with beta estrogen receptor agonist (DPN) group in terms of gastric acid secretion in different studied sub-groups	97
7-B	Comparison between ovariectomized group and ovariectomized with beta estrogen receptor agonist (DPN) group in terms of number of ulcers in different studied sub-groups	97
8-A	Comparison between ovariectomized with estrogen replacement group ovariectomized with alpha estrogen receptor agonist (PPT) group in terms of gastric acid secretion in different studied sub-groups	101
8-B	Comparison between ovariectomized with estrogen replacement group ovariectomized with alpha estrogen receptor agonist (PPT) group in terms of number of ulcers in different studied sub-groups	101
9-A	Comparison between ovariectomized with estrogen replacement group and ovariectomized with beta estrogen receptor agonist (DPN) group in terms of gastric acid secretion in different studied sub-groups	105
9-B	Comparison between ovariectomized with estrogen replacement group and ovariectomized with beta estrogen receptor agonist (DPN) group in terms of number of ulcers in different studied sub-groups	105

List of Abbreviations

AE	Anion exchanger isoform
ANOVA	Analysis of variance
ANP	Atrial natriuretic peptide
cAMP	Cyclic adenosine mono phosphate
CFTR	cystic fibrosis transmembrane conductance regulator
cGMP	Cyclic guanosine mono phosphate
CGRP	Calcitonine gene related peptide
Cl	Chloride
CNQX	6-cyano-7-nitroquinoxaline-2,3-dione
CO ₂	Carbone dioxide
COMT	catechol O-methyl transferase
COX	Cyclooxygenase
Da	Dalton
DIDS	4,4'-diisothiocyanostilbene-2,2'-disulphonic acid
DMSO	Dimethyl sulphoxide
DNA	Deoxy riboneucleic acid
DPN	2, 3-bis (4-Hydroxyphenyl) Propionitrile
DRG	Dorsal Root Ganglions
ECL	Enterochromaffine like
EGF	Epidermal growth factor
EGFR	Epidermal growth factor receptor
EP receptors	E series of prostaglandins receptors
ER	Estrogen receptor
EST	Estradiol
EST	Estradiol
FSH	Follicular stimulating hormone
GABAA	g-amino-butyric acid A receptor
GIT	Gastro intestinal tract
GLP	Glucagon like peptide
GPR30	G protein-coupled receptor
GRP	gastrin-releasing peptide
GSK	Glycogen synthesis kinase
H.PYLORI	Helicobacter pylori
H ₂ O	Water
Hcl	Hydrochloric acid
HCO ₃	Bicarbonate
HDL	High density lipoproteins

HSV	High selective vagotomy
ICMJE	International Committee of Medical Journal Editors
IGF	Insulin like growth factor
IM	Intra muscular
IV	Intra venous
Kcl	Potassium chloride
LDL	Low density lipoproteins
LH	Luteinizing hormone
MAPK	mitogen-activated protein kinase
MBS	Mucosal bicarbonate secretion
Na cl	Sodium chloride
NBC	Sodium bicarbonate co transport
NMDA	N-methyl-D-aspartate
NO	Nitric oxide
NOS	Nitric oxide synthetase
NSAID	Non steroidal anti inflammatory drugs
OCD	Obsessive compulsive disorders
OV	Ovariectomy
OVX	Ovariectomized
OX1R	Orexin receptor one
OX2R	Orexin receptor two
PG	Prostaglandins
PI3K	phosphoinositide 3-kinase
PPT	1, 3, 5-tris (4-hydroxyphenyl)-4-propyl-1H-pyrazole
PUD	Peptic ulcer disease
RNA	Ribonucleic acid
SD	Standard deviation
TFF	Trefoil factor family
TRH	Thyrotropine releasing hormone
TRPV1	Transient receptor potential vanilloid receptor 1
USA	United states of America
VIP	Vasoactive intestinal peptide

Acknowledgement

*Thanks to **ALLAH** who have lightened my path to become a humble student of a noble profession, and granted me the ability to accomplish this work,*

*I would like to express my sincere gratitude and appreciation to **Prof.Dr.Mohamed Mahmoud Emran** Professor of Physiology Faculty of Medicine Cairo University & head of physiology department Misr University for Science and Technology, for his generous help, supervision and extreme kindness, spending much of his valuable time advising me.*

*I am also deeply grateful and would like to express my sincere thanks to my spiritual father **Prof.Dr.Akef Abdel-Haleem Khowaileed** Professor of Physiology Faculty of Medicine Cairo University, for his great help and his efforts that made this work come to light. Also I want to thank him for his continuous assistance and valuable guidance to me in my life.*

*I am greatly thankful and grateful to **Dr.Nashwa Mohamed Saied El-tablawy** Lecturer of Physiology Faculty of Medicine Cairo University, for her keen supervision, kind help and valuable instructions.*

*Great thanks and appreciation go to **Prof.Dr. Mohamed Hany Gamal El-Din**, professor of physiology, Faculty of Medicine, Cairo and MUST Universities for his sincer help.*

*Special appreciation and gratitude go to **Dr. Walid El-Sherbiny** and **Dr. Inas Abdel Rehim** for their great help and support.*

Lastly I am deeply grateful to everyone who has participated in this work,



قَالُوا سُبْحَانَكَ
لَا عِلْمَ لَنَا
إِلَّا مَا عَلَّمْتَنَا
أَنَّكَ أَنْتَ
الْعَلِيمُ الْحَكِيمُ

صرق الله العظيم

سورة البقرة

(الآية ٣٢)



*I would like to dedicate this thesis to my Father
& Sunshine Mother; to them I will never find
adequate words to express my gratitude.*

*Also to my beloved Brother and Sisters, for
dealing so patiently, tactfully during this work.*

Introduction

Originally identified as reproductive hormones, estrogens are now generally thought to play an important roles in bone, cardiovascular, gastrointestinal and in the central nervous system. This expanded view of estrogen action reflects the findings of a large number of clinical studies and huge body of empirical observational data gathered on the effect of exogenous estrogen administration on the health of menopausal women. Parallel studies reporting the distribution of estrogen receptors and others describing specific estrogenic responses in functionally related tissues have led to an appreciation of the sphere of influence of this hormone (*Mc Donnell, 2003*).

Aim of the work

This study aims to investigate the effect of estrogen on basal, charbacol stimulated acid secretion and also its effect on indomethacin induced ulcer and whether it acts through alpha or beta estrogen receptors.

Gastric Acid Secretion

Hydrochloric acid (HCl) is the best-known component of the parietal (oxyntic) cell secretion which is known as the gastric juice. The capacity of the stomach to secrete HCl is directly proportionate to the parietal cell number, which on stimulation secretes HCl at a concentration of roughly 160 millimol (mM)/liter that is equivalent to a pH of 0.8. Gastric acid facilitates the digestion of protein and absorption of iron, calcium, and vitamin B12, and prevents bacterial overgrowth and enteric infection (*Wei and Mitchell, 2006*).

Functions of Gastric Acid:

Hydrochloric acid (HCl) secretion assists protein digestion by activating pepsinogen to pepsin, renders the stomach sterile against orally-ingested pathogens, prevents bacterial or fungal overgrowth of the small intestine, encourages the flow of bile and pancreatic enzymes, and facilitates the absorption of a variety of nutrients, including folic acid, ascorbic acid, beta-carotene, non-heme iron, and some forms of calcium, magnesium, and zinc. Moreover normal gastric acid secretion is necessary for effective absorption of thyroxine (*Centanni et al 2006*).

Mechanism of Acid Secretion:

Hydrogen ions are generated within the parietal cell from electric dissociation of water. The hydroxyl ions formed in this process rapidly combine with carbon dioxide to form bicarbonate ion, a reaction catalyzed by carbonic anhydrase.

Bicarbonate is transported out of the basolateral membrane in exchange for chloride. The outflow of bicarbonate into blood results in a slight elevation of blood pH known as the "alkaline tide". This process serves to maintain intracellular pH in the parietal cell. Chloride and potassium ions are transported into the lumen of the canaliculus by conductance channels, and such is necessary for secretion of acid. Hydrogen ion is pumped out of the cell, into the lumen, in exchange for potassium through the action of the proton pump; potassium is thus effectively recycled. Accumulation of osmotically-active hydrogen ion in the canaliculus generates an osmotic gradient across the membrane that results in outward diffusion of water. The resulting gastric juice is 155 mM HCl and 15 mM KCl with a small amount of NaCl (*Yao and Forte, 2003*).

Phases of acid secretion:

The physiologic stimulation of acid secretion has classically been divided into three interrelated phases: cephalic, gastric, and intestinal. The cephalic phase is activated by the thought, taste, smell and sight of food, and swallowing. It is mediated mostly by cholinergic/vagal mechanisms. The gastric