

Ain Shams University, Women's College for Arts, Science and Education, Cairo- Egypt.

Effect of Temperature on The Removal of Some Heavy Elements by Friedel's Salt

A Thesis

Submitted to the Chemistry Department,
Women's College, Ain Shams University
In Partial Fulfillment of the Requirements for
the Degree of M. Sc. in Chemistry.

Presented by Heba Amine El-said Ahmed Sharaf (B.Sc., 2008)

Supervisors

Prof. Dr. Essam A. Kishar

Prof. of Inorganic Chemistry Women's College, Ain Shams University, Cairo, Egypt.

Prof. Dr. Tarek M. El-sokary

Prof. of Physical Chemistry, Housing and Building National Research Center, Cairo, Egypt.

Dr. Maha R. Mohamed

Ass. Prof. of Inorganic Chemistry Women's College, Ain Shams University, Cairo, Egypt.

Effect of Temperature on The Removal of Some Heavy Elements by Friedel's Salt

THESIS ADVISORS

APPROVED

Prof. Dr. Essam A. Kishar

Prof. of Inorganic Chemistry

Prof. Dr. Tarek M. El-sokary

Prof. of Chemistry in Housing and Building National Research Center

Ass. Prof. Dr. Maha R. Mohamed

Ass. Prof. of Inorganic Chemistry

APPROVED

Head of Chemistry Department

Prof. Dr. Mansoura Ismail

سم الله الرحمن الرحيح

الحمد لله الذي له ما في السموات وما في الأرض وله الحمد في الاخره وله الحدد في الاخره وهو وهو الحديم الخير

صدق الله العظيم

سوره سبأ الآبيه (١)



Ain Shams University, Women's College for Arts, Science and Education, Cairo- Egypt.

Student Name: Heba Amine El-said Ahmed Sharaf

Scientific Degree: B. Sc.

Department: Chemistry

Name of Faculty: Faculty of Girls

University: Ain Shams University

B. Sc. Graduation Date: 2008

ACKNOWLEGMENT

Praise and Thanks be to **ALLAH**, the most merciful for assisting and directing me the right way.

I would like to submit my gratitude, sincere thanks and appreciation to *Prof. Dr. Essam A. Kishar*, Professor of Inorganic Chemistry, Women's College, Ain Shams University, for suggesting the subject of this work, useful guidance, kind help, continuous interest, and fruitful discussion, which have facilitated the interpretation of the data throughout the chapters of this study as well as preparation the thesis in its final form.

I would like to express my deepest thanks to *Prof. Dr. Tarek M. El-sokary*, Prof. of Physical Chemistry in Housing and Building National Research Center, for his kind help, talented supervision and criticism, useful directions, valuable and fruitful discussion during all the steps of the study.

My deepest gratitude and appreciation to *Ass. Prof. Dr. Maha R. Mohamed*, Ass. Prof. of Inorganic Chemistry,
Women's College, Ain Shams University, for her deep concern
in this work, brilliance and effort in guiding and continuous
help throughout the whole work.

Thanks are also to all members of the Chemistry

Department, Women's College, Ain Shams University, for their

help.

Heba Amine

NOTE

Beside the work done in this thesis, the candidate has attended post -graduate courses for one year in inorganic and analytical chemistry including the following topics:

- Instrumental Analysis.
- Thermodynamic Chemistry.
- Kinetics and Catalysis.
- Nuclear Chemistry.
- Structural Inorganic Chemistry.
- Electrochemical Analysis.
- Special Topics.
- Spectroscopy.
- Coordination Chemistry.
- Photo Chemistry.
- Advanced Chemical Reactions.
- English Language.

She has successfully passed written examinations in the above mentioned topics.

Contents

	Page
Acknowledgement	
List of Tables	i
List of Figures	I
Abstract	a
<u>CHAPTER (I)</u>	
Introduction and Aim of the Work	
I. A. Literature Survey	1
I. A. 1. Friedel's salt composition	1
I. A. 2. Structure of Friedel's salt	2
I. A. 3. Preparation of Friedel's salt	9
I. A. 4. Stability and solubility of Friedel's salt	14
I.A.5. Friedel's salt as an adsorbent	16
I. A. 5.i. Adsorption of anions by Friedel's salt	17
I. A. 5.ii. Adsorption of cations by Friedel's salt	25
I. B. Aim of the work	29
CHAPTER (II)	
Materials, Techniques and Instruments	
II. A. 1. Materials	31
II.B. Techniques	31
II.B.1. Preparation of samples	31
II.B.1.i. Preparation of tricalcium aluminate, C ₃ A,	31

$(3\text{CaO.Al}_2\text{O}_3)$		
II.B.1.ii. Preparation of Friedel's salt		
II.B.2. Procedures	34	
II.B.2.i. Determination of the solubility and stability of		
Friedel's salt at different pH values:	34	
II.B.2.ii. Preparation of Friedel's salt metal complexes.	34	
II.B.2.iii. Determination of the effect of temperature on		
the adsorption kinetics of iron and cobalt on FS:	35	
II.C. Instrumentation.	35	
II.C.1. X-Ray Diffraction Analysis (XRD)	35	
II.C.2. Atomic Absorption (AA)	35	
II.C.3. Infrared spectroscopic measurements (IR)		
CHAPTER (III)		
Results and Discussion		
III. A. The solubility and stability of Friedel's salt at		
different pH values.	37	
III. B. Kinetics of adsorption of iron on Friedel's salt.	43	
III.B.1. Iron adsorption on Friedel's salt at room	42	
temperature.	43	
III.B.1.i. Adsorption of different concentrations of iron	42	
at pH=4.	43	
III.B.1.ii. Adsorption of different concentrations of iron		
.	55	

III.B.2. Effect of temperature on the adsorption of 0.5mM FeCl ₃ .6H ₂ O on Friedel's salt at pH 4.	
III.C. Kinetics of adsorption of cobalt on Friedel's salt.	
III.C.1. Cobalt adsorption at room temperature.	71
III.C.1.i. Adsorption of different concentrations of cobalt at pH=4.	71
III.C.2. Effect of temperature on adsorption of 0.5 mM of CoCl ₂ .2H ₂ O at pH=4 on Friedel's salt.	79
Summary and Conclusion	
References	
Arabic Summary	

List of Tables

Table No.	Title of Table	Page
Table (1)	The concentration of the released Ca ²⁺ in the solution at different pH's.	38
Table (2)	Table (2): Non-adsorbed and adsorbed iron ions in solutions having different initial concentrations at pH= 4 at different time intervals.	50
Table (3)	Non-adsorbed and adsorbed iron ions in solutions having different initial concentrations at pH= 6 at different time intervals.	61
Table (4)	The effect of temperature on adsorption of 0.5mM of iron at pH=4 at different time intervals.	68
Table (5)	Non-adsorbed and adsorbed cobalt ions in solutions having different initial concentrations at pH= 4 and at different time intervals.	77
Table (6)	The effect of temperature on adsorption of 0.5mM of cobalt at different time intervals at pH=4.	80

i

List of Figures

Figure No.	Title of Figure	Page
Figure (1)	Hydrogen environment of chloride anion: six H atoms from hydroxyl groups at 2.644Å and six H atoms (at maximum) at 2.478Å from water molecules, which are bonded to Ca ²⁺ cation.	4
Figure (2)	anapshot of the simulation of hydrocalumite / water interface at 25° C. Octahedral portray Al layers. Small spheres are Ca larger spheres are Cl.	7
Figure (3)	Interlayer structure of hydrocalumite (view down the c direction): (a) ordered structure from the X-ray diffraction data interpretation (b) a snapshot of the orientationally disordered structure resulting from the MD simulations at 0° C.	8
Figure (4)	Sevenfold-coordinated environment of a Ca atom in the hydroxide layer and the Ca- OH_2 electrostatic bonding in hydrocalumite.	9
Figure (5)	The reaction of NO ²⁻ with Friedel's salt.	25

ı

Figure (6)	XRD of the prepared C_3A .	32
Figure (7)	XRD of the prepared Friedel's salt.	33
Figure (8)	FTIR spectrum of prepared FS.	33
Figure (9)	Solubility of Friedel's salt at different pH values.	38
Figure (10)	X-ray diffraction patterns of FS at pH= 4.	40
Figure (11)	X-ray diffraction patterns of FS at pH=6.	41
Figure (12)	X-ray diffraction patterns of FS at pH= 8.	42
Figure (13)	XRD patterns of FS with 0.1 mM iron at pH= 4.	45
Figure (14)	XRD patterns of FS with 0.5 mM $FeCl_3.6H_2O$ at $pH=4$.	46
Figure (15)	XRD patterns of FS with 3 mM FeCl ₃ .6H ₂ O at pH= 4.	47
Figure (16)	XRD patterns of FS with 8 mM FeCl ₃ .6H ₂ O at pH= 4.	48
Figure (17)	Concentration of residual iron ions in solution at pH= (4) at different time intervals.	51
Figure (18)	FTIR of FS with 0.5 mM FeCl ₃ .6H ₂ O at pH= 4.	53

Figure (19)	FTIR of FS with 3 mM FeCl ₃ .6H ₂ O at pH=4.	53
Figure (20)	FTIR of FS with 8 mM FeCl ₃ .6H ₂ O at pH=4.	54
Figure (21)	XRD patterns of FS with 0.1 mM FeCl ₃ .6H ₂ O at pH= 6.	56
Figure (22)	XRD patterns of FS with 0.5 mM $FeCl_3.6H_2O$ at $pH=6$.	57
Figure (23)	XRD patterns of FS with 3 mM FeCl ₃ .6H ₂ O at pH= 6.	58
Figure (24)	XRD patterns of FS with 8 mM FeCl ₃ .6H ₂ O at pH= 6.	59
Figure (25)	Concentration of residual iron ions in solution at pH= 6 at different time intervals.	62
Figure (26)	FTIR of FS with 0.5 mM FeCl ₃ .6H ₂ O at pH= 6.	64
Figure (27)	FTIR of FS with 3 mM FeCl ₃ .6H ₂ O at pH=6.	64
Figure (28)	FTIR of FS with 8 mM FeCl ₃ .6H ₂ O at pH=6.	65
Figure (29)	XRD patterns of Fe treated FS at pH= 4 and 50°C.	67
Figure (30)	XRD patterns of the mix FS with 0.5 mM CoCl ₂ .2H ₂ O at pH= 4.	73
Figure (31)	XRD patterns of the mix FS with 3 mM CoCl ₂ .2H ₂ O at pH=4.	74

Figure (32)	XRD patterns of the mix FS with 8 mM CoCl ₂ .2H ₂ O at pH= 4.	75
Figure (33)	Concentration of residual cobalt ions at pH= 4 and at different time intervals.	78
Figure (34)	XRD patterns of Co- treated FS at pH= 4 and 50°C.	80
Figure (35)	The concentration of residual cobalt ions in solution after adsorption of CoCl ₂ .2H ₂ O at pH=4 at room temperature and at 50°C at different time intervals.	81