

PHYSIOLOGICAL AND BIOCHEMICAL STUDIES ON MORINGA SPECIES IN EGYPT

A THESIS

Submitted for the degree of Master of Science as a Partial Fulfillment for requirements of the Master of Science

Botany (Plant Physiology)

By Heba Mohamed El- Saied Faramawy

M. Sc. In Science (Botany) (2008)

Ain Shams University Faculty of Science Botany Department (2016)



PHYSIOLOGICAL AND BIOCHEMICAL STUDIES ON MORINGA SPECIES IN EGYPT

A THESIS

Submitted for the degree of Master of Science as a Partial Fulfillment for requirements of the Master of Science Botany (Plant Physiology)

By Heba Mohammed El- Saied Faramawy B. Sc. In Science (Botany) (2008)

Supervisors

Prof. Dr. Raifa Ahmed Hassanein

Prof. of Plant Physiology

Faculty of Science

Ain-Shams University

Dr. Amal Fadl Abd El-Kader

Assist. Prof. of Plant Physiology

Faculty of Science

Ain-Shams University

Ain Shams University Faculty of Science Botany Department (2016)



Approval Sheet

Title of Thesis:

Physiological and Biochemical Studies on *Moringa* Species in Egypt

Degree:

Master of Science in Botany (Physiology)

Name of Student:

Heba Mohammed El- Saied Faramawy

This Thesis has been approved by:

- 1- Prof. Dr
- 2- Prof. Dr.
- 2- Prof. Dr.

Prof. Dr. Maher Ahmed Shehata Head Department of Botany



M.Sc. Thesis

Name: Heba Mohamed El- Saied Faramawy

Title: "Physiological and Biochemical Studies on *Moringa*

Species in Egypt"

Degree: Master of Science (Botany, Physiology)

Supervisors:

Prof. Dr. Raifa Ahmed Hassanein

Prof. of Plant Physiology

Botany Department, Faculty of Science

Ain-Shams University

Dr. Amal Fadl Abd El-Kader

Assist. Prof. of Plant Physiology

Botany Department, Faculty of Science

Ain-Shams University

Prof. Dr. Maher Mohamed Shehata

Head of Botany Department Faculty of Science

Declaration

These thesis is the original work of the author and does not done previously at this degree or any other university. It was valuable and beneficial for me and I wish to be useful for other authors and researchers.

Heba Mohamed Faramawy



ACKNOWLEDGMENT

Firstly and Finally Thanks to Allah for helping me to finish this work.

I would like to express my special thanks and gratitude to **Prof.Dr. Raifa A. Hassanein**, Professor of Plant Physiology, Botany Department, Faculty of Science, Ain Shams University, For suggesting the point of research, supervision, continuous help, encouragement support, following up throughout the work.

Thanks are also to **Prof. Dr. Amal F. Abd El-Kader,** Assistant Professor of Plant Physiology, Botany Department, Faculty of Science, Ain Shams University, for his encouragement.

Special thanks for **Prof. Dr. Maher Shehata**, Head of Botany Department, Faculty of Science, Ain Shams University, for his encouragement and valuable help.

My thanks are also to all staff members in Botany Department, Faculty of Science, Ain Shams University.

My thanks to **Dr. Nesma Maher** Professor of Plant Physiology for their help and support throughout my work.

Special appreciation to **Dr. Hend Kamel** assistant Professor of Plant Ecology For help in Statistical Analysis

Special appreciation and sincere gratefulness is extended to my parents, my sisters, my brothers, my lovely husband and his continuous spiritual support, encouragement, sympathy, love and my sweety sons for their patience.

List of Tables	ı
List of Figures	V
List of Plates	X
List of photos	XIII
List of Abbreviation	XIV
Abstract	1
General Introduction and Aim of Work	4
Materials and Methods	40
Plant materials	40
Methods	40
Preparation of crude extract of <i>Moringa</i> leaves	40
The preparation of <i>Moringa</i> seed-cake	41
Extraction, separation and determination of growth regulating	41
substances	
Determination of the percentage of relative water content	45
(%RWC)	
Extraction and Estimation of Photosynthetic Pigments	45
Extraction and estimation of carbohydrates	46
Extraction and estimation of nitrogenous constituents	48

Extraction and estimation of lipid peroxidation product (MDA	53
content)	
Extraction and estimation of total reduced glutathione	54
Extraction and estimation of ascorbic acid	54
Extraction and estimation of anthocyanin content	55
Extraction and estimation of total phenols	56
Extraction and determination of flavonoids	57
Extraction and assaying activity of antioxidant enzymes	58
Extraction and estimation of protein using (SDS-PAGE)	64
Extraction and determination of certain elements	68
Chemical composition of fibers	69
Extraction and determination of oil content	71
Experimental Results	72
Biochemical Composition of Leaves of Moringa oleifera and	72
Moringa peregrina	
Phenolic Composition of <i>Moringa</i> Leaves	75
Protein Banding Patterns of Moringa Leaves	75
Experiment I:	80
Effect of Soil Treatment with Moringa Leaf Extract, (MLE) and/or Salinity	
on Growth Characters and some Metabolic Activities of Sweet Basil Plant	
(Ocimum basillcum)	

Experimental design and growth conditions	80
Changes in growth parameters	82
Changes in potosynthetic pigments and anthocyanin contents of leave	99
Changes in carbohydrate content	102
Changes in proline content	106
Changes in lipid peroxidation product (MDA)	107
Changes in antioxidant enzyme activities	110
Biochemical Composition of Fully Mature Seeds of Moringa oleifera and	130
Moringa peregrina Plants	
Phenolic compounds of Moringa seeds	131
Protein banding pattern of Moringa seeds	133
Experiment II:	
•	407
Mitigation of Cadmium Toxicity in Wheat Plant (Triticum asetivum L.)	137
Using the Coagulant Defatted Moringa oleifera or Moringa peregrina	
Seeds Powder	
Experimental design and growth conditions	137
Changes in growth parameters	138
Changes in relative water content	147
Changes in Photosynthetic Pigments	154
Change in The Cadmium Content	154
Protein Banding Pattern of Wheat Plants	157
Discussion	163
Summary	183

References	189
الملخص العربي	

LIST OF TABLES

Table number	Table title	Page number
1	Biochemical composition of fully expanded leaves of <i>Moringa oleifera</i> and <i>Moringa peregrina</i> plants	74
2	Phenolic compounds of leaves of <i>Moringa oleifera</i> and <i>Moringa peregrina</i> plants	76
3	Protein banding pattern of <i>Moringa oleifera</i> and <i>Moringa peregrina</i> leaves (molecular weight; Mw and amounts)	78
4	Effect of <i>Moringa oleifera</i> leaf extract (OLE) and <i>Moringa peregrina</i> leaf extract (PLE) on the growth parameters of sweet basil (<i>Ocimum basilicum</i> L.) plant grown under normal condition and/or salinity (S) stress. Each value is the mean of three variables. a) Shoot length b) Number of branches c) Shoot Fresh and Dry Weights d) Area of Leaves (cm²) e) Root Length (cm) f) Root Fresh and Dry Weights (g/plant)	84 87 90 93 96 100
5	Effect of <i>Moringa oleifera</i> leaf extract (OLE) and <i>Moringa</i> peregrina leaf extract (PLE) on photosynthetic pigments & anthocyanin content of leaves of sweet basil (<i>Ocimum basilicum</i>) plant grown under normal and salinity stress. Each value is the mean of three variables.	103

Ī

6	Effect of <i>Moringa oleifera</i> leaf extract (OLE) and <i>Moringa</i> peregrina leaf extract (PLE) on carbohydrate content (g/100g DW) of leaves sweet basil (<i>Ocimum basilicum</i>)	108
	plant grown under normal and salinity stress. Each value is the mean of three variables.	
7	Effect of <i>Moringa oleifera</i> leaf extract (OLE) and of <i>Moringa peregrine</i> leaf extract (PLE) on proline content (mg/g DW)of leaves and roots of sweet basil (<i>Ocimum basilicum</i>) plant grown under normal and salinity stress. Each value is the mean of three variables.	111
8	Effect of <i>Moringa oleifera</i> leaf extract (OLE) and <i>Moringa peregrina</i> leaf extract (PLE) on lipid peroxidation (MDA) content (mM/100g FW) of leaves and roots of sweet basil (<i>Ocimum basilicum</i>) plant grown under normal and salinity stress. Each value is the mean of three variables.	113
9	Effect of <i>Moringa oleifera</i> leaf extract (OLE) and <i>Moringa peregrina</i> leaf extract (PLE) on the activity of some antioxidant enzymes in leaves and roots of sweet basil (<i>Ocimum basilicum</i>) plant grown under normal and salinity stress condition. Each value is the mean of three variables. a) Superoxide dismutase (SOD) activity (unit/g Fw) b) Catalase Activity (CAT) (μM H ₂ O ₂ destroyed/g FW) c) Peroxidase activity (POX) d) Ascorbate peroxidase (APX) activity e) Poly phenole oxidase (ppo) activity f) Ascorbic acid oxidase (ASO) activity	118 120 122 124 126 128

10	Biohemical composition of fully matured seeds of <i>Moringa oleifera</i> and <i>Moringa peregrina</i> plants	132
11	Phenolic compounds of seeds of <i>Moringa oleifera</i> and <i>Moringa peregrina</i> plants	134
12	Protein banding pattern of <i>Moringa oleifera</i> and <i>Moringa peregrina</i> seeds (molecular weight; MWt and amounts)	135
13	Effect of soil application of defatted <i>Moringa oleifera</i> (OS) and <i>Moringa peregrina</i> (PS) seeds on growth parameter of 35 day old seedlings of wheat (<i>Triticum aestivum</i>) grown under Cd-acetate stress. Each value is the mean of three variables. A) Shoot Length (cm) B) Shoot Fresh and Dry Weights (mg/plant) C) Root Length (cm) D) Root Fresh and Dry Weights (mg/plant)	140 144 146 148
14	Effect of soil application of defatted <i>Moringa oleifera</i> (OS) and <i>Moringa peregrina</i> (PS) seeds on percentage of relative water content (% RWC) of leaves of 35 day old seedlings of wheat (<i>Triticum aestivum</i>) grown under Cdacetate stress. Each value is the mean of three variables.	152
15	Effect of soil application of defatted <i>Moringa oleifera</i> (OS) and <i>Moringa peregrina</i> (PS) seeds powder on photosynthetic pigments content of leaves (μg g ⁻¹ FW) of 30 day old seedlings of wheat (<i>Triticum aestivum</i>) grown under Cd- stress. Each value is the mean of three variables.	155

16	Effect of soil treatment of defatted <i>Moringa oleifera</i> seed powder (OS) and <i>Moringa peregrina</i> seed powder (PS) on Cd content of shoots and roots of 35 day old seedlings of wheat (<i>Triticum aestivum</i>) grown under Cd ²⁺ stress. Each value is the mean of three variables.	158
17	Protein banding pattern of leaves of wheat plant (<i>Triticum aestivum</i>) treated with cadmium alone or in combination with defatted seed powder of <i>Moringa oleifera</i> (OS) or <i>Moringa peregrina</i> (PS), (MWt & amount).	161