



شبكة المعلومات الجامعية

بسم الله الرحمن الرحيم



شبكة المعلومات الجامعية
@ ASUNET



شبكة المعلومات الجامعية التوثيق الالكتروني والميكرو فيلم



شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الالكتروني والميكرو فيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأفلام قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيدا عن الغبار

في درجة حرارة من ١٥-٢٥ مئوية ورطوبة نسبية من ٢٠-٤٠%

To be Kept away from Dust in Dry Cool place of
15-25- c and relative humidity 20-40%

بعض الوثائق الأصلية تالفة

بالرسالة صفحات لم ترد بالاصل

RECENT ADVANCES IN RIGID GAS PERMEABLE CONTACT LENSES

*Essay Submitted for Partial Fulfillment of the
Master Degree in Ophthalmology*

B5623

By
Inas Hassan Hashem

*M.B., B.Ch.
Ain Shams University*

Supervised by
Prof. Dr. Fatma M. S. El Hennawy

*Professor of Ophthalmology
Faculty of Medicine - Ain Shams University*

Dr. Mamdouh H. El Kafrawy

*Assistant Professor of Ophthalmology
Faculty of Medicine - Ain Shams University*

**Faculty of Medicine
Ain Shams University
Cairo
2000**

میں نے
اپنے
میں

میں نے
میں نے
میں نے

۲۲

بسم الله الرحمن الرحيم

قالوا سبحانك لا علم لنا إلا ما علمتنا،
إنك أنت العليم الحكيم

صدق الله العظيم

سورة البقرة- الآية: ٣٢

17

18

19

20

21

22

23

Acknowledgment

I would like to express my respect and gratitude to Prof. Dr. Fatma M. S. El Hennawy, Professor of Ophthalmology, Faculty of Medicine, Ain Shams University, for her continuous support and meticulous supervision, without her great help and advices, this work could not be completed.

Also I present my thanks and appreciations to Dr. Mamdouh H. El Kafrawy, Assistant Professor of Ophthalmology, Faculty of Medicine - Ain Shams University, for his fruitful advices, continuous encouragement and worthy remarks.

Last but not least, I would like to present my sincere thanks to all staff members of the Ophthalmology Department, Faculty of Medicine, Ain Shams University, for their help and encouragement during this work.

1000000000

1000000000

1000000000

1000000000

1000000000

1000000000

1000000000

List of Contents

	Page
Introduction	1
Aim of the Work	6
RGP Materials	7
Parameters and Design	16
Corneal Topography	28
Indications of RGP Contact Lenses	43
Relative Contraindications for RGP Contact Lenses	57
Fitting of Rigid Gas Permeable (RGP) Contact Lenses	59
Current RGP Contact Lens Care Systems	95
Complications	103
Summary	126
References	128
Arabic Summary	

8000

11

8

20

20

20

20

20

20

20

20

20

20

20

20

List of Figures

	<i>Page</i>
Fig. (1): The contact angle.	11
Fig. (2): Tricurve lens showing intermediate and peripheral radii.	18
Fig. (3a): A spheric design.	22
Fig. (3b): Posterior surface of a spherical lens has a series of secondary curves that must be blended at each junction and flatten toward the lens periphery (top). The posterior surface of the aspheric Boston Envision lens flattens without any junctions that interfere with comfort or disturb corneal integrity (bottom).	23
Fig. (4): A high plus lenticular lens (Right) contrasted with a standard single cut plus lens (Left).	26
Fig. (5): Top: Photokeratoscope Bottom: appearance of normal placido image.	29
Fig. (6): Keratotomy showing colour coded topographical map of normal cornea.	31
Fig. (7): Normal topographic patterns.	33
Fig. (8): Astigmatism is seen on keratotomy as a distortion of the circular projections into concentric oval reflections.	34
Fig. (9): The line of sight is the straight line from the fixation point to the center of the entrance pupil (E). Current videokeratoscopes align on a reflected corneal image of a target that is symmetric with the videokeratoscope optic axis, which is detected toward the center of curvature (C) for the corneal position measured. Videokeratotomy axis deviates from where the line of sight crosses the cornea.	36
Fig. (10): Computerized stimulated fluorescein pattern showing an aligned spherical corneal lens.	40
Fig. (11): Computerized stimulated fluorescein pattern showing a steep fit spherical lens.	41
Fig. (12): A computerized stimulated fluorescein pattern showing a flat fit spherical lens.	42

Fig. (13a): Corneal topography demonstrates the steep irregular astigmatism of the cone with its peak in the inferior cornea below fixation.	47
Fig. (13b): The cone is completely eccentric and acuity is reduced to 20/200.	48
Fig. (14): Corneal mapping in central keratoconus showing irregularity.	49
Fig. (15): Relative size and shape of three cone forms in keratoconus (A) nipple shaped cone, (B) an oval cone, (C) globus cone	51
Fig. (16): Lenticulation.	66
Fig. (17): Top Rt: Lid attachment fit with the lens resting in superior position. Bottom Rt: Interpalpebral fit with centralizing lens. Top Lt: Inferior lens position. Bottom Lt: Temporal ride of aphakic lens.	69
Fig. (18): In this photograph, a dark area can be seen centrally, surrounded by bright green. The dark zone indicated that there is moderate apical bearing caused by a fitting relationship that is too flat.	73
Fig. (19): This photograph shows an evenly aligned edge-to-edge fluorescein pattern with slight bearing in the mid-periphery of about 45° on each side in the horizontal meridian. Mid-peripheral bearing must be kept to less than 180° to prevent seal-off and ensure an adequate tear exchange. This lens meets the criteria for an ideal fit	74
Fig. (20): The slight pooling of fluorescein centrally indicates apical clearance.	75
Fig. (21): In this lens-cornea relationship, severe apical clearance can be seen, as evidenced by substantial central pooling of fluorescein and 360° of heavy mid-peripheral bearing.	76
Fig. (22): A dark zone can be seen centrally with a moderately large area of central touch, causing severe apical bearing. The fit is so flat that it causes inferior standoff with pooling of fluorescein.	77