





ثبكة المعلومات الجامعية





جامعة عين شمس

التوثيق الالكتروني والميكروفيلم



نقسم بللله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأفلام قد اعدت دون آية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15-20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of 15-25c and relative humidity 20-40 %



ثبكة المعلومات الجامعية







PHYSIOLOGICAL STUDIES ON CITRUS ROOTSTOCK SEEDLINGS

By

ALAA AHMED MOKHTAR EL-ATEEK

A Thesis Submitted in Partial Fulfillment of The Requirements for the Degree of

Master of Science

In

Agriculture science (Pomology)

Department of Horticulture
Faculty of Agriculture, Moshtohor
Zagazig University, Benha Branch

3

 $\Lambda \subset \Lambda$

2003

PHYSIOLOGICAL STUDIES ON CITRUS ROOTSTOCK SEEDLINGS

Ву

ALAA AHMED MOKHTAR EL-ATEEK

B. Sc. Agric. Sci., 1982

Under the Supervision of

PROF. DR.

MOHAMED ABD - EL WAHAB KHAMIS

Prof. of Pomology

DR. KHALED ALI BAKRY

Lecture of Pomology

Department of Horticulture
Faculty of Agriculture, Moshtohor
Zagazig University
Benha Branch

2003

APPROVAL SHEET

PHYSIOLOGICAL STUDIES ON CITRUS ROOTSTOCK SEEDLINGS

By

ALAA AHMED MOKHTAR EL-ATEEK

B. Sc. Agric. Sci., 1982

This thesis for M. Sc. Degree has been Approved by:

Prof. Dr.: M. Fahmy

Prof. Dr.: M.A.Khan

Prof. Dr.: Thanks II. heraf

Date of Examination: / /2003

CONTENTS

	Page
I – INTRODUCTION	1
II – REVIEW OF LITERATURE	3
III- MATERIALS AND METHODS	37
IV- RESULTS AND DISCUSSION	44
IV.1. Vegetative growth	44
IV.1.1. Stem length	45
IV.1.2. Root length	48
IV.1.3. Plant height	49
IV.1.4. Stem diameter	53
IV.1.5. Number of leaves per plant	55
IV.1.6.Leaf area	57
IV.1.7. Number of roots per plant	61
IV.1.8. Top fresh weight	63
IV.1.9. Root fresh weight	66
IV.1.10. Total plant fresh weight	68
IV.1.11. Top dry weight	72
IV.1.12. Root dry weight	74
IV.1.13. Total plant dry weight	78
IV.1.14. Top /root ratio	79
IV.2. Effect of citrus rootstock kind, effective	
microorganism (EM) levels and method of	,
application on leaf chemical content of three	
citrus rootstocks seedlings	83
IV.2.1. Determination on leaf pigments	83
IV.2.1.1. Leaf chlorophyll "A" content	83
IV.2.1.2. leaf chlorophyll "B" content	86
IV.2.1.3. leaf carotenoids content	88
IV.2.2. Leaf mineral content	90
IV.2.2.1. Leaf "N" content	90
IV.2.2.2. Leaf phosphorus content	92

IV.2.2.3. Leaf potassium content	95
IV.2.2.4. Leaf calcium content	99
IV.2.2.5. Leaf magnesium content	102
IV.2.2.6. Leaf Iron content	103
IV.2.2.7. Leaf manganese content	108
IV.2.2.8. Zinc content	109
V- SUMMARY AND CONLUSION	114
VI -RFERENCES	120
VII – ARABIC SUMMARY	

LIST OF TABLE

Table (1)	Preparing 100 liter of different concentration EM solution.	40
Table (2)	Stem length of citrus rootstock seedlings as effected by	47
	specific effect of rootstock species; level, application	
	method of effective Microorganism (EM) and	
	interaction effect of their combinations during 1999 –	
	2000 and 2000 – 2001 seasons.	
		50
Table (3)	Root length of citrus rootstock seedlings as effected by	50
	specific effect of rootstock species; level, application	
	method of effective Microorganism (EM) and	
	interaction effect of their combinations during 1999 -	
	2000 and 2000 - 2001 seasons.	
Table (4)	Plant height of citrus rootstock seedlings as effected by	52
	specific effect of rootstock species; level, application	
	method of effective Microorganism (EM) and	
1	interaction effect of their combinations during 1999 –	
	2000 and 2000 – 2001 seasons.	
Table (5)	Stem diameter of citrus rootstock seedlings as effected	54
Table (3)	by specific effect of rootstock species; level,	
	application method of effective Microorganism (EM)	
	••	
	and interaction effect of their combinations during	
	1999 – 2000 and 2000 – 2001 seasons.	50
Table (6)	Number of leaves/plant of citrus rootstock seedlings as	5,8
	effected by specific effect of rootstock species; level,	
	application method of effective Microorganism (EM)	
	and interaction effect of their combinations during	
	1999 – 2000 and 2000 – 2001 seasons.	

Table (7)	Leaf area of citrus rootstock seedlings as effected by	60
	specific effect of rootstock species; level, application	
	method of effective Microorganism (EM) and	
	interaction effect of their combinations during 1999 -	
	2000 and 2000 - 2001 seasons.	
Table (8)	Number of roots/plant of citrus rootstock seedlings as	62
	effected by specific effect of rootstock species; level,	
	application method of effective Microorganism (EM)	
	and interaction effect of their combinations during	
	1999 2000 and 2000 2001 seasons.	
Table (9)	Top fresh weight of citrus rootstock seedlings as	65
	effected by specific effect of rootstock species; level,	
	application method of effective Microorganism (EM)	
	and interaction effect of their combinations during	
	1999 - 2000 and 2000 - 2001 seasons.	
Table (10)	Root fresh weight of citrus rootstock seedlings as	70
	effected by specific effect of rootstock species; level,	
	application method of effective Microorganism (EM)	
	and interaction effect of their combinations during	
	1999 – 2000 and 2000 – 2001 seasons.	
Table (11)	Total plant fresh weight of citrus rootstock seedlings as	71
	effected by specific effect of rootstock species; level,	
	application method of effective Microorganism (EM)	
	and interaction effect of their combinations during	
	1999 – 2000 and 2000 – 2001 seasons.	
Table (12)	Top dry weight of citrus rootstock seedlings as	76
	effected by specific effect of rootstock species; level,	
	application method of effective Microorganism (EM)	
	and interaction effect of their combinations during	
	1999 – 2000 and 2000 – 2001 seasons.	