

**REVISION OF EGYPTIAN HERBARIUM SPECIMENS
OF THE MOSS GENUS FISSIDENS HEDW.**

A Thesis

Submitted to the Faculty of Science
Ain Shams University

In Partial Fulfilment
of the Requirements for the Degree of
Master of Science

In
Botany

By

Samar Nour-El-Deen Mahmoud Nour-EL-Deen

B.Sc. Ain Shams University, special degree in Botany

Supervised by

Prof. Wagieh El-Sayed El-Saadawi

Prof. of Botany and ex-head of Botany Department
Faculty of Science
Ain Shams University

Prof. Hanaa Moustafa Shabbara

Prof. of Botany
Faculty of Science
Ain Shams University

2008

**REVISION OF EGYPTIAN HERBARIUM SPECIMENS
OF THE MOSS GENUS FISSIDENS HEDW.**

A Thesis
Submitted In Partial Fulfilment
of the Requirements for the Degree of
Master of Science
In
Botany

By
Samar Nour-El-Deen Mahmoud Nour-EL-Deen
B.Sc. Ain Shams University, special degree in Botany

Faculty of Science
Ain Shams University
2008

Examined by

Prof Mohamed Ahmed Hamouda

Prof of Botany

Faculty of Girls

Ain Shams University, Egypt

Prof Nady Ahmed El-Bassuony Ghanem

Prof of Ecology and Cryptogamic Plants

Faculty of Science (For Boys)

Al-Azhar University, Nasr City, Egypt

Prof. Wagieh El-Sayed El-Saadawi

Prof. of Botany and ex-head of Botany Dept.

Faculty of Science

Ain Shams University

Prof. Hanaa Moustafa Shabbara

Prof. of Botany

Faculty of Science

Ain Shams University

This thesis entitled:
Revision of Egyptian Herbarium Specimens
of the Moss Genus *Fissidens* Hedw.

written by: Samar Nour-El-Deen Mahmoud Nour-El-Deen
has been approved for Botany Department, Ain Shams University

Date

Date

Date

Date

The final copy of this thesis has been examined by the signatories, and we find that both the content and the form meet acceptable presentation standards of scholarly work in the above mentioned discipline.

I hereby declare that the work contained in this thesis, now submitted for the degree of Master of Science (Botany) to Ain Shams University, is the result of my own investigations.

_____ Candidate (Samar Nour-El-Deen)

_____ Supervisor (Wageih El-Saadawi)

_____ Supervisor (Hanaa Shabbara)

I hereby certify that the work embodied in this thesis has not already been accepted in substance for any degree, and is not currently being submitted for any other degree.

_____ Candidate (Samar Nour-El-Deen)

I dedicate this thesis to the most precious personalities of my life, my parents, without whom I would literally not be here—my mother ‘Nerman’ and my father ‘Nour-El-Deen’, whose number one question “Is she ever going to finish her MSc thesis?” can now be answered. I owe you all a debt of gratitude that can never be repaid for all the support you given me. To my sister ‘Sahar’, for her prayers and for pushing when I thought I could not be pushed anymore. To my brother ‘Mohammad’, who has always reassured and encouraged me in pursuit of my endeavors. To my little sweeties ‘Yousuf’ and ‘Mohammad’, who illuminated my days with their smiles. Without all of you I could not have reached this far. If I am looking a little taller, it is because I am standing on the shoulders of all of you and I owe you a lot.

Contents

List of Figures	v
List of Tables	xiii
Acknowledgements	xv
Abstract	xix
Preface	xxi
List of Abbreviations	xxiii
I General Part	1
1 Introduction	3
1.1 Bryophytes in Egypt	3
1.2 The Aim of This Thesis	6
1.3 History of Fissidens in Egypt	8
2 Materials and Methods	13
2.1 Materials	13
2.1.1 Specimens studied	13
2.1.2 Geographic area covered in the present study	22
2.2 Methods	30
2.2.1 General preparation of specimens	30
2.2.2 Cleaning of herbarium specimens	30
2.2.3 Permanent slide preparation	32
2.2.4 Staining	33
2.2.5 Illustrations and measurements	34
2.2.6 Distribution maps	35

3	General Considerations of Fissidens	37
3.1	Family Fissidentaceae	37
3.1.1	Conspectus of the genus Fissidens	38
3.1.2	Classification of the genus Fissidens	39
3.2	Fissidens in Egypt	40
3.2.1	Subgenus Fissidens section Fissidens	40
3.2.2	Description of the genus Fissidens	40
3.2.3	Notes on morphology and terminology used in de- scription	42
II	Special part	47
4	Floristic Treatment	49
4.1	Identification of Egyptian Specimens	49
4.1.1	Characters of value in delimiting Fissidens species	50
4.2	Key to the Egyptian Species of the Genus Fissidens	51
4.3	Fully-Identified Taxa	53
4.3.1	Fissidens crassipes subsp warnstorffii	53
4.3.2	Fissidens leucocinctus	89
4.3.3	Fissidens obtusifolius	107
4.4	Insufficiently Studied Species	123
4.4.1	Fissidens sp.1	123
4.4.2	Fissidens sp.2	133
4.5	Excludenda	143
4.6	Comment on a synonym included in the 1999's updated list	147
4.7	Dubious record	147
5	General Discussion	149
5.1	Conclusions and Challenges For the Future	153
	Literature Cited	157
	Summary	167
	Appendices	171
A	Section Cutting	173
B	Gametangia Arise on Long Stalks	181
B.1	Observations	182
B.2	Discussion	183

Literature Cited	191
C Chytrid Galls on Fissidens Rhizoids	193
C.1 Observations and Discussion	194
C.1.1 Fungus identification and taxonomy of chytrids	197
C.1.2 Morphology of the infected host	198
Literature Cited	207
Glossary	209

List of Figures

1.1	Map of Lower Egypt showing the place of Heliopolis city (ancient Egypt).	9
2.1	Map showing phytogeographical territories in Egypt.	23
2.2	Map showing the geographic area covered in the present study.	25
2.3	Map showing the provinces in Nile Valley where herbarium specimens were collected.	26
2.4	Map showing collection sites of the studied herbarium specimens collected from Nile Delta.	27
2.5	Map showing collection sites of the studied herbarium specimens collected from El-Faiyum province.	28
2.6	Map showing collection sites of the studied herbarium specimens collected from El-Sharqiyah province.	29
3.1	Different zones of ornamentation of peristome teeth (<i>bryoides</i> -type). . .	45
4.1	<i>Adiantum capillus-veneris</i> L.	57
4.2	Map showing the approximate extent of <i>Fissidens crassipes</i> subsp <i>warnstorffii</i> (M. Fleisch.) Brugg.-Nann. in Egypt.	58
4.3	Map showing floristic regions of the mosses in the World.	59
4.4	<i>Fissidens crassipes</i> subsp <i>warnstorffii</i> (M. Fleisch.) Brugg-Nann. Anatomical abnormalities in perichaetial leaves.	77
4.5	<i>Fissidens crassipes</i> subsp <i>warnstorffii</i> (M. Fleisch.) Brugg-Nann. (a)–(g) Parts of stems showing aerial rhizoids.	78
4.6	<i>Fissidens crassipes</i> subsp <i>warnstorffii</i> (M. Fleisch.) Brugg-Nann. Habit. (a)–(g) Unbranched plants; (h), (i) stems denuded of leaves showing axillary rhizoids in clusters; (j)–(o) branched plants; (p)–(s) plants branched from older parts.	79

- 4.7 *Fissidens crassipes* subsp *warnstorffii* (M. Fleisch.) Brugg-Nann. (a)–(e) Infertile leaves; (f)–(i) fertile leaves; (j), (k) areolation at leaf apex; (l), (m) areolation at middle of dorsal lamina; (n) costa forked; (o), (p) cells at basal part of leaf; (q) axillary hyaline nodule; (q') close-up view of (q). 80
- 4.8 *Fissidens crassipes* subsp *warnstorffii* (M. Fleisch.) Brugg-Nann. (a)–(c) Intralaminar limbidia on vaginant laminae; (d) marginal limbidia; (e), (f) cross-section of distal part of leaf; (g), (h) cross-section of distal part of leaf passed across the apex of minor lamina; (i)–(k) cross-section of proximal part of leaf; (l) cross-section of costa at proximal part of leaf; (m) cross-section of stem at caulogenous branch; (n)–(o) cross-sections of stem. 81
- 4.9 *Fissidens crassipes* subsp *warnstorffii* (M. Fleisch.) Brugg-Nann. Arrows points to caulogenous branches. (a), (c)–(e) caulogenous branches of indeterminate growth (not ended with gametangia); (f) stem was denuded of leaves to show distribution of gametangia along stem and caulogenous branch; (b), (g)–(o) caulogenous branches of determinate growth (ended with gametangia or sporophyte). 82
- 4.10 *Fissidens crassipes* subsp *warnstorffii* (M. Fleisch.) Brugg-Nann. (a)–(d) Reduced caulogenous branches ended with sporophyte, (c'), (d') close-up views of (c), (d), respectively; (e) archegonia in axil of reduced caulogenous branch, stem was denuded of leaves; (e') close-up view of (e); (f)–(i) caulogenous branches associated with gametangia; (j) naked solitary archegonium, stem was denuded of leaves. 83
- 4.11 *Fissidens crassipes* subsp *warnstorffii* (M. Fleisch.) Brugg-Nann. (a)–(d) Fertile branches; (e)–(h) elongated fertile branches; (i)–(q) variation of sporophyte. 84
- 4.12 *Fissidens crassipes* subsp *warnstorffii* (M. Fleisch.) Brugg-Nann. (a)–(c) Polysetous plants, (a'), (a'') close-up views of (a); (d)–(h) variations of shape of operculum; (i)–(o) cucullate calyptra; (p) outer surface of capsule; (q) surface view of exothecial cells; (r), (s) stomata; (t) peristome teeth; (u), (v) spores. 85
- 4.13 *Fissidens crassipes* subsp *warnstorffii* (M. Fleisch.) Brugg-Nann. (a)–(m) Lateral sporophyte, (f'), (g') close-up views of (f), (g), respectively; (n)–(q) disetous plants; (r)–(t) multiple fertilization. 86
- 4.14 *Fissidens crassipes* subsp *warnstorffii* (M. Fleisch.) Brugg-Nann. (a)–(g) Long foliar filamentous gemmae; (h)–(l) gemmae produced on rhizoids; (m)–(p) close-up views of rhizoidal gemmae, arrow in (n) points to an algal filament. 87

- 4.15 *Fissidens crassipes* subsp *warnstorffii* (M. Fleisch.) Brugg-Nann. (a)–(c) Rhizoidal tubers; (d), (e) germinated tubers; (f) dormant rhizoidal tuber adjacent to terminal sporophyte, arrow points to an aborted sporophyte; (g), (h) close-up views of the tuber; (i) stem denuded of leaves to show axillary chloronemata; (j) deciduous stem apex; (k) fertile stem denuded of leaves showing young gametophytes emerging from the vaginula; (l) close-up view of (k). 88
- 4.16 Map showing the approximate extent of *Fissidens leucocinctus* Hampe in Egypt. 91
- 4.17 Map of the African continent showing the ten countries which make up the Nile river basin and the African distribution of *Fissidens leucocinctus* Hampe. 93
- 4.18 *Fissidens leucocinctus* Hampe. (a) Population site, (b) fertile plant bearing two sporophytes; (c) intact capsule; (d) calyptra; (e) operculum; (f) peristome teeth. All photos by Lashin (1990). 98
- 4.19 *Fissidens leucocinctus* Hampe. (a)–(l) Variations of plant size and branching pattern. (g) plant underwent growth resumption; (l) branches from older parts; (m) arrow points to aerial rhizoid. 101
- 4.20 *Fissidens leucocinctus* Hampe. (a) Arrow points to a germinated rhizoidal bud; (b) arrows indicate young branches on older parts; (c) arrow points to a just separated deciduous stem apex; (d) close-up view of (c); (e)–(g) deciduous stem apices; (h) arrow points to deciduous stem apex developing new shoot; (i)–(n) germinated shoots, different developmental stages. 102
- 4.21 *Fissidens leucocinctus* Hampe. (a)–(j) Variations of leaf shape; (k) leaf apex; (l) middle of dorsal lamina; (m)–(p) variations of dorsal lamina; (q) arrows point to axillary hyaline nodules, stem was denuded of leaves; (r) close-up view of a hyaline nodule. 103
- 4.22 *Fissidens leucocinctus* Hampe. (a) Cross-section of distal part of leaf; (b)–(d) cross-section of distal part of leaf passed across the apex of minor lamina; (e)–(i) cross-section of proximal part of leaf; (j) cross-section of leaf showing recurved dorsal lamina; (k), (l) bistratose laminar cells; (m) bistratose juxtacostal cells; (n), (o) cross-section of costa at proximal part of leaf; (p) cross-section of costa at distal part of leaf; (q)–(s) cross-section in pluristratose limbidial cells of vaginant laminar cells; (t)–(y) cross-sections of stem. 104
- 4.23 *Fissidens leucocinctus* Hampe. (a)–(g) Areolations at leaf apex and variations of costa extension, (a'), (b') lower portions of (a) and (b), respectively, showing costa ending far below apex; (h)–(j) costa extending into leaf apex ended by a single large cell. 105

-
- 4.24 *Fissidens leucocinctus* Hampe. (a) Areolation at middle of dorsal lamina; (b)–(g) variations of limbidia development on dorsal lamina; (c), (d) intralaminar limbidia. 106
- 4.25 *Fissidens leucocinctus* Hampe. (a), (b) Cross-section of stem with part of leaf; (c)–(f) cross-section of stem through a hyaline nodule. 106
- 4.26 Map showing the Egyptian distribution of *F. obtusifolius* Wilson. 110
- 4.27 *Fissidens obtusifolius* Wilson. (a)–(f) Habit, (a)–(d) branched, (e), (f) unbranched; (g) plant with seta; (h) plant with sporophyte. All photos were taken of the specimen WE70 (CAIA), except (a), (b) of the specimen WE69 (CAIA). 117
- 4.28 *Fissidens obtusifolius* Wilson. (a)–(c) Stem leaves; (d), (e) perichaetial leaves; (f) perichaetial leaf with archegonia; (g) perigonal leaf with antheridia; (h) perigonal leaf; (i) cells at margin of vaginant laminae. All photos were taken of the specimen WE70 (CAIA). 118
- 4.29 *Fissidens obtusifolius* Wilson. (a) Cells at leaf apex; (b) cells at dorsal lamina, apex of minor lamina; (c) leaf insertion; (d) cells at mid-dorsal lamina; (e) cross-section of proximal part of leaf; (f) cross-section of costa (enlarged); (g) cross-section of distal part of leaf; (h), (i) cross-sections of stem. All photos were taken of the specimen WE70 (CAIA). 119
- 4.30 *Fissidens obtusifolius* Wilson. (a), (b) Female branches from base of stem; (c) upper part of stem showing distribution of female inflorescence (terminal and axillary), stem was denuded of leaves; (d), (e) terminal archegonia (enlarged); (f) male branch with terminal antheridium. All photos were taken of the specimen WE70 (CAIA). 120
- 4.31 *Fissidens obtusifolius* Wilson. (a) Exothecial cells, arrows point to stomata; (b) exothecial cells (enlarged); (c) spore sac with spores; (d), (e) spores (enlarged); (f) basal part of peristome teeth; (g) basal part (undivided) of a broken tooth (enlarged). All photos were taken of the specimen WE70 (CAIA), one of the intact capsules was dissected and used to prepare the illustrations. 121
- 4.32 *Fissidens obtusifolius* Wilson. Rhizoidal tubers. (a) young pale yellowish brown tuber on basal rhizoid; (b) mature dark red tubers, arrow points to a short rhizoidal stalk; (c)–(f) some rhizoidal tubers, arrows point to germinated tubers; (g), (h) tubers on rhizoidal stalks, arrows point to irregular shapes of tubers; (i)–(k) germinated tubers, different developmental stages. All photos were taken of the specimen WE69 (CAIA), except (a), (b), (d), (g) of the specimen WE70 (CAIA). 122

- 4.33 *Fissidens* sp1. (a)–(c) Habit, unbranched; (d) habit, branched from base; (e)–(g) branched from older parts; (h)–(k) terminal sporophyte of different sizes; (l), (m) sporophyte in subterminal positions; (n) intact sporophyte; (o) two sporophytes developed on one stem, arrow points to the incomplete one in subterminal position. 127
- 4.34 *Fissidens* sp1. (a)–(f) Variations of apex of minor lamina, dorsal lamina extension, and shape of stem leaves; (g)–(i) perichaetial leaves; (j)–(l) variations of leaf apex; (m) cells at dorsal lamina; (n), (o) intralaminar limbidial cells at base of vaginant laminae; (p) cross-section of distal part of leaf; (q) cross-section of costa at distal part of leaf; (r), (s) cross-sections of proximal part of leaf; (t) cross-section of costa at proximal part of leaf; (u) cross-section in pluristratose limbidial cells of vaginant laminae; (v) cross-section of stem. 128
- 4.35 *Fissidens* sp1. (a)–(g) Distribution of perichaetia and perigonia on stem (stems were denuded of leaves); (h), (i) close-up view of gametangia, arrow in (h) points to archegonium with inclined tip; (j), (k) caulogenous branch associated with gametangia; (l) close-up view of (k). 129
- 4.36 *Fissidens* sp1. (a) Outer surface of capsule; (b) surface view of exothecial cells; (c), (d) close-up views of stomata; (e) rostellate operculum; (f) peristome teeth; (g) spores; (h) young rhizoidal tuber on axillary rhizoids; (i), (j) close-up views of (h). 130
- 4.37 *Fissidens* sp1. (a)–(d) Plantlets developed on axillary rhizoids, arrows in (c) and (d) point to plantlets developed on subterranean and axillary rhizoids, respectively; (e)–(g) stems were denuded of leaves to show several plantlets on axillary rhizoids, (g) close-up view of (f); (h) arrow points to subterminal branch, (i) two branches developed per leaf axil; (j) arrows point to axillary chloronemata; (k) close-up view of (j); (l)–(n) stems were denuded of leaves to show axillary chloronemata; (o) rhizoidal bud developed on subterranean rhizoids; (p) close-up view of (o). 131
- 4.38 *Fissidens* sp2. (a) Infertile plant unbranched; (b), (c) sympodial branching; (d)–(f) branched infertile plants, arrows point to caulogenous branches; (g) fertile plants with terminal sporophyte, arrows point to seta; (h), (i) fertile plants with lateral sporophyte (only basal parts of sporophyte available); (j) fertile plant, left arrow indicates vaginula in subterminal position, right arrow indicates short caulogenous branch ended with part of seta. All photos were taken of the specimen II-C(16), except (e), (f) of the specimen II-B(16). 137

- 4.39 *Fissidens* sp2. (a) Upper part of stem denuded of leaves, arrows point to caulogenous branches; (b) lateral sporophyte; (c)–(e) stem leaves; (f) cells at leaf apex; (g) cells at dorsal lamina, apex of minor lamina; (h) leaf insertion; (i) cells in the middle of the dorsal lamina; (j) part of stem denuded of leaves, arrow points to a weakly differentiated hyaline nodule; (k) close-up view of (j). 138
- 4.40 *Fissidens* sp2. (a) Cross-section of proximal part of leaf; (b)–(d) close-up views of (a), (b) cross-section of costa (enlarged), showing 5 guide cells; (c), (d) cross-section in pluristratose limbidial cells of vaginant laminae; (e) cross-section of stem. 139
- 4.41 *Fissidens* sp2. (a) Axillary perigonia enclosed with small perigonial leaf, stem leaf was detached; (b) fertile stem showing subterminal perichaetia and naked axillary gametangia composed of one archegonium and one antheridium; (c) part of stem denuded of leaves, showing naked axillary cluster of antheridia; (d) close-up view shows naked axillary cluster of antheridia; (e) part of stem showing terminal and axillary synoious inflorescence (in both clusters antheridia outnumbered archegonia); (f), (g) caulogenous branches associated with naked archegonia; (h), (i) close-up views of (f) and (g), respectively. 140
- 4.42 *Fissidens* sp2. (a) Part of stem showing reduced caulogenous branch with terminal gametangia; (b) close-up view of (a) showing naked gametangia associated with caulogenous branch; (c)–(e) distribution of perichaetia and perigonia along stem. 141
- A.1 Method used in cutting leaf sections. (a) Arranging leaves on 3" × 1" glass slide; arrows indicate proposed line of sectioning. (b) Using a second slide as a guide, half a double-edged razor blade is drawn across the leaves. (c) Side view of sectioning; by altering the angle (Θ) of the blade, the thickness of the sections can be finely controlled. When the blade has passed the vertical ($\Theta = 0^\circ$), the fulcrum moves from P_1 to P_2 , giving even finer control over the last few sections. (d) Photograph of leaf sectioning. Photos by Sean Edwards. 176
- A.2 (a) Vaginant laminae (also called conduplicate part, true, or sheathing laminae), (b) ventral (apical or superior) lamina, (c) dorsal (inferior) lamina, (d) a cross-section of distal half of *Fissidens* leaf, (e) a cross-section of distal half of leaf passed across the apex of minor lamina, (f) a cross-section of proximal half of leaf, (g) photo showing the position of leaf to be sectioned, arrows point to the directions of cuttings. 178