



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ





شبكة المعلومات الجامعية



شبكة المعلومات الجامعية

التوثيق الالكتروني والميكرو فيلم

جامعة عين شمس

التوثيق الالكتروني والميكروفيلم

قسم

نقسم بللّاه العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأفلام قد اعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15 – 20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of
15 – 25c and relative humidity 20-40 %



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



شبكة المعلومات الجامعية



بالرسالة صفحات

لم ترد بالأصل

**CLOMIPHENE CITRATE AND
TAMOXIFEN CITRATE COMBINED THERAPY
FOR TREATMENT OF POLYCYSTIC OVARIAN
DISEASE "APROSPECTIVE CLINICAL STUDY"**

THESIS

Submitted in Partial Fulfillment of
The MASTER DEGREE
In
Obstetrics and Gynaecology

BY

SOZAN MOHAMED SALEH.
(M.B.B.Ch.)

Mansoura University

SUPERVISORS

Dr. AHMED ALAA YASSIN

Professor of
Obstetrics and Gynaecology
Benha Faculty of Medicine

Dr. MOHAMED ABDUL KARIM

Assist. Professor of
Obstetrics and Gynaecology
Benha Faculty of Medicine

Dr. EL-SAYED EL-NAGAR

Assist. Professor of
Obstetrics and Gynaecology
Benha Faculty of Medicine

Benha Faculty of Medicine
Zagazig University

Zagazig University
Benha Faculty of Medicine
Obstetrics and Gynaecology Department

**CLOMIPHENE CITRATE AND
TAMOXIFEN CITRATE COMBINED THERAPY
FOR TREATMENT OF POLYCYSTIC OVARIAN
DISEASE "APROSPECTIVE CLINICAL STUDY"**

THESIS

Submitted in Partial Fulfillment of
The MASTER DEGREE
In
Obstetrics and Gynaecology

BY

**SOZAN MOHAMED SALEH
(M.B.B.Ch.)**

Mansoura University

SUPERVISORS

Dr. AHMED ALAA YASSIN

Professor of
Obstetrics and Gynaecology
Benha Faculty of Medicine

Dr. MOHAMED ABDUL KARIM

Assist. Professor of
Obstetrics and Gynaecology
Benha Faculty of Medicine

Dr. EL-SAYED EL-NAGAR

Assist. Professor of
Obstetrics and Gynaecology
Benha Faculty of Medicine

**Benha Faculty of Medicine
Zagazig University**

1998

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

ACKNOWLEDGMENT

First and Foremost, Thanks to GOD

I would like to express my sincere thanks and deepest gratitude to Prof. Dr. Ahmed Alaa Yassin, Professor of Obstetrics and Gynaecology, Faculty of Medicine, Benha University for giving me the privilege of working under his supervision, for his great help and advice, his continuous encouragement, marvellous support and eminent fatherly guidance have made this work possible.

My sincere thanks and utmost appreciation are presented to Dr. Mohamed Abdul Karim, Assistant Professor of Obstetrics and Gynaecology, Faculty of Medicine, Benha University for his kind enthusiastic supervision, energetic help, moral support, constructive advice, criticism and great aid during this work.

I wish to express my deep thanks to Dr. EL-Sayed EL-Nagar, Associate Professor of Obstetrics and Gynaecology, Faculty of Medicine, Benha University for his valuable assistance in this work.

Lastly, my gratitude goes to all the patients with polycystic ovarian disease who made this study possible.

CONTENTS

	Page
INTRODUCTION	1
AIM OF THE WORK	3
REVIEW OF LITERATURE	4
Epidemiology of polycystic ovarian disease	5
Anatomic pathology of polycystic ovarian disease ..	7
Pathophysiology of polycystic ovarian disease	10
Etiology of polycystic ovarian disease	15
Clinical features of polycystic ovarian disease	25
Diagnosis of polycystic ovarian disease	30
Management of polycystic ovarian disease	39
SUBJECT AND METHODS	65
RESULTS	69
DICUSSION	81
SUMMARY AND CONCLUSION	85
REFERENCES	87
ARABIC SUMMARY	

INTRODUCTION

INTRODUCTION

Anovulation is a very common problem present in a variety of clinical manifestations, including amenorrhea, irregular menses and/or hirsutism. Serious consequences of chronic anovulation are infertility and a greater risk for developing carcinoma of the endometrium and the breast (**Speroff et al., 1994**).

Polycystic ovarian disease (PCOD) represents the common form of normogonadotropic anovulation associated with non cyclic gonadotropins and oestrogens, and in which oestrogen function is intact but cycle failure and androgen excess are evident (**Kase et al., 1990**).

Gross sclerocystic changes in the human ovary were clearly described by **Chereau in 1845**. More interest was aroused in 1935, when this anatomical abnormality was described by **Stein and Leventhal** to be a clinical syndrome consisting of menstrual irregularity featuring amenorrhoea, a history of sterility, masculine type hirsutism and obesity (**Stein and Leventhal, 1935**).

The diagnosis of this disorder relies mostly on the clinical presentation and hormonal profiles of the patient (**Yen, 1980 - Goldzieher, 1981**). Laparoscopy had been used to confirm the diagnosis (**Rojanaskul et al., 1989**).

The development of high resolution real-time ultrasound during the past decade led to the new approach for ovarian visualization (**Huckeloer et al., 1979 - Nitscke-Debelstein, 1980**). In contrast to Laparoscopy where only the external surface of the ovary could be seen, ultrasonic examination allow a direct visualization of the internal structure of the ovaries. It has now replaced Laparoscopy as the method for assessing the ovarian morphology in many clinical conditions (**Rajanasakul et al., 1989**).

Vaginal ultrasounds is now available and allows one to perform a much more precise analysis of internal ovarian morphology. As the probe can be brought closer to the ovaries than by abdominal ultrasound, higher ultrasound frequencies (5 to 7.5 MH) can be used yielding high resolution ovarian picture (**Ardaens et al., 1991**).

AIM OF THE WORK

AIM OF THE WORK

This work was performed to comparison the use of clomiphene citrate and tamoxifen citrate combined therapy followed by HCG for treatment of patients with PCOD and to compare results with the protocol of ovulation induction using clomiphene citrate and HCG only.

Also, comparison of the protocol of ovulation induction; using clomiphene citrate, tamoxifen citrate and HCG after ovarian inhibition with oral contraceptive pills.

The three protocol of ovulation induction were evaluated according to ovulation response and occurrence of pregnancy.