

# **A PHARMACEUTICAL STUDY OF LIPOSOMES AS A VEHICLE FOR DRUG DELIVERY**

**A Thesis**

Presented to the Graduate School  
Faculty of Pharmacy, Alexandria University  
In Partial Fulfillment of the  
Requirements for the Degree

Of

**Master of Pharmaceutical Sciences**

In

**Industrial Pharmacy**

**By**

**Mohammed Maher Mohammed Riyad Mehanna**  
B. Pharm. Sci., University of Alexandria

**June ٢٠٠٨**



# **A PHARMACEUTICAL STUDY OF LIPOSOMES AS A VEHICLE FOR DRUG DELIVERY**

**Presented by**

**Mohammed Maher Mohammed Riyad  
Mehanna**

**B. Pharm. S., University of Alexandria**

**For The Degree of**

**Master of Pharmaceutical Sciences**

**In**

**Industrial Pharmacy**

**Examiner's committee:**

**Approved**

**Prof. Adel M. Motawaa**

-----

**Prof. Magda W. Samaha**

-----

**Prof. Samia A. Nour**

-----

**Dr. Hoda A. Elmaradny**

-----

## **Advisor's Committee**

**Prof. Magda Wadih Samaha**  
Professor of Industrial Pharmacy  
Head of Industrial Pharmacy Department  
Faculty of Pharmacy  
Alexandria University  
Egypt

**Dr. Hoda Ahmed Elmaradny**  
Ass. Professor of Industrial Pharmacy  
Department of Industrial Pharmacy  
Faculty of Pharmacy  
Alexandria University  
Egypt

DEDICATED TO

**PROF. NAWAL KHALAFALLA**  
**MY IDEAL**

**&**

**MRS. ELHAM MEHANNA**  
**MY REAL MOTHER**

## ACKNOWLEDGMENT

First, thanks **Allah** for help and generosity.

I wish to express my deepest thanks, sincere gratitude and everlasting appreciation to **Professor Dr. Magda**

**Wadih Samaha**, Head of Industrial Pharmacy department, Faculty of Pharmacy, University of Alexandria, for giving me the chance to share her creativity and inspiring ideas. On the top of that, her infinite support, patience and continues encouragement throughout the progress of the study.

I also like to extend my great indebtedness and cordial thanks to **Dr. Hoda Ahmed ElMaradny**, Assistant Professor of Industrial Pharmacy, Faculty of Pharmacy, University of Alexandria, for her kind supervision, laborious efforts, wise guidance, valuable supervision and willing assistance during the course of this work and preparation of the thesis.

My best thanks to **Prof. Dr. Adel M. Motawaa**, professor of Industrial Pharmacy, former Dean of Faculty of Pharmacy, University of Alexandria and Beirut University, for his encouragement and support.

Appreciation is expressed for the help received from the Departments of Pharmaceutics and Microbiology, staff and

*members; especially to my dear colleague A. Allam, Ass. Lecturer, Department of Pharmaceutics, for the endless discussions and helpful tips and O. El-halfawy, Demonstrator, Department of Microbiology for sincere help during microbiological studies.*

*Finally, my indebtedness is due to my parents and my sisters for their interest in addition, sincere help that were of great value in accomplishing this thesis.*

*Thank you  
Mohammed M. Mehanna*

# CONTENTS

LIST OF TABLES	i
LIST OF FIGURES	iii
LIST OF ABBREVIATIONS	vi
GENERAL INTRODUCTION	v
OBJECTIVES OF THE THESIS	۳۷
<b><u>Part I</u></b>	
<b>UTILIZATION OF FACTORIAL DESIGN TO STUDY THE PREPARATION OF CIPROFLOXACIN HCL OCULAR LIPOSOMES AND THEIR OPTIMIZATION</b>	
۱. Introduction	۴۰
Aim of Part I	۴۳
۲. Experimental	
۲.۱. Materials	۴۴
۲.۲. Equipment	۴۵
۲.۳. Methodology	
۲.۳.۱. U.V. spectrophotometry of ciprofloxacin HCl in isotonic phosphate buffer pH ۴.۵ and ۷.۴	۴۶
۲.۳.۲. Construction of calibration curve of ciprofloxacin HCl in isotonic phosphate buffer pH ۴.۵ and ۷.۴	۴۶
۲.۳.۳. Preparation of ciprofloxacin HCl-loaded liposomes	۴۶
۲.۳.۴. Separation of free untrapped ciprofloxacin HCl from liposomes	۴۷
۲.۳.۵. Experimental design	۴۷
۲.۳.۶. Characterization of ciprofloxacin HCl-loaded liposomes	۵۰
۲.۳.۷. Differential scanning calorimetry (DSC) analysis	۵۱
۲.۳.۸. Data analysis of factorial design	۵۱
۲.۳.۹. Check point analysis (CPA)	۵۲
۳. Results and discussion	
۳.۱. Spectrophotometerical analysis of ciprofloxacin HCl	۵۳
۳.۲. Selection of ciprofloxacin HCl liposomes preparation method	۵۳
۳.۳. Factorial analysis	۵۷
۳.۳.۱. Entrapment efficiency (EE%)	۵۷
۳.۳.۲. Particle size	۶۳
۳.۳.۳. Burst release from liposomes	۷۶
۳.۳.۴. Percentage of drug released after ۱۰ hr	۸۳
۳.۴. Check point analysis (CPA)	۸۴
۳.۵. Kinetic analysis of in vitro release	۹۴
۳.۶. Ciprofloxacin HCl-membrane interaction	۹۵
۴. Conclusion	۱۰۳

## **Part II**

### **FORMULATION OF PROLONGED RELEASE OCULAR LIQUID DOSAGE FORMS BASED ON CIPROFLOXACIN HCl LIPOSOMES**

1. Introduction	105
Aim of Part II	107
2. Experimental	
2.1. Materials	108
2.2. Equipment	109
2.3. Methodology	
2.3.1. Preparation of chitosan coated liposomes (Chitosomes)	110
2.3.1.1. Determination of entrapment efficiency percent	110
2.3.1.2. Morphological evaluation and particle size analysis of chitosomes	110
2.3.1.3. Determination of sedimentation volume	111
2.3.1.4. In vitro release of ciprofloxacin HCl from chitosomes	111
2.3.1.5. Kinetic analysis of the release data	111
2.3.1.6. Rheological evaluation of chitosomes	111
2.3.1.7. Ex-in vivo permeation studies	111
2.3.2. Preparation of poloxamer gel and dispersion of liposomes	112
2.3.2.1. In vitro release of ciprofloxacin HCl from Pluronic F-127 gel	112
2.3.2.2. Drug release kinetics	112
2.3.2.3. Rheological evaluation of the PF-127 gels	113
3. Results and discussion	
3.1. Chitosan – coated liposomes (chitosomes)	114
3.1.1. Morphological evaluation and particle size analysis	114
3.1.2. Entrapment Efficiency	117
3.1.3. Sedimentation volume	118
3.1.4. In vitro release studies	122
3.1.5. Kinetic models of in vitro drug release	124
3.1.6. Rheological studies	124
3.1.7. Ex-in vivo permeation studies	126
3.2. Liposomes dispersed in Pluronic F-127 gel	127
3.2.1. Ciprofloxacin HCl release kinetics from PF-127 gel	127
3.2.2. Rheological behaviour of Pluronic F-127 gel	130
4. Conclusion	134

## **Part III**

### **MICROBIOLOGICAL AND IN VIVO EVALUATION OF CIPROFLOXACIN HCl PROLONGED RELEASE OCULAR LIQUID DOSAGE FORMS**

1. Introduction	136
Aim of Part III	138
2. Experimental	
2.1. Materials	139
2.2. Equipment	140
2.3. Methodology	
2.3.1. Microbiological Evaluation	141

۲.۳.۱.۱. Preparation of the culture media	۱۴۱
۲.۳.۱.۲. Isolation of microorganisms and maintenance of the isolates	۱۴۱
۲.۳.۱.۳. Determination of the minimum inhibitory concentration (MIC) of liposomes and chitosomes against the tested microorganisms using agar dilution technique	۱۴۱
۲.۳.۱.۴. Determination of the antimicrobial efficacy of Pluronic F-۱۲۷ gel containing liposomes against the tested microorganisms using cup-plate technique	۱۴۲
۲.۳.۲. In vivo Evaluation	۱۴۳
۲.۳.۲.۱. Rabbit eye study	۱۴۳
۲.۳.۲.۲. Animal studies: pharmacodynamic properties	۱۴۳
۲.۳.۲.۳. Statistical analysis	۱۴۳
۳. Results and discussion	
۳.۱. Microbiological studies	
۳.۱.۱. In vitro antimicrobial activity of liposomes containing ciprofloxacin HCl	۱۴۴
۳.۱.۲. In vitro antimicrobial activity of chitosomes containing ciprofloxacin HCl	۱۴۴
۳.۱.۳. In vitro antimicrobial activity of Pluronic F-۱۲۷ in situ gel containing liposomal ciprofloxacin HCl	۱۴۷
۳.۲. In vivo Evaluation	
۳.۲.۱. Single dose study	۱۴۸
۳.۲.۲. Evaluation of different formulations using in vivo model (rabbit eye conjunctivitis)	۱۴۹
۴. Conclusion	۱۵۲
SUMMARY	۱۵۳
REFERENCES	۱۵۸
ARABIC SUMMARY	

## LIST OF TABLES

Table	Title	Page
1	Some head group alcohols of the phosphoglycerides	7
2	Liposomes classification based on their method of preparation	10
3	Liposomes classification based on composition and mode of drug delivery	12
4	Liposomes classification by size and number of lamellae	13
5	Applications of liposomes in sciences	22
6	Variables and responses in 2 <sup>o</sup> full factorial design	48
7	Composition of different liposomal dispersions prepared by reverse phase evaporation method according to 2 <sup>o</sup> full factorial design	49
8	Formulation variables in check point analysis	52
9	Entrapment efficiency, mean particle diameter and Span Index of liposomes produced by various methods	56
10	Entrapment efficiency response of ciprofloxacin HCl-loaded liposomes prepared by reverse phase evaporation method according to 2 <sup>o</sup> factorial design	58
11	Analysis of variance table for the entrapment efficiency response	60
12	The experimental (actual) and predicted values of entrapment efficiency (%) response and percent bias	60
13	Mean particle size diameter response of ciprofloxacin HCl-loaded liposomes prepared by reverse phase evaporation method according to 2 <sup>o</sup> factorial design along with its Span Index	66
14	Analysis of variance table for mean particle size diameter response	73
15	Percentage of drug released after 1 hr (burst release) and 10 hr responses of ciprofloxacin HCl-loaded liposomes prepared by reverse phase evaporation method according to 2 <sup>o</sup> factorial design	77
16	Analysis of variance table for percentage drug released at 1 hr (burst effect) response	80
17	The experimental (actual) and predicted values of burst release response and percent bias	80
18	Analysis of variance table for % drug released after 10 hr response	90
19	The experimental and predicted values of % drug released after 10 hr response and percent bias	93
20	Actual and predicted responses of check point	94
21	Drug release kinetic parameters of reverse phase evaporation liposomes	97
22	Diffusion and erosion coefficient values of reverse phase evaporation liposomes calculated using Kopcha equation	98
23	Thermotropic parameters of interaction between ciprofloxacin HCl and different types of liposomes with regard to endothermic peak of phase transition	102
24	Composition of different chitosomal formulations	113
25	Effect of coating with different concentrations and molecular weights of chitosan on particle size of chitosomes	116
26	Entrapment efficiency of different chitosomal dispersions	119
27	The influence of chitosan concentration and liposomes surface charge on sedimentation volume of chitosomes	121
28	Release kinetic model by fitting in vitro release studies	124

२९	Viscosity of different chitosomal dispersions	१२०
३०	Corneal permeability and apparent permeability coefficient of selected chitosomal preparations	१२६
३१	kinetic analysis of ciprofloxacin HCl release from PF-१२१ gel	१२८
३२	Drug release mechanism in relation to diffusion exponent value	१२८
३३	Kinetic parameters of ciprofloxacin HCl release obtained by fitting the experimental results to equations (१६) and (११)	१२८
३४	Rheological behaviour of different gel formulations	१३२
३०	In vitro activity of different liposomes formulations containing ciprofloxacin HCl against reference strains and clinical isolates in nutrient agar	१४०
३६	In vitro activity of chitosomes formulation containing ciprofloxacin HCl against reference strains and clinical isolates in nutrient agar	१४६
३१	Antimicrobial efficacy of the prepared in situ PF-१२१ gel containing ciprofloxacin HCl-loaded liposomes by agar diffusion technique	१४१
३८	In vivo single dose study of the prepared formulations	१४८
३९	Assessment scale of experimentally induced bacterial conjunctivitis in rabbits' eye	१४९
४०	Mean grades for severity of redness, lacrimal secretion, mucoidal discharge, response to ocular stimulus and swelling of eyelid in bacterial conjunctivitis rabbit model	१०१

## LIST OF FIGURES

Figure	Title	Page
1	Schematic representation of liposome structure	5
2	The general chemical structure of phospholipid has a glycerol backbone	6
3	Chemical structure of cholesterol	7
4	Effect of molecular geometry on phase properties of lipids	9
5	Predominant mechanisms of intracellular drug delivery by liposomes	11
6	Mechanism of remote loading of doxorubicin (DOX) into the intraliposome aqueous phase by ammonium sulfate gradient	18
7	Anatomical features of the eye in cross-section, illustrating major routes of administration	28
8	Diagrammatic representation of different factors attributing to poor ocular bioavailability of an ophthalmic formulation	29
9	Chemical structure of ciprofloxacin	40
10	Absorption curve of ciprofloxacin HCl in isotonic phosphate buffer pH 7.4 (a) and phosphate buffer pH 5.0 (b)	54
11	Calibration curve of ciprofloxacin HCl in isotonic phosphate buffer pH 7.4 at $\lambda_{max}$ 272 nm (a) and phosphate buffer pH 5.0 at $\lambda_{max}$ 277 nm (b)	55
12	Optical photographs (I) and transmission electron microscopical photomicrographs (II) of ciprofloxacin HCl-loaded liposomes prepared by different methods	56
13	Normal probability plot of residuals for the entrapment efficiency response	59
14	Plot of outlier T (externally studentized residuals) versus run number for the entrapment efficiency response	59
15	Plot of residuals versus predicted values of entrapment efficiency response	59
16	Normal probability plot of the effects before (a) and after (b) selection of the main effects for entrapment efficiency response	61
17	One factor plot of main effect of cholesterol molar concentration on the entrapment efficiency	62
18	Two (a) and three-dimensional (b) plots for the multifactor effect of dicetylphosphate concentration and pH of aqueous phase on the entrapment efficiency response	64
19	The cumulative size distribution curve of some prepared liposomal dispersions	67
20	Photographs of some prepared liposomal dispersions	68
21	Normal probability plot of residuals for the particle size response	70
22	Plot of externally studentized residuals versus run number for the particle size response	70
23	Plot of residuals versus predicted values for the particle size response	70
24	Box-Cox plot of particle size response before (a) and after (b) power transformation	71
25	Normal probability plot of residuals for the particle size response after transformation	72
26	Plot of externally studentized residuals versus run number for the particle size response after transformation	72
27	Half normal probability plot of the effects before (a) and after (b) selection of the main effects for particle size response	74
28	Contour plot (a) and interaction curve (b) for the multifactor effect of cholesterol and dicetylphosphate concentration on the particle size response	75
29	Normal probability plot of residuals for the % drug released at 1 hr response	78

३०	Plot of studentized residuals versus predicted values for the % drug released at १ hr response	१४
३१	Plot of studentized residuals versus run number for the % drug released at १ hr response	१४
३२	Plot of outlier T (externally studentized residuals) versus run number for the response % drug released at १ hr response	१५
३३	Box-Cox plot of the % drug released at १ hr response	१५
३४	Normal probability plot of the effects before (a) and after (b) selection of the main effects for % drug released at १ hr response	१६
३०	One factor plot of main effect of dicetyl phosphate molar concentration on the % drug released at १ hr response	१७
३६	Interaction plot between dicetyl phosphate molar concentration and pH of the aqueous phase on the % drug released at १ hr response	१७
३१	In vitro release profiles of ciprofloxacin-loaded liposomes prepared using reverse phase evaporation method according to १ factorial design (F <sup>१</sup> , F <sup>०</sup> , F <sup>११</sup> , F <sup>११</sup> )	१७
३४	In vitro release profiles of ciprofloxacin-loaded liposomes prepared using reverse phase evaporation method according to १ factorial design (F <sup>१</sup> , F <sup>१</sup> , F <sup>११</sup> , F <sup>११</sup> )	१७
३५	In vitro release profiles of ciprofloxacin-loaded liposomes prepared using reverse phase evaporation method according to १ factorial design (F <sup>१</sup> , F <sup>१</sup> , F <sup>११</sup> , F <sup>११</sup> )	१७
४०	In vitro release profiles of ciprofloxacin-loaded liposomes prepared using reverse phase evaporation method according to १ factorial design (F <sup>१</sup> , F <sup>१</sup> , F <sup>११</sup> , F <sup>११</sup> )	१७
४१	In vitro release profiles of ciprofloxacin-loaded liposomes prepared using reverse phase evaporation method according to १ factorial design (F <sup>१</sup> , F <sup>११</sup> , F <sup>११</sup> , F <sup>११</sup> )	१८
४२	In vitro release profiles of ciprofloxacin-loaded liposomes prepared using reverse phase evaporation method according to १ factorial design (F <sup>११</sup> , F <sup>११</sup> , F <sup>११</sup> , F <sup>११</sup> )	१८
४३	In vitro release profiles of ciprofloxacin-loaded liposomes prepared using reverse phase evaporation method according to १ factorial design (F <sup>११</sup> , F <sup>११</sup> , F <sup>११</sup> , F <sup>११</sup> )	१९
४४	In vitro release profiles of ciprofloxacin-loaded liposomes prepared using reverse phase evaporation method according to १ factorial design (F <sup>११</sup> , F <sup>११</sup> , F <sup>११</sup> , F <sup>११</sup> )	१९
४०	Normal probability plot of the effects before (a) and after (b) selection of the main effects for % drug released at १.० hr response	१९
४१	One factor plot of main effect of cholesterol molar concentration on the % drug released at १.० hr response	१९
४१	One factor plot of main effect of pH of the aqueous phase on the % drug released at १.० hr response	१९
४४	DSC thermogram of ciprofloxacin HCl	१९
४५	DSC thermogram of cholesterol	१९
००	DSC thermogram of plain liposomes made up with cholesterol	१९
०१	DSC thermogram of plain neutral liposomes	१००
०२	DSC thermogram of neutral liposomes-loaded ciprofloxacin HCl at pH ४.०	१००
०३	DSC thermogram of neutral liposomes-loaded ciprofloxacin HCl at pH १.४	१००
०४	DSC thermogram of plain negative liposomes	१०१
००	DSC thermogram of negative liposomes-loaded ciprofloxacin HCl at pH ४.०	१०१
०१	DSC thermogram of negative liposomes-loaded ciprofloxacin HCl at pH १.४	१०१

57	Representative images of three formulations of chitosomes. Images were obtained by optical (I) and transmission electron microscopes (II)	110
58	Effect of coating with different concentrations, molecular weights of chitosan and type of liposomes surface charge on particle size of chitosomes	116
59	Effect of chitosan concentration on entrapment efficiency (%) of different chitosomal dispersions	120
60	Effect of chitosan molecular weight on entrapment efficiency (%) of different chitosomal dispersions	120
61	Effect of charge type of liposomes on entrapment efficiency (%) of different chitosomal dispersions	120
62	The influence of different variables on sedimentation volume of chitosomes	121
63	In vitro release profiles of different chitosomal formulations (C <sup>1</sup> , C <sup>2</sup> , C <sup>3</sup> , C <sup>4</sup> ) into isotonic phosphate buffer pH 7.4	123
64	In vitro release profile of a chitosomal formulation (C <sup>1</sup> ) into isotonic phosphate buffer pH 7.4 in comparison to drug solution and liposomal dispersion (F <sup>1</sup> )	123
65	Effect of coating with different chitosan concentrations and molecular weights on the rheological properties of chitosomes at 30°C	125
66	Cumulative amount of ciprofloxacin HCl released from Pluronic F-127 (2.0% w/v) control gel and liposomal gel containing phosphatidylcholine/cholesterol/dicetylphosphate (1:1:1 mol/mol) liposomes	129
67	Higuchi release plot of ciprofloxacin HCl from PF-127 gel	129
68	Higuchi release plot of liposomal ciprofloxacin HCl from PF-127 gel	129
69	Rheogram of a 2.0% aqueous of Pluronic F-127 at 4°C	131
70	Rheogram of a 2.0% aqueous of Pluronic F-127 at 34°C	131
71	Rheological profile of temperature- triggered in situ gelling system (PF-127) in presence of NaCl salt	132
72	Rheological profile of temperature- triggered in situ gelling system (PF-127) of different formulations	133
73	The growth inhibition zone with liposomal gel (1), control gel (2) and drug solution (3) using <i>Pseudomonas aeruginosa</i>	147
74	Normal rabbit eye showing no redness	150
75	Bacterial conjunctivitis in rabbit eye showing redness	150
76	Normal rabbit eye showing normal eyelid	150
77	Bacterial conjunctivitis in rabbit eye showing eyelid edema	150
78	Normal rabbit eye showing no mucoidal exudates	150
79	Bacterial conjunctivitis in rabbit eye showing mucoidal exudates	150
80	Normal rabbit eye with normal tearing	150
81	Bacterial conjunctivitis in rabbit eye showing excessive tearing	150