ELECTROCHEMICAL PROPERTIES OF SOME SOILS OF EGYPT IN RELATION TO MINERALOGICAL COMPOSITION

BY

MOHAMED SALAH KOTB

(B. Sc. Agric. Soil Science)

Cairo University, 2000

THESIS

Submitted to the Graduate Division in Partial Fulfilment of the

Requirements for the Degree of Master Science

In

Agricultural Sciences (Soils Science)

Faculty of Agriculture Cairo University Egypt

2006

Advisory Committee

Senior Supervisor Prof. Dr. R.R.Shahin Professor of Soil Chemistry, Soil Science Department, Faculty of Agricultural, Cairo University.

Assistant Supervisors

Dr. Y.A.Nasr Associate Professor of Soil Chemistry, Soil Science Department, Faculty of Agricultural, Cairo University. Dr. M.M.Abdel-Tawab Associate Professor of Soil Chemistry, Soil Science Department, Faculty of Agricultural, Cairo University.

APPROVAL SHEET

Name: Mohamed Salah Kotb Ramadan

Title: ELECTROCHEMICAL PROPERTIES OF SOME SOILS OF EGYPT IN RELATION TO MINERALOGICAL COMPOSITION

THESIS Submitted for the Degree of Master

In

Agricultural Sciences (Soils Science)

Thesis has been approved by:

Prof. Dr.

Prof. Dr.

Prof. Dr.

(committee in charge)

Date: / / 2006

ACKNOWLEDGEMENT

The author wishes to express his gratitude and sincere appreciation to Prof. Dr. Reda.R.Shahin, Soils Department, Faculty of Agriculture, Cairo University, for his kind supervision, advice, helpful criticism throughout the study and valuable guidance during the preparation of the manuscript.

Thanks are extended to Dr. Y.A.Nasr and M.M.Abdel-Tawab, Assistant Professors, Soils Department, Faculty of Agriculture, Cairo University for their supervision and kindly assistance.

CONTENTS

	Page
LIST OF FIGURES	
LIST OF TABLES	
1- INTRODUCTION	1
2- REVIEW OF LITERATURE	2
2.1. Charge development in soil	2
2.1.1. Permanent charge	4
2.1.2. pH-Dependent	8
2.2. Charge balance of mineral surfaces :	14
2.3. Point of zero charge:	16
2.3.1. Definitions of points of zero charge:	16
2.3.2. Determination methods of points of zero charge:	18
	19
2.3.2.1. Potentiometric titration curves:	
2.3.2.2. Ion adsorption method:	20
2.3.2.3. Electrophoretic mobility	21
2.4 Factors affecting charge characteristic parameters in soils	21
2.4.1. Ionic strength and temperature	21
2.4.2. Mineralogical composition	23
2.4.3. Iron oxides	25
2.4.4. Organic matter content	26
2.5. P-oxyanion adsorption impact on charge characteristics:	27
3. MATERIALS AND METHODS	30
3.1. Materials	30
3.1.1. Soil sampling	30
3.1.2. Standard clay materials	30
3.1.3. Separation of soil clay fractions	30
3.2. Methods of analysis	32
3.2.1. Physical determinations	32

3.2.2. Chemical determinations	32
3.2.3. Mineralogical analysis	33
1- X-ray analysis	33
2- Determination of amorphous materials	33
3.2.4. Potentiometric titration curves	34
3.2.5. Ion – adsorption experiment	36
3.2.6. P-oxyanion adsorption isotherms	36
3.2.7. Cation exchange isotherms :	37
4- RESSULTS AND DISCUSSION	38
4.1. Distribution of amorphous materials in clay fractions:	38
4.2. Specific surface area of clay fractions:	38
4.3. Mineralogical composition of soil clay fractions:	40
4.4. Charge characteristics of soil clay separates :	50
4.4.1. Potentiometric titration curves:	50
(a) Reference clay material	52
(b) Soil clay fractions:	55
4.4.2. Ion adsorption curves:	65
4.5. Cation exchange isotherms of clay separates:	86
4.6. Phosphate adsorption isotherms of clay fractions:	105
4.7. Charge characteristics of P-saturated clay separates :	125
4.7.1. Potentiometric titrations of P-saturated clays:	125
4.7.2. The effect of P-oxyanion adsorption on potntiometric	
titration curves and their charge parameters:	135
4.7.3. Ion adsorption curves of P-saturated clays	145
5. SUMMARY	173
6. REFERENCES	179
7. APPENDIX	193
8. ARABIC SUMMARY	232

LIST OF FIGURES	
Fig. (1) Distribution of the total amorphous materials as	39
percentage of the clay fraction (%).	
Fig. (2) X-ray diffractograms of Aswan fine and coarse clay	43
separates .	
Fig. (3) X-ray diffractograms of El-menia fine and coarse clay	44
separates .	
Fig. (4) X-ray diffractograms of Fayium fine and coarse clay	45
separates .	
Fig. (5) X-ray diffractograms of Tanta fine and coarse clay	46
separates .	
Fig. (6) X-ray diffractograms of Kafr El-Sheikh fine and coarse	47
clay separates .	
Fig. (7) X-ray diffractograms of Belbeis fine and coarse clay	48
separates .	
Fig. (8) X-ray diffractograms of Nubaryia fine and coarse clay	49
separates .	
Fig. (9) Distribution of expanding and non-expanding minerals in	51
fine and coarse clay fractions of the investigated soils.	
Fig. (10) Potentiometric titration curves for Kaoline and Bentonite	53
with and without amorphous material.	
Fig.(11) Charge parameters as calculated from potentiometric	54
titration curves.	
Fig. (12) Potentiometric titration curves for Aswan fine and	57
coarse clay with and without amorphous material.	
Fig. (13) Potentiometric titration curves for El-menia fine and	58
coarse clay with and without amorphous material.	

Fig. (14) Potentiometric titration curves for Fayium fine and	59
coarse clay with and without amorphous material.	
Fig. (15) Potentiometric titration curves for Tanta fine and coarse	60
clay with and without amorphous material.	
Fig. (16) Potentiometric titration curves for Kafr El-Sheikh fine	61
and coarse clay with and without amorphous material.	
Fig. (17) Potentiometric titration curves for Belbeis fine and	62
coarse clay with and without amorphous material.	
Fig. (18) Potentiometric titration curves for Nubaryia fine and	63
coarse clay with and without amorphous material.	
Fig. (19) Ion adsorption curves for Kaoline and Bentonite with	66
and witout amorphous material.	
Fig. (20) Ion adsorption curves for Aswan fine and coarse clay	67
with and without amorphous material.	
Fig. (21) Ion adsorption curves for El-menia fine and coarse clay	68
with and without amorphous material.	
Fig. (22) Ion adsorption curves for Fayium fine and coarse clay	69
with and without amorphous material.	
Fig. (23) Ion adsorption curves for Tanta fine and coarse clay with	70
and without amorphous material.	
Fig. (24) Ion adsorption curves for Kafr El-Sheikh fine and coarse	71
clay with and without amorphous material.	
Fig. (25) Ion adsorption curves for Belbeis fine and coarse clay	72
with and without amorphous material.	
Fig. (26) Ion adsorption curves for Nubaryia fine and coarse clay	73
with and without amorphous material.	
Fig.(27) Charge parameters as calculated from ion adsorption	74
curves.	
L	1

Fig. (28) Na-Cl adsorption curves and net charge for Kaoline and	76
Bentonite with and witout amorphous material.	
Fig. (29) Na-Cl adsorption curves and net charge for Aswan fine	77
and coarse clay with and without amorphous material.	
Fig. (30) Na-Cl adsorption curves and net charge for El-menia	787
fine and coarse clay with and without amorphous material.	
Fig. (31) Na-Cl adsorption curves and net charge for Fayium fine	79
and coarse clay with and without amorphous material.	
Fig. (32) Na-Cl adsorption curves and net charge for Tanta fine	80
and coarse clay with and without amorphous material.	
Fig. (33) Na-Cl adsorption curves and net charge for Kafr El-	81
Sheikh fine and coarse clay with and without	
amorphous material.	
Fig. (34) Na-Cl adsorption curves and net charge for Belbeis fine	82
and coarse clay with and without amorphous material.	
Fig. (35) Na-Cl adsorption curves and net charge for Nubaryia	83
fine and coarse clay with and without amorphous	
material .	
Fig.(36) Charge parameters as calculated from Na-Cl adsorption	85
curves.	
Fig. (37) Sodium - calcium exchange isotherms for Kaoline and	87
Bentonite before and after the removal of amorphous	
material.	
Fig. (38) Sodium - calcium exchange isotherms for clay separates	88
of Aswan before and after the removal of amorphous	
material.	
Fig. (39) Sodium - calcium exchange isotherms for clay separates	89
of El-menia before and after the removal of amorphous material.	
	1

Fig. (40) Sodium - calcium exchange isotherms for clay separates	90
of Fayium before and after the removal of amorphous	<i>y</i> v
material.	
	0.1
Fig. (41) Sodium - calcium exchange isotherms for clay separates	91
of Tanta before and after the removal of amorphous	
material.	
Fig. (42) Sodium - calcium exchange isotherms for clay separates	92
of Kafr El-Sheikh before and after the removal of	
amorphous material.	
Fig. (43) Sodium - calcium exchange isotherms for clay separates	93
of Belbeis before and after the removal of amorphous	
material.	
Fig. (44) Sodium - calcium exchange isotherms for clay separates	94
of Nubaryia before and after the removal of amorphous	
material.	
Fig. (45) Gapon relationship of Ca-Na exchange of the reference	97
Kaoline and Bentonite clays with and without	
amorphous material.	
Fig. (46) Gapon relationship of Ca-Na exchange of fine and	98
coarse clays of Aswan with and without amorphous material.	
Fig. (47) Gapon relationship of Ca-Na exchange of fine and	99
coarse clays of El-menia with and without amorphous material.	
Fig. (48) Gapon relationship of Ca-Na exchange of fine and	100
coarse clays of Fayium with and without amorphous material.	
Fig. (49) Gapon relationship of Ca-Na exchange of fine and	101
coarse clays of Tanta with and without amorphous material.	
Fig. (50) Gapon relationship of Ca-Na exchange of fine and	102
coarse clays of Kafr El-Sheikh with and without a.m.	

Fig. (51) Gapon relationship of Ca-Na exchange of fine and	103
coarse clays of Belbeis with and without amorphous material.	
Fig. (52) Gapon relationship of Ca-Na exchange of fine and	104
coarse clays of Nubaryia with and without amorphous material.	
Fig. (53) Adsorped - P vs. final soluble - P at pH = 3 (\blacktriangle) and 9	106
(•) For Kaoline and Bentonite .	
Fig. (54) Adsorped - P vs. final soluble - P at pH = 3 (\blacktriangle) and	107
(•) For fine clay separates of the studied soils.	
Fig. (55) Adsorped - P vs. final soluble - P at pH = 3 (\blacktriangle) and 9 (108
•) For fine clay separates of the studied soils.	
Fig. (56) Adsorped - P vs. final soluble - P at pH = 3 (\blacktriangle) and 9 (109
•) For coarse clay separates of the studied soils.	
Fig. (57) Adsorped - P vs. final soluble - P at pH = 3 (\blacktriangle) and 9 (110
•) For coarse clay separates of the studied soils.	
Fig. (58) P - adsorption isotherms at pH = 3 (\blacktriangle) and 9 (\bullet) of	116
Kaoline and Bentonite (with / without amorphous	
material) according to linearlized - langmuir model.	
Fig. (59) P - adsorption isotherms at pH = 3 (\blacktriangle) and 9 (\bullet) of	117
fine clay fractions (with / without amorphous material)	
according to linearlized - langmuir model.	
Fig. (60) P - adsorption isotherms at pH = 3 (\blacktriangle) and 9 (\bullet) of	118
fine clay fractions (with / without amorphous material)	
according to linearlized - langmuir model.	
Fig. (61) P - adsorption isotherms at pH = 3 (\blacktriangle) and 9 (\bullet) of	119
coarse clay fractions (with / without amorphous	
material) according to linearlized - langmuir model.	
Fig. (62) P - adsorption isotherms at pH = 3 (\blacktriangle) and 9 (\bullet) of	120
coarse clay fractions (with / without amorphous material)	

according to linearlized - langmuir model.	
Fig. (63) Potentiometric titration curves for P-saturated Kaoline	126
and Bentonite with and without amorphous material.	
Fig. (64) Potentiometric titration curves for P-saturated Aswan	127
fine and coarse clay with and without amorphous material.	
Fig. (65) Potentiometric titration curves for P-saturated El-menia	128
fine and coarse clay with and without amorphous material.	
Fig. (66) Potentiometric titration curves for P-saturated Fayium	129
fine and coarse clay with and without amorphous material.	
Fig. (67) Potentiometric titration curves for P-saturated Tanta fine	130
and coarse clay with and without amorphous material.	
Fig. (68) Potentiometric titration curves for P-saturated Kafr El-	131
Sheikh fine and coarse clay with and without	
amorphous material.	
Fig. (69) Potentiometric titration curves for P-saturated Belbeis	132
fine and coarse clay with and without amorphous material.	
Fig. (70) Potentiometric titration curves for P-saturated Nubaryia	133
fine and coarse clay with and without amorphous material.	
Fig. (71) Possible configuration of the P-surface complexation	136
(Ryden et al., 1977).	
Fig.(72) Potentiometric titration curves at ionic strength (I = 0.01	137
M) for Kaoline and Bentonite with and without A.M. [
• (-P) ' • (+P)]	
Fig.(73) Potentiometric titration curves at ionic strength (I = 0.01	138
M) for Aswan with and without A.M. [\bullet (-P) ' \bullet (+P)]	
Fig.(74) Potentiometric titration curves at ionic strength (I = 0.01	139
M) for El-menia with and without A.M. [• (-P) ' • (+P)]	

 Fig.(75) Potentiometric titration curves at ionic strength (I = 0.01 M) for Fayium with and without A.M. [• (-P) ' • (+P)] Fig.(76) Potentiometric titration curves at ionic strength (I = 0.01 M) for Tanta with and without A.M. [• (-P) ' • (+P)] Fig.(77) Potentiometric titration curves at ionic strength (I = 0.01 M) for Kafr El-Sheikh with and without A.M. [• (-P) ' • (+P)] Fig.(78) Potentiometric titration curves at ionic strength (I = 0.01 M) for Belbeis with and without A.M. [• (-P) ' • (+P)] Fig.(79) Potentiometric titration curves at ionic strength (I = 0.01 M) for Nubaryia with and without A.M. [• (-P) ' • (+P)] Fig.(80a,b,c) Effect P-saturation on total charge (σ_T) of the studied clay separates. Fig. (81) Ion adsorption curves for P-saturated Kaoline and Bentonite with and witout amorphous material .
M) for Tanta with and without A.M. [• (-P) ' • (+P)] Fig.(77) Potentiometric titration curves at ionic strength (I = 0.01 142 M) for Kafr El-Sheikh with and without A.M. [• (-P) ' • (+P)] Fig.(78) Potentiometric titration curves at ionic strength (I = 0.01 143 M) for Belbeis with and without A.M. [• (-P) ' • (+P)] Fig.(79) Potentiometric titration curves at ionic strength (I = 0.01 144 M) for Nubaryia with and without A.M. [• (-P) ' • (+P)] Fig.(80a,b,c) Effect P-saturation on total charge (σ _T) of the studied clay separates. Fig. (81) Ion adsorption curves for P-saturated Kaoline and Bentonite with and witout amorphous material .
 Fig.(77) Potentiometric titration curves at ionic strength (I = 0.01 M) for Kafr El-Sheikh with and without A.M. [• (-P) ' • (+P)] Fig.(78) Potentiometric titration curves at ionic strength (I = 0.01 M) for Belbeis with and without A.M. [• (-P) ' • (+P)] Fig.(79) Potentiometric titration curves at ionic strength (I = 0.01 M) for Nubaryia with and without A.M. [• (-P) ' • (+P)] Fig.(80a,b,c) Effect P-saturation on total charge (σ_T) of the studied clay separates. Fig. (81) Ion adsorption curves for P-saturated Kaoline and Bentonite with and witout amorphous material .
 M) for Kafr El-Sheikh with and without A.M. [• (-P) ' • (+P)] Fig.(78) Potentiometric titration curves at ionic strength (I = 0.01 M) for Belbeis with and without A.M. [• (-P) ' • (+P)] Fig.(79) Potentiometric titration curves at ionic strength (I = 0.01 M) for Nubaryia with and without A.M. [• (-P) ' • (+P)] Fig.(80a,b,c) Effect P-saturation on total charge (σ_T) of the studied clay separates. Fig. (81) Ion adsorption curves for P-saturated Kaoline and Bentonite with and witout amorphous material .
 Fig.(78) Potentiometric titration curves at ionic strength (I = 0.01 M) for Belbeis with and without A.M. [• (-P) ' • (+P)] Fig.(79) Potentiometric titration curves at ionic strength (I = 0.01 M) for Nubaryia with and without A.M. [• (-P) ' • (+P)] Fig.(80a,b,c) Effect P-saturation on total charge (σ_T) of the studied clay separates. Fig. (81) Ion adsorption curves for P-saturated Kaoline and Bentonite with and witout amorphous material .
M) for Belbeis with and without A.M. [• (-P) ' • (+P)] Fig.(79) Potentiometric titration curves at ionic strength (I = 0.01 M) for Nubaryia with and without A.M. [• (-P) ' • (+P)] Fig.(80a,b,c) Effect P-saturation on total charge (σ _T) of the studied clay separates. Fig. (81) Ion adsorption curves for P-saturated Kaoline and Bentonite with and witout amorphous material.
 Fig.(79) Potentiometric titration curves at ionic strength (I = 0.01 M) for Nubaryia with and without A.M. [• (-P) ' • (+P)] Fig.(80a,b,c) Effect P-saturation on total charge (σ_T) of the studied clay separates. Fig. (81) Ion adsorption curves for P-saturated Kaoline and Bentonite with and witout amorphous material .
M) for Nubaryia with and without A.M. [• (-P) ' • (+P)] Fig.(80a,b,c) Effect P-saturation on total charge (σ _T) of the studied clay separates. Fig. (81) Ion adsorption curves for P-saturated Kaoline and Bentonite with and witout amorphous material.
 Fig.(80a,b,c) Effect P-saturation on total charge (σ_T) of the studied clay separates. Fig. (81) Ion adsorption curves for P-saturated Kaoline and Bentonite with and witout amorphous material.
studied clay separates. Fig. (81) Ion adsorption curves for P-saturated Kaoline and Bentonite with and witout amorphous material.
Fig. (81) Ion adsorption curves for P-saturated Kaoline and Bentonite with and witout amorphous material.
Bentonite with and witout amorphous material.
Fig. (92) I am a larger time and a Comp. A state of the s
Fig. (82) Ion adsorption curves for P-saturated Aswan fine and 150
coarse clay with and without amorphous material.
Fig. (83) Ion adsorption curves for P-saturated El-menia fine and 151
coarse clay with and without amorphous material.
Fig. (84) Ion adsorption curves for P-saturated Fayium fine and 152
coarse clay with and without amorphous material.
Fig. (85) Ion adsorption curves for P-saturated Tanta fine and 157
coarse clay with and without amorphous material.
Fig. (86) Ion adsorption curves for P-saturated Kafr El-Sheikh 154
fine and coarse clay with and without amorphous material.
Fig. (87) Ion adsorption curves for P-saturated Belbeis fine and 156
coarse clay with and without amorphous material.
Fig. (88) Ion adsorption curves for P-saturated Nubaryia fine and 157
coarse clay with and without amorphous material.

159
160
161
162
163
164
165
166
167

LIST OF TABLES	
Table (1) Some properties of reference clay minerals.	7
Table (2) Points of zero charges of the main soil minerals as	27
reported in references.	
Table (3) Main physical and chemical characteristics of the	32
investigated soils.	
Table (4) The distribution of amorphous materials as percentage	40
of the clay fraction (%).	
Table (5) Total surface area of the clay fractions before and after	40
the removal of amorphous materials (m ² /g)	
Table (6) Semiquantitative mineralogical composition of the	42
studied clay separates.	
Table (7) Charge characteristics of the investigated clay separates	56
as calculated from potentiometric titration curves.	
Table (8) Charge parameters of the investigated clay separates as	75
calculated from Ion-adsorption curves.	
Table (9) Gaines – Thomas (K _{GT}) selectivity coefficients of fine	95
and coarse clay separates.	
Table (10) Gapon selectivity coefficients (K _G) of the studied clay	105
separates.	
Table (11) Amounts of phosphate sorbed (mg/kg) on Kaoline and	111
Bentonite at different P- initial concentrations and	
constant pH values.	
Table (12) Amounts of phosphate sorbed (mg/kg) on fine clay at	113
different P-initial concentrations and constant pH values.	
Table (13) Amounts of phosphate sorbed (mg/kg) on coarse clay	114
at different P-initial concentrations and constant pH values.	