
References

- **Alamo PO, Rodríguez TR, and Fernández RS (2005):** The effect of AM3 in the resolution of otitis media with effusion (OME) in paediatric patients: *Acta Otorrinolaringol Esp.*;56:1-5.
- **American Academy of Family Physicians, the American Academy of Otolaryngology-Head and Neck Surgery, and the American Academy of Pediatrics Subcommittee on Otitis Media with Effusion (2004):** Otitis Media with Effusion Clinical Practice Guideline; *Am Fam Physician.* ;69:2776-2779.
- **Arguedas AG, Zaleska M, Stutman HR, Blumer JL, and Hains CS: (1991)** Comparative trial of cefprozil vs. amoxicillin clavulanate potassium in the treatment of children with acute_ otitis media with effusion: *Pediatr Infect Dis J.* 10:375-80
- **Bakaletz LO (1989):** Effect of formalin-fixed *Haemophilus influenzae* and *Streptococcus pneumoniae* on dye transport by the chinchilla eustachian tube, *Acta Otolaryngol* 107:235–243
- **Balle V, Sederberg-Olsen J, Thomsen J, and Hartzen S (1998):** Treatment of children with secretory otitis media (SOM) with amoxicillin and clavulanic acid (Spektramox) or penicillin-V

- (Primcillin). Bacteriological findings in the nasopharynx before and after treatment. *Int J Pediatr Otorhinolaryngol.* 45:77-82.
- **Blanshard JD; Maw AR and Bawden R (1993):** Conservative treatment of otitis media with effusion by autoinflation of the middle ear. *Clin Otolaryngol.*, 18:188-192.
 - **Bluestone CD, Beery QC, and Paradise JL (1973) :** Audiometry and tympanometry in relation to middle ear effusions in children. *Laryngoscope* 83: 594-604
 - **Bluestone CD (1990),** *Ann Otol Rhinol Laryngol* 99:43
 - **Bluestone CD, Stool SE, Alper CM and Klien JO (2003):** Otitis media and Eustachian tube dysfunction In *Paediatric otorhinolaryngology*, 4th ed. Philadelphia: Sanders, 2003: 474-685
 - **Brown DT, Litt M and Potsic WP (1985):** A study of mucus glycoproteins in secretory otitis media. *Arch Otolaryngol.* ; 111(10):688-695.
 - **Butler CC and van Der Voort JH (2001):** Steroids for otitis media with effusion: a systematic review. *Arch Pediatr Adolesc Med.*; 155:641-647.

- **Casselbrant ML(1985)** : Otitis media in a population of black American and white American infants, 0–2 years of age, *Int J Pediatr Otorhinolaryngol* 33:1–16.
- **Casselbrant ML, Brostoff LM, Flaherty MR, Blueston CD, Doyle WJ and Canteken E (1985)**: otitis media with effusion in preschool children. *Laryngoscope.* ; 95: 428-436
- **Casselbrant ML, and Mandel EM (2003)**: Epidemiology. In: Rosenfeld RM, Bluestone CD, eds. *Evidence-Based Otitis Media*. 2nd ed: Hamilton, Ontario: BC Decker:147–162
- **Clarke S, and De RW (2005)**: otitis media in children. *Hosp Med.*; 66:268-270
- **Corwin MJ, Weiner LB and Daniels D (1986)**: Efficacy of oral antibiotics for the treatment of persistent otitis media with effusion: *International Journal of Pediatric Otorhinolaryngology* , 11:109-112
- **Coyte PC, Croxford R, McIsaac W, Feldman W and Friedberg J (2001)**: The role of adjuvant adenoidectomy and tonsillectomy in the outcome of insertion of tympanostomy tubes:*N Engl J Med.* 344:1188–1195.
- **Daly K, Giebink GS, Batalden PB, Anderson RS, Le CT, and Lindgren B(1991)**: Resolution of otitis media with effusion with

the use of a stepped treatment regimen of trimethoprim-sulfamethoxazole and prednisone: *Pediatr Infect Dis J*;10:500-6.

- **Donaldson JD, Martin GF, Maltby CC, and Seywerd EB (1990):** The efficacy of pulse-dosed antibiotic therapy in the management of persistent otitis media with effusion: *J Otolaryngol.*;19:175-8.
- **Ehrlich GD (2002):** Mucosal biofilm formation on middle-ear mucosa in the chinchilla model of otitis media: *JAMA* 287:1710–1715.
- **Erdem IC, McGuire TW, and Griffith TL (1991):** Antimicrobial Therapy for Otitis Media With Effusion ('Secretory' Otitis Media): *JAMA.* ;266:3309-3317
- **Ernstson S and Sundberg L. (1984):** Erythromycin in the treatment of Otitis Media with Effusion (OME): *The Journal of Laryngology and Otology* . Vol. 98. pp. 767-769
- **Ernstson S, and Anari M (1985):** Cefaclor in the treatment of otitis media with effusion: *Acta Otolaryngol (Stockh); Suppl.* 424: 17-21.

-
- **Eser OK, Ipci K, Alp S, Akyol U, Unal OF, Hascelik G, Sennaroglu L, and Gür D (2009):** Efficacy of nasopharyngeal culture in identification of pathogen in middle ear fluid in chronic otitis media with effusion: Indian J Med Microbiol. (3):237-41.
 - **Fergie N, Bayston R, Pearson JP, Birchall JP (2004):** Is otitis media with effusion a biofilm infection; 29(1):38-46.
 - **Fernau JL, Hirsch BE, Derkay C, Ramasastry S, and Schaeter SE (1992)** hyperbaric oxygen therapy: effect on middle ear and Eustachian tube function: laryngoscope, 102: 48-52
 - **Foshee WS (1992):** Loracarbef (LY163892) versus amoxicillin-clavulanate in the treatment of bacterial acute otitis media with effusion: , J Pediatr. ;120:980-6.
 - **Gates GA (1986):** Differential otomanometry: Am J Otolaryngol 7:147–150
 - **Gates GA (1986):** Medical treatment of chronic otitis media with effusion (secretory otitis media): Otolaryngol Head Neck Surg 94:350–354.
 - **Gates GA (1987):** Effectiveness of adenoidectomy and tympanostomy tubes in the treatment of chronic otitis media with effusion: N Engl J Med 31:1444–1451.

- **Giebink GS, Batalden PB, Le CT, Lassman FM, Buran DJ, and Seltz AE (1990):** A controlled trial comparing three treatments for chronic otitis media with effusion: *Pediatr Infect Dis J* 1990;9:33-40.
- **Gooch WM 3rd, Blair E, Puopolo A, Paster Z, Schwartz RH, Miller HC, Smyre HL, Giguere gGC, and Collins JJ. (1995):** Clinical comparison of cefuroxime axetil suspension and amoxicillin/clavulanate suspension in the treatment of pediatric patients with acute otitis media with effusion: *Clin Ther.* ;17:838-51
- **Grote JJ (1997):** Antibiotics in otitis media with effusion: *Ned Tijdschr Geneeskd.* ;141:76-7.
- **Hardy SM, Heavner SB, White DR, Prazma J and Pillsbury HC (2001):** Transient inflammatory and dysfunction of the eustachian tube secondary to multiple exposures of simulated gastroesophageal refluxant: *Ann Otol Rhinol Laryngol*; 110: 928- 934.
- **Healy GB (1984):** Antimicrobial therapy of chronic ofitis media with Effusion: *International Journal of Pediatric Otorhinolaryngology*; 8: 13 - 17

- **Hurst DS (1996):** Association of otitis media with effusion and allergy as demonstrated by intradermal skin testing and eosinophil cationic protein levels in both middle ear effusions and mucosal biopsies: *Laryngoscope* 106:1128–1137.
- **Iwona T, Kinoshita T, Hamada E, Doi T, Ushito K, and Kumazawa T(1993):**OME & Eustachian tube dysfunction in adults and children: *Acta otorhinolaryngology Supplement*, 500: 66-9.
- **Jerger J (1970):** Clinical experience with impedance audiometry: *Arch Otolaryngol* 92:311–324
- **Klein JO, Tos M, Hussl B, Naunton RF and Ohyama M (1989):** definition and classification of secretory otitis media, Panel reports 1A. *Ann Otol Rhinol. Laryngol*; 98 (Suppl. 139): 10
- **Kobayashi K (2001):** Applicability of long-term low-dosage macrolide therapy for children with serous otitis media: *Jpn J Antibiot.* Dec;54 Suppl C:26-9
- **Lai D, Li W, Xian J, and Liu S (2008):** Multifrequency tympanometry in adults with otitis media with effusion. *Eur Arch Otorhinolaryngol* ;265:1021-1025.
- **Leach AJ, Morris PS, and Mathews JD (2008):** Compared to placebo, long-term antibiotics resolve otitis media with effusion (OME) and prevent acute otitis media with perforation

(AOMwiP) in a high-risk population: A randomized controlled trial: BMC Pediatr. ; 8: 23

- **Lim DJ, and Moon SK (2011):** Establishment of cell lines from the human middle and inner ear epithelial cells. Human cell transformation advances in experimental medicine and biology; 720: 15-25
- **Low CR, Kumar V, Robins SL and Robins CM (1997):** Pathologic basis of disease. 6th ed. Philadelphia, Pa: WB Saunders: 332.
- **Mandel EM, Rockette HE, Bluestone CD, Paradise JL, and Nozza RJ (1987):** Efficacy of amoxicillin with and without decongestant- antihistamine for otitis media with effusion in children: N Engl J Med ;316:432-7
- **Mandel EM, Rockette HE, Paradise JL, Bluestone CD, and Nozza RJ (1991):** Comparative efficacy of erythromycin-sulfisoxazole, cefaclor, amoxicillin or placebo for otitis media with effusion in children: Pediatr Infect Dis J. ;10:899-906.
- **Mandel EM, Casselbrant ML, Kurs-Lasky M, and Bluestone CD (1996):** Efficacy of ceftibuten compared with

-
- amoxicillin for otitis media with effusion in infants and children: *Pediatr Infect Dis J.* ;15:409-14.
- **Mandel EM, and Casselbrant ML (2004):** Antibiotics for otitis media with effusion: *Minerva Pediatr* ;56:481-95
 - **Marchant CD, Shurin PA, Turczyk VA, Feinstein JC, Johnson CE, Wasikowski DE, Knapp LJ, and Tutihasi MA (1984):** A randomized controlled trial of cefaclor compared with trimethoprim-sulfamethoxazole for treatment of acute otitis media: *J Pediatr.*; 105(4):633-8.
 - **Marks NJ, Mills RP, and Shaheen OH (1981):** A controlled trial of cotrimoxazole in serous otitis media: *J Laryngol Otol* 95:1003-9.
 - **Marra F, Patrick DM, Chong M, and Bowie WR (2006):** Antibiotic use among children in British Columbia, *Journal of Antimicrobial Chemotherapy*; Vol. 58, issue 4:830–9.
 - **McLinn SE (1987):** Randomized, open label, multicenter trial of cefixime compared with amoxicillin for treatment of acute otitis media with effusion: *Pediatr Infect Dis J.* ;6:997-1001.
 - **McLinn SE, Moskal M, Goldfarb J, Bodor F, Aronovitz G, Schwartz R, Self P, and Ossi MJ (1994):** Comparison of cefuroxime axetil and amoxicillin-clavulanate suspensions in

- treatment of acute otitis media with effusion in children: Antimicrob Agents Chemother. ;38:315-8.
- **Odio CM, Kusmiesz H, Shelton S, and Nelson JD (1985):**Comparative treatment trial of augmentin versus cefaclor for acute otitis media with effusion: Pediatrics. ;75:819-26.
 - **Ozturk k, Gode S, Ogut F, Bilgen C, and Kirazli T (2011):** The Value of Multifrequency Tympanometry in the Management of Otitis Media with Effusion ;Int. Adv. Otol; 7: 4-8
 - **Paparella MM, Timothy TK and Goycoolea MV (1991):** Otitis media with effusion in Paparella, Shumrick, Gluckman and Myerhoff Otolaryngology, Philadelphia, W.B. Saunders Co., Vol. II, Chap. 27, pp. 1317-1340.
 - **Paradise JL, Smith GC, and Bluestone CD (1976):** Tympanometric detection of middle ear effusion in infants and children: Pediatrics 58:198–210
 - **Paradise JL, Rockette HE, and Colborn DK (1997):** Otitis media in 2253 Pittsburgh area infants: prevalence and risk factors during the first two years of life:. Pediatrics; 99 :318–333
 - **Paradise JL, Feldman HM, and Campbell TF (2007):** Tympanostomy tubes and developmental outcomes at 9 to 11

years of age. The New England journal of medicine ;356:248-261.

- **Perera R, Haynes J, Glasziou P, and Heneghan CJ(2006):** Autoinflation for hearing loss associated with otitis media with effusion.Cochrane Database Syst Rev. ;4:CD006285.
- **Poetker DM, Lindstrom DR, Edmiston CE, Krepel CJ, Link TR, and Kerschner JE (2005):** Microbiology of middle ear effusions from 292 patients undergoing tympanostomy tube placement for middle ear disease: International Journal of Pediatric Otorhinolaryngology ;69:799–804.
- **Ponsioen BP (1997):**Antibiotics in otitis media with effusion, Ned Tijdschr Geneeskd. ;141:702.
- **Post JC, Preston RA, Aul JJ, Larkins-Pettigrew M, Rydquist-White J, Anderson K W, Wadowski R M, Reagan D R, Walker E S, Kingsley L A, Magit A E, and Ehrlich G D (1995):** Molecular analysis of bacterial pathogens in otitis media with effusion. JAMA.;273:1598–1604.
- **Rayner MG (1998):** Evidence of bacterial metabolic activity in culture-negative otitis media with effusion: JAMA 279:296–299,

- **Robinson PJ, Lodge S, Jones BM, Walker CC, and Grant HR (1992):** The effect of palate repair on otitis media with effusion: *Journal of plastic surgery*, 8: 640-645
- **Rosenfeld RM and Bluestone CD (2003):** Clinical efficacy of surgical therapy. In: Rosenfeld RM, Bluestone CD, eds. *Evidence-Based Otitis Media*. 2nd ed. Hamilton, ON, Canada: BC Decker; 227–240.
- **Rosenfeld RM, Culpepper KJ, Doyle KM, Grundfast A, Hoberman MA, Kenna AS, Lieberthal M, Mahoney RA, Wahl CR, Woods Jr and Yawn BL (2004) :**Clinical practice guideline: otitis media with effusion: *Otolaryngol. Head Neck Surg*130: (Suppl. 5): S95- S118.
- **Roydhouse N (1991):** Antibiotic treatment of otitis media with effusion. *New Zealand Medical Journal*;104(919): 380–2.
- **Ryan AF (1985):** Immunologically mediated models of chronic otitis media with effusion. In Sade H, editor: *Acute and secretory otitis media*, 10: 220-234
- **Ryding M, White P and Kalm O (2004):** Eustachian tube function and tympanic memberane findings after chronic secretory otitis media. *Int J Pediatr Otorhinolaryngol*Feb; 68 (2); 197-204

-
- **Saafan ME, Ibrahim WS, and Tomoum MO (2012):** Role of adenoid biofilm in chronic otitis media with effusion in children: Eur Arch Otorhinolaryngol. 3: 50-76
 - **Sade J and Fuchs C (2003):** Middle ear gas deficiency. Is secretory otitis media a contraindication to air travel?: Ann Otol Rhinol Laryngol; 112:230-235.
 - **Safak MA, Kilic, R, Haberal I, Go"çmen H, and OÈ zeri C (2001):** A comparative study of azithromycin and pseudoephedrine hydrochloride for otitis media with effusion in children: Acta Otolaryngol ; 121: 925–929.
 - **Sade J: (1971):** Mucociliary flow in the middle ear: Ann Otol Rhinol Laryngol 80:336-431.
 - **Sando I, and Takahashi H (1990):** Otitis media in association with various congenital disease: Ann Otol Rhinol Laryngol 99:13–16
 - **Senturia BH and others (1958):** Studies concerned with tubotympanitis : Ann Otol Rhinol Laryngol 67:440–467
 - **Shekelle P, Takata G, and Chan LS (2003):** Diagnosis, Natural History, and Late Effects of Otitis Media With Effusion. Evidence Report/Technology Assessment No. 55: AHR55:3-23

- **Stool SE, Berg AO, and Berman S (1994):** Otitis Media With Effusion in Young Children. Clinical Practice Guideline, Number 12: AHCPR12:94-622
- **Takahashi H, Honjo I, and Fujita A (1996):** endoscopic findings at the pharyngeal orifice of the Eustachian tube in OME: European archives of otorhinolaryngology 253: 42-4.
- **Takahashi H, Fujita A and Honjo I (2003):** Eustachian tube compliance in cleft palates. In: Mogi G, Honjo I, Iskii T, et al (Eds): Recent Advances in Otitis Media. Amsterdam, Kugler, p. 303.
- **Teele DW, Klein JO, and Rosner B (1989):** The Greater Boston Otitis Media Study Group: Epidemiology of otitis media during the first seven years of life in children in Greater Boston: a prospective cohort study: J Infect Dis 160:83–94
- **Terris MH, Magit AE and Davidson TM (1995):** Otitis media with effusion in infants and children. Primary care concerns addressed from an otolaryngologists perspective. Postgraduate Med., 97:137-8, 143-4.
- **Tharsher RD III, Talaver F, and Meyers AD (2011):** Otitis media with effusion workup: Medscape

-
- **Thomsen J, Sederberg-Olsen J, Stangerup SE, Balle V and Vejlsgaard R(1988):** Long-term antibiotic treatment of children with secretory otitis media: A double-blind placebo-controlled study: *Acta Otolaryngol (Stockh)* ; 449: 49-50
 - **Thomsen J, Sederberg-Olsen J, Balle V, and Hartzen S (1997):**Antibiotic treatment of children with secretory otitis media. Amoxicillin-clavulanate is superior to penicillin V in a double-blind randomized study: *Arch Otolaryngol Head Neck Surg.* ;123:695-9.
 - **Thornton RB, Rigby PJ, Wiertsema SP, Filion P, Langlands J, Coates HL, Vijayasekaran S, Keil AD, and Richmond PC (2011):** Multispecies Bacterial Biofilm and intracellular Infection in Otitis Media
 - **Tos M, and Paulsen G (1979):** Tympanometry in 2-year-old children. Seasonal influence of frequency of secretory otitis and tubal function: *ORL J Oto* 41:1–10
 - **Tos M, Poulsen G, and Hancke AB(1979):** Screening tympanometry during the first year of life: *Acta Otolaryngol* 88:388–394.
 - **Tos M(1984):** Epidemiology and natural history of secretory otitis: *Am J Otol.*1984;5 :459– 462