
Body Contouring After Bariatric Surgery

Essay

Submitted for Partial Fulfillment of Master
Degree in General Surgery

Presented by

Azza Mohamed El-Sayed Mohamed

M.B, B.CH

Faculty of Medicine
Ain Shams University

Supervised by

Prof. Dr. Ashraf Abd El-moghny Mostafa

*Professor of General Surgery
Ain Shams University*

Dr. Karim Khalil El-Lamie

*Lecturer of Plastic Surgery
Ain Shams University*

**Faculty of Medicine
Ain Shams University**

2010

Contents

<i>Title</i>	<i>Page</i>
• Introduction & aim of the work	1
• Obesity	5
• Anatomical considerations	25
• Pathophysiology of skin problems after bariatric surgery	69
• Management of skin problems after bariatric surgery:	
- Contouring of the trunk	83
- Lower body lift & thigh lift	131
- Brachioplasty	160
- Upper body lift	186
- Breast reshaping procedure	195
• Summary & Conclusions	220
• References	225
• Arabic summary	—



(وَقَالَ رَبِّ أَوْزِعْنِي أَنْ أَشْكُرَ
نِعْمَتَكَ الَّتِي أَنْعَمْتَ عَلَيَّ وَعَلَى
وَالِدَيَّ وَأَنْ أَعْمَلَ صَالِحاً تَرْضَاهُ
وَأَدْخِلْنِي بِرَحْمَتِكَ فِي عِبَادِكَ
الصَّالِحِينَ)

صدق الله العظيم

سورة النمل آية [] []

Acknowledgement

*First of all, I thank **ALLAH** who had blessed me and gave me the power to finish this work,*

*I would like to express my deepest gratitude and appreciation to **Prof. Dr. Ashraf Abd El-moghny Mostafa**, Professor of general surgery, Faculty of Medicine, Ain Shams University, for his guidance, supervision, and support.*

*I would also want to express my appreciation to **Dr. Karim Khalil El-Lamie**, Lecturer of Plastic and Reconstructive Surgery, Faculty of Medicine, Ain Shams University, who was patient and considerate, saving no time and effort helping me with this work,*

Last but not least I would like to express my best regards and thanks to my family who gave me a hand while working in this research and I would like to dedicate this work to everyone who supported me to complete this work,

List of Abbreviations

BMI	Body Mass Index
BPD	Biliopancreatic Diversion
FDA	Federal Drug Administration
HDL	High Density Lipoproteins
IMF	Infra mammary fold
LAGB	Laparoscopic Adjustable Gastric Band
LCFA	Lateral Circumflex Femoral Artery
LDL	Low Density Lipoproteins
MWL	Massive weight loss
NAC	Nipple areola complex
NHLBI	The National Heart, Lung, and Blood Institute
NIDDKD ..	National Institute for Diabetes & Digestive and Kidney Diseases
PRS	Plication of anterior rectus sheath
RD	Rectus diastasis
RYGBP	Roux-en-Y Gastric bypass
SA	Suction area
SAL	Suction assisted liposuction
SES	Socio Economic Status

SFS Superficial Fascial System

SMAS Superficial Musculo Aponeurotic System

TFL Tensor Fascia Lata

UAL Ultrasound assisted liposuction

VBG Vertical Banded Gastroplasty

WHR Waist Hip Ratio

List of Figures

<i>Figure No.</i>	<i>Page</i>
Figure (1): Gastric bypass procedure	18
Figure (2): Roux-en-Y gastric bypass procedure	19
Figure (3): Illusration of Biliopancreatic diversion	20
Figure (4): Vertical Band Gastroplasty	21
Figure (5): Illustration of intestinal bypass	22
Figure (6): Laparoscopic gastric banding	23
Figure (7): Gross anatomy of fatty tissue	25
Figure (8): Cross-section through the middle of upper arm	30
Figure (9): The axillary artery and its branches	33
Figure (10): The brachial artery	34
Figure (11): Diagram of segmental distribution of the cutaneous nerves of the right upper extremity. Anterior view	36
Figure (12): Cutaneous nerves of right upper extremity Posterior view	37
Figure (13): Diagram of segmental distribution of the cutaneous nerves of the right upper extremity. Posterior view	38
Figure (14): Nerves of the left upper extremity	39
Figure (15): Lymphatics of the mammary gland, and the axillary glands	41

Figure (16):	The External, internal and transverses abdominis muscles	48
Figure (17):	The rectus sheath at various levels	48
Figure (18):	A perforator is demonstrated branching at the level of Scarpa's fascia (dotted line) and anastomosing with adjacent perforators at a subdermal level	50
Figure (19):	A perforator is demonstrated branching at the level of Scarpa's fascia (dotted line) and anastomosing with adjacent perforators at a subdermal level	50
Figure (20):	Zones of blood supply to the abdominal wall	51
Figure (21):	The anatomic regions (suction areas) for suction lipectomy in abdominoplasty: SA1 (safe), SA2 (limited), SA3 (cautious) and SA4 (unrestricted)	53
Figure (22):	The cutaneous distribution of the thoraco-abdominal nerves	54
Figure (23):	Boundaries of the thigh	57
Figure (24):	Arterial supply of the thigh: Gray's Anatomy, 1985	59
Figure (25):	Superficial veins of the lower limb	60
Figure (26):	Diagram of superficial fascial system zones of adherence (black bands, most adherent zones; white zones, least adherent)	63
Figure (27):	Diagram of superficial fascial system anatomic variations in males (left) and females (right). The superficial fascial system zone of adherence is at the iliac crest in males and several centimeters inferior to the iliac crest in females	67

Figure (28):	Hanging panniculus and hanging abdominal apron	87
Figure (29):	The female abdomen consists of seven aesthetic units	100
Figure (30):	The male abdomen consists of six aesthetic units	101
Figure (31):	Modern abdomen accepted as the standard of beauty	104
Figure (32):	The aesthetic mons pubis	105
Figure (33):	Different shapes of umbilicus (top left) T-shaped umbilicus, (top right) oval shaped umbilicus, (below left) vertical shaped umbilicus, (below right) horizontally shaped umbilicus	106
Figure (34):	(Left) Type A deformity showing rectus diastasis (RD) secondary to pregnancy. (Right) Correction of the rectus diastasis by the plication of the anterior rectus sheath	108
Figure (35):	(Left) Patient with type A deformity with rectus diastasis. (Right) One year postoperatively after plication of the anterior rectus sheath	108
Figure (36):	(Left) Type B deformity with demarcation at the area of rectus diastasis and the L-shaped and inverted-L-shaped plication of the external oblique aponeurosis. (Right) After correction of the rectus diastasis and plication of the external oblique aponeurosis	109

- Figure (37):** (*Left*) Patient with type B deformity with rectus diastasis and laxity of the inferior abdominal wall. The patient also presented with breast ptosis. (*Right*) Seven months postoperatively with improvement of the abdominal contour by plication of the anterior rectus sheath associated with an L-shaped plication of the external oblique aponeurosis. Correction of breast ptosis was also performed109
- Figure (38):** (*Left*) Type C deformity showing congenital rectus diastasis (*CRD*). (*Right*) Undermining of both rectus muscles (*RM*), exposing the posterior recti sheaths (*PS*)110
- Figure (39):** (*Above, left*) Oblique view of a patient with type C deformity with lateral insertion of the rectus muscles diagnosed at the preoperative examination. (*Above, right*) Six months after correction of the muscular layer by release and advancement of the rectus muscles. (*Below, left*) Profile view of the same patient showing the forward projection of the abdomen. (*Below, right*) Six months postoperatively, showing the repositioning of the muscular layer110
- Figure (40):** (*Left*) Type D deformity showing elevation of the external oblique muscles. (*Right*) Advancement of the external oblique muscles and correction of rectus diastasis111
- Figure (41):** (*Left*) Patient with type D deformity who wished to improve the waistline. (*Right*) Six months

after the procedure, showing improvement of the waistline caused by the external oblique muscles advancement. Plication of the anterior sheath of the rectus muscle was also performed ...111

- Figure (42):** A: Pre-op marking B: Infiltration with xylocain & adrenalin with saline C: Dissection of umbilical stump114
- Figure (43):** A: Epifascial dissection up to xiphoid process b; Repair of the diverication of recti115
- Figure (44):** a, b The patient is flexed prior to demarcation of excess tissue. A temporary suture is placed in the midline to equally distribute and fix the flaps, and a long Pitanguy flap demarcator is used to estimate amount of excess tissue116
- Figure (45):** After demarcation each flap is incised, beveling the adipose tissue117
- Figure (46):** **a, b** A cone of subcutaneous tissue is resected, allowing for the exteriorization of the umbilicus and its pedicle118
- Figure (47):** Final sutures are done by planes, with the assistant pulling the flap inwards so as to decrease lateral extension of the final scars118
- Figure (48):** Anatomical areas for suction in combined abdominoplasty- liposuction. SA 1, safe; SA 2, limited; SA 3, cautious; SA 4, unrestricted120
- Figure (49):** (*Above, left*) Design of skin excision for standard transverse abdominoplasty treats only the vertical dimension of skin excess; areas of

skin excision are shaded. (*Above, right*) Modified vertical (fleur-de-lis) abdominoplasty treats both vertical and transverse skin excision in the front of the abdomen. (*Below*) Circumferential lower body procedures treat vertical excess and partially treat transverse skin excess by translating soft tissue caudally over the widest skeletal structures123

Figure (50): (*Above, left*) Preoperative view of a 32-year-old woman with a maximum body mass index of 38 and a body mass index of 22 at the time of body contouring. (*Above, right*) View of the inverted T incision resulting from modified vertical abdominoplasty. (*Below*) Image of excised skin124

Figure (51): Technique for determining vertical triangle excision and eliminating lateral dog-ears. (*Above*) After pannus dissection and abdominal wall plication, the lateral dog-ears are worked out toward the center of the abdomen to the level of the iliac crests, and provisional closure of the lateral edges is secured with towel clamps. Only then is the lower end of the pannus excised, to prevent over resection caused by the cephalad rotation of the upper flap that occurs with medial translation. (*Below, left*) The central skin excess is gathered gently to one side, and the midline is marked (*dotted line*), outlining one side of the vertical triangular excision. (*Below, right*) The central skin excess

	is then moved contralaterally and the remaining side of the vertical triangular excision is determined	125
Figure (52):	The hip pinch test	133
Figure (53):	Preoperative markings of vertical thighplasty	139
Figure (54):	Positioning using spreader bars	141
Figure (55):	Vertical thigh lift is complete; drains are placed and exited distally	142
Figure (56):	The patient had bilateral lymphedema after vertical thigh lift, left greater than right so a venous duplex ultrasound was recommended to exclude DVT, it was negative and the swelling resolved with compressive wrapping and leg elevation	146
Figure (57):	Thigh lift scar	147
Figure (58):	Medial thigh lift with scar migration and irregular fullness of upper inner thigh	147
Figure (59):	Preoperative markings of lower body lift	151
Figure (60):	Preoperative markings of lower body lift (anterior view)	153
Figure (61):	The dissection then sweeps posteriorly in the same plane (Fig.26), leaving deep fat posterior to the iliac crest	154
Figure (62):	The SFS fibrous adherence in the trochanteric region must be released with discontinuous cannula undermining to allow distal transmission of the lifting forces	154

Figure (63):	Undermining cannula used to loosen the fibrous superficial fascial system attachments circumferentially t to the knee	155
Figure (64):	Stacked towels are temporarily laterally sutured to reduce pressure when surgery is performed on the opposite side	156
Figure (65):	Lower body lift and medial thighplasty, one year postoperative	158
Figure (66):	Presentation of upper arm excess	161
Figure (67):	A T-type brachioplasty pattern	168
Figure (68):		169
Figure (69):		172
Figure (70):	Position of examination	173
Figure (71):	The anterior and posterior marks are connected to create the outer ellipse, starting at the distal upper arm and extending onto the lateral chest wall	174
Figure (72):	Central line	176
Figure (73):	Upper arrow shows bicipital groove & lower one shows the most inferior point of the upper arm	177
Figure (74):	Retraction clamps are placed along the central line of the ellipse. Traction on these clamps during the resection will help prevent development of edema	178
Figure (75):	Once the first crosshatch mark is reached, sharp, penetrating clamps are used to bring the wound	

together at the hatch marks. All bleeding is stopped and the resected segment is closed in layers. 0-PDS is used to approximate the full thickness of tissues between the skin and the underlying muscle fascia. This includes the superficial fascial system of the arm and is the strength of the closure. The skin is approximated with interrupted 2-0 and 3-0 (Monocryl) inverted subcuticular sutures. Once the first distal segment is closed, the next more proximal segment is incised, resected, and closed in a similar manner179

Figure (76): A Z-plasty is then designed and executed, based on the axillary crease position180

Figure (77): The resection is shown after completion of the second segmental closure. If the closure is deemed too tight at this stage, then the wound is reopened and a portion of the proximal based skin-fat flap is reintroduced between the open edges of the wound181

Figure (78): Before and after brachioplasty185

Figure (79): Thoracic deformities188

Figure (80): Superior extent of the lateral breast-upper back roll is incised and taken down to the level of the of of the underlying muscle fascia. Next, a skin-fat flap is elevated192

Figure (81): (Breast profile: A ducts, B lobules, C dilated section of duct to hold milk, D nipple, E fat, F pectoralis major muscle, G chest wall/rib cage