



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ
بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ





شبكة المعلومات الجامعية



شبكة المعلومات الجامعية

التوثيق الالكتروني والميكرو فيلم



شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الالكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأفلام قد اعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15 – 20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of
15 – 25c and relative humidity 20-40 %



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



شبكة المعلومات الجامعية



بالرسالة صفحات

لم ترد بالأصل

STUDIES ON FORAGE CROPS IN EGYPT

By

Shadia Mossad Ali Shahwan

B.Sc., Tanta Univ., 1978
M.Sc., Menofiya Univ., 1987

Thesis

Submitted in Partial Fulfillment
of the Requirements for the degree of

DOCTOR OF PHILOSOPHY

IN
AGRONOMY

Department of AGRONOMY,
Faculty of Agriculture,
Menofiya University

1998

Approved by :

Prof. Dr. A. A. El-Hosary

Prof. Dr. S. M. Abdel-Aal

Prof. Dr. A. I. Ramadan

Prof. Dr. M. S. El-Haroun

Committee in Charge

Date : \ \ 1998

B
1-210

SUPERVISION COMMITTEE

Prof. Dr. Mohamed El- Sayed El- Haroun.

professor of Agronomy , Faculty of Agriculture,
Minufiya University . *El-Haroun*

Prof. Dr. Saadia Ali Shaaban.

professor of Agronomy , Faculty of Agriculture,
Minufiya University . *S.A-Shaaban*

Prof. Dr. Ahmed Mahrous Rammah.

Head of Forage crops Research program Agricultural
Research Center. *Rammah*

Prof. Dr. Ahmed Abd El-Hay El-Sayed Ali.

professor of Agronomy , Faculty of Agriculture,
Minufiya University . *A-A-Ali*

ACKNOWLEDGEMENT

The auther wishes to express his deep and sincere gratitude to Dr. M.S. El- Haroun, professor of Agronomy , Faculty of Agriculture, Menofiya University, for his keen and continuous supervision, generous guidance, his valuable criticism throughout the entire study and review of the manuscript .

The deepest gratitude to Dr.S.A. Shaaban, professor of Agronomy, Faculty of Agriculture , Menofiya University, for her supervision, and guidance.

The deepest gratitude to Dr. A.M. Rammah, Head of Forage crops Research program Agricultural Research Center, for his supervision and guidance.

The deepest gratitude to Dr. A. A. Ali , professor of Agronomy ,Faculty of Agriculture , Menofiya University, for his supervision, and guidance.

The deepest gratitude to Dr. S. M. Abdel- Aal, professor of Agronomy ,Faculty of Agriculture , Menofiya University, for providing facilities during the entire research work .


I am also greatly indebted to Dr. Z.M. Marei, Head of Forage department, Agricultural Research Center ,for providing facilities during the entire research work .

I feel deeply thankful to all Staff members and technicians of forage Research Section at Gemmeiza who extended to me helping hand and sacrificed a great deal from their time during the progress of this work.

I thank and dedicate this work to whom I adore ,my parents and my family .

CONTENTS

1-	Introduction .	1
II	Review of Literature .	2
1-	Effect of residues extract of preceding crops .	2
2-	Effect of temperature and seeding depth .	8
3-	Effect of salinity .	13
III	Materials and methods .	19
IV	Results and discussion .	24
1-	The first experiment	
A-	Effect of residues extract of summer crops on seed germination and seedling development of winter forage crops.	24
1-	Germination percentage .	24
2-	Germination rate .	26
3-	Radicle length .	27
4-	Plumule length .	29
5-	Seedling length .	29
6-	Seedling fresh weight.	31
B-	Effect of residues extract of winter crops on seed germination and seedling development of summer forage crops.	32
1-	Germination percentage .	32
2-	Germination rate .	34
3-	Radicle length .	34
4-	Plumule length .	36
5-	Seedling length .	37
6-	Seedling fresh weight .	39
	The second experiment	40
2-	Effect of temperature and seeding depth on the emergence and development of forage crops .	40
1-	Effect of temperature .	40
2-	Effect of seeding depth .	45
3-	Effect of the interaction between temperature and seeding depth .	49
	The third experiment	62
3-	Effect of different concentrations of sodium chloride , calcium chloride and isoosmolal solutions of mannitol on seed germination and seedling development of winter and summer forage crops .	62
1-	Germination percentage .	62
2-	Plumule length .	68
3-	Radicle length .	73
4-	Seedling fresh weight .	79
V-	Summary .	85
VI	References.	90
	Arabic summary.	



INTRODUCTION


INTRODUCTION

Great efforts have been oriented to increase the area occupied with forage crops as well as to develop and increase the productivity and quality, which could be gained using proper varieties and optimum practices .

It is well known that forage crops production is highly affected by many factors such as preceding crop, prevailing temperature , soil salinity, soil temperature, planting depth -----etc. So , it is necessary to manage such factors properly to achieve the yield potential .

The role of allelopathy in agriculture as well as natural ecosystems has received increased attention because of its greater effects on germination , seedling development and the economic and marketable yields. More evidence has accumulated recently to show that many plants contain germination and growth inhibitors and show that toxic compounds are produced upon the decomposition of the crop residues in the soil .

So in the light of the above mentioned factors, the present research work was conducted to find out the extent to which germination and seedling characteristic of summer and winter forage crops were influenced .



**REVIEW OF
LITERATURE**

REVIEW OF LITERATURE

The review of literature of this study will be arranged under the following topics:-

- (1) Effect of residues extract of preceding crops.
- (2) Effect of temperature and seeding depth.
- (3) Effect of salinity.

(1)Effect of residues extract of preceding crops .

Bonner (1950) found that toxic substances given out to soil by higher plants can affect the growth of the same or other species.

Guenzi and McCalla (1962) studied the effect of water-soluble substances of wheat and oat straw, soybean and sweet clover hay ,and corn and sorghum stalks on germination and growth of wheat, sorghum and corn seeds. They found that all residues contained water-soluble substances depressed germination, shoot and root growth of corn, wheat and sorghum.

The effect of water extracts of field soil and decomposing plant residues, i. e barley , rye , wheat , vetch, broccoli and sudangrass on germination and growth of lettuce was studied by Patrick et al., (1963) . They found an inhibition in germination and growth of lettuce by various extracts.

Guenzi et al, (1964) found that the water soluble extracts from forage of 2 alfalfa varieties at 3 cuttings and 6 stages of growth contained water-soluble substances reduced shoot and root growth of corn seedlings.

Hoveland (1964) studied the effects of root extracts, of 6 grasses i.e bermudagrass (*Cynodon dactylon* (L.) Pers), bahiagrass (*Paspalum notatum* Flugge), dallisgrass (*Paspalum dilatatum* Poir), johnsongrass (*Sorghum halapense* (L.) Pers), sorghum alnum (*Sorghum alnum* Parodi) and tall fescue (*Festuca arundinacea* Schreb) on germination and radicle elongation of 4 clovers, i.e white clover (*Trifolium repens* L.), ball clover (*Trifolium nigrescens* Viv.), crimson clover (*Trifolium incarnatum* L.) and arrowleaf clover (*Trifolium vesiculosum* Savi). He found that johnsongrass and sorghum alnum extracts caused the most severe reduction in germination and seedling vigor. Johnsongrass and coastal bermuda grass delayed the germination of crimson clover and alfalfa initially but did not affect the final germination count after 4 days. Tall fescue reduced the radicle length of ball and crimson clovers but did not inhibit germination.

Guenzi et al., (1967) determined the toxicity of various plant parts of corn (*Zea mays* L.), sorghum (*Sorghum vulgare* Pers.), wheat (*Triticum aestivum* L) and oats (*Avena sativa* L.) collected at harvest, to seed germination and the growth of corn and wheat seedlings. They found that wheat and corn residue extracts were more toxic to corn seedlings whereas extracts of oats and sorghum residues were more toxic to the growth of wheat seedling. The germination of wheat and corn seeds was not significantly reduced by any extract.

Peters (1968) found that water extracts from roots of fescue (*Festuca arundinacea elator* Schreb.) reduced growth of the roots and germination of rape (*Brassica nigra* L.) and birdsfoot trefoil (*Lotus corniculatus* L.).

Bieber and Hoveiand (1968) studied the phytotoxic potential of 10 plant species on crownvetch and crimson clover germination and seedling vigor. They found that water extracts (1 : 15) of virginia pepperweed, evening primrose, crabgrass and crownvetch were most toxic to seed germination. Tall fescue and weeping lovegrass were least toxic. Pepperweed residues incorporated into soil for 10 weeks were toxic to crownvetch seed germination. Kobe lespedeza residues incorporated into soil did not affect germination percentage but decreased seedling growth.

Sharma and Singh (1969) in India, studied the effect of three winter forage legumes, i.e. Egyptian clover (*Trifolium alexandrinum* Juslen.), sweet clover of senji (*Melilotus indica* All.) and pea (*Pisum sativum* L.) on the germination of the succeeding crop of maize (*zea mays* L.). They found that the preceding legumes slightly delayed the germination of maize. The speed of germination of maize were faster after follow than those sown after berseem.

Peters and Mohammed Zam (1981) found that after 10 and 14 days, germination of birdsfoot trefoil (*Lotus corniculatus* L.) and red clover (*Trifolium pratense* L.) were significantly inhibited by tall fescue (*Festuca arundinacea schreb*) extracts as compared with distilled water.