



*Ain Shams University  
Faculty of Medicine  
Diagnostic Radiology department*

# **Role of Interventional Radiology in Management of Abnormal Uterine Bleeding**

*An Essay  
Submitted for Fulfillment of the Master Degree in Diagnostic Radiology*

*Presented by*

***Mohammed Hazem Mohammed Ghaly***  
*M.B.B.Ch. Zagazig University*

*Under Supervision*

*Of*

**Prof. Dr. Maha Mohammed Abdel-Raouf**

*Prof. of Diagnostic Radiology  
Faculty of Medicine  
Ain Shams University*

**Dr. Ahmed Mohammed Hussein**

*Lecturer of Diagnostic Radiology  
Faculty of Medicine  
Ain Shams University*

*Faculty of Medicine  
Ain Shams University  
2015*

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

قالوا

لسببائك لا علم لنا  
إلا ما علمتنا إنك أنت  
العليم العظيم

صدق الله العظيم

سورة البقرة الآية: ٣٢



## *Acknowledgement*

*First and foremost, thanks to **Allah** who granted me the ability to accomplish this work,*

*Words can never express my deepest gratitude and sincere appreciation to **Prof. Dr. Maha Mohammed Abdel-Rgouf**, Professor of Radiodiagnosis, Faculty of Medicine, Ain Shams University, for her precious and close supervision, and step by step guidance throughout planning and completing this work,*

*Also my deep thanks are to **Dr. Ahmed Mohammed Hussein**, lecturer of Radiodiagnosis, Faculty of Medicine, Ain Shams University, for his persistent effort, valuable guidance and meticulous revision of this work,*

*Finally, I would like to thank my family as well as my colleges for their support and care.*



**Mohammed Ghaly**  
**Cairo 2015**

---

# Contents

---

<b>List of Abbreviations .....</b>	<b>I</b>
<b>List of Tables .....</b>	<b>II</b>
<b>List of Figures .....</b>	<b>III</b>
<b>Introduction and aim of work.....</b>	
<b>Chapter 1: Anatomy of the female genital system .....</b>	
<b>Chapter 2: Causes and pathology of abnormal uterine bleeding .....</b>	
<b>Chapter 3: Technique of interventional procedures .....</b>	
<b>Summary&amp;Conclusion.....</b>	
<b>References.....</b>	
<b>Arabic Summary .....</b>	<b>--</b>

---

## Table of Abbreviations

---

Abbreviations	
<b>AVM</b>	Arteriovenous malformation
<b>CM</b>	Centimeter
<b>CEUS</b>	Contrast-enhanced ultrasound
<b>DVT</b>	Deep venous thrombosis
<b>F</b>	French
<b>Fig.</b>	Figure
<b>GnRH</b>	Gonadotropin releasing hormone
<b>HIFU</b>	High intensity focused ultrasound
<b>IIA</b>	Internal iliac artery
<b>IR</b>	Interventional radiology
<b>IV</b>	Intravenous

<b>M</b>	Microns
<b>MHZ</b>	Megahertz
<b>MI</b>	Milliliter
<b>Mmol/kg</b>	Millimole/kilogram
<b>MRI</b>	Magnetic resonance imaging
<b>PID</b>	Pelvic inflammatory disease
<b>PVA</b>	Polyvinyl alcohol
<b>PZT</b>	Spherical lead zirconate titanate
<b>RF</b>	Radio Frequency
<b>TBA</b>	Thermal balloon ablation
<b>TOA</b>	Tubo- ovarian abcess

<b>UAE</b>	Uterine artery embolization
<b>UFE</b>	Uterine fibroid embolization
<b>UP</b>	Universal protocol
<b>US</b>	Ultrasonography

---

## List of tables

---

<b>No.</b>	<b><i>Tables</i></b>	<b>page</b>
<b>(1)</b>	Terms used to describe abnormal uterine bleeding.	<b>24</b>
<b>(2)</b>	Causes of premenopausal bleeding.	<b>25</b>
<b>(3)</b>	Causes of postmenopausal bleeding.	<b>35</b>
<b>(4)</b>	FIGO and TNM staging for uterine carcinoma.	<b>40</b>
<b>(5)</b>	Differential Diagnosis Abnormal uterine Bleeding.	<b>41</b>
<b>(6)</b>	Comparison of Various Ablation Techniques.	<b>148</b>



---

## Table of Figures

---

<i>No.</i>	<i>Fig.</i>	<b>Page</b>
<b>1</b>	Anatomy of the uterus and its relation to urinary bladder and rectum.	<b>6</b>
<b>2</b>	Divisions of the uterus.	<b>7</b>
<b>3</b>	Variations in uterine position and their terminology.	<b>10</b>
<b>4</b>	Lymph drainage of the uterus.	<b>12</b>
<b>5</b>	Uterine ligaments.	<b>13</b>
<b>6</b>	Fallopian tubes.	<b>17</b>
<b>7</b>	Proliferative endometrium.	<b>20</b>
<b>8</b>	Secretory endometrium.	<b>21</b>
<b>9</b>	Multiple Leiomyomata with partial cystic degeneration.	<b>32</b>
<b>10</b>	Sequential steps through the differential diagnosis of abnormal uterine bleeding in women of childbearing age.	<b>42</b>
<b>11</b>	Tools used to guide into the uterine arteries.	<b>49</b>

<b>12</b>	(A): Microcoils in different shapes with trombogenic synthetic fibers. (B): Embolization coils.	<b>55</b>
<b>13</b>	Gelfoam Slurry Preparation.	<b>59</b>
<b>14</b>	Microspheres.	<b>61</b>
<b>15</b>	Microscopic appearances of gelatin microspheres.	<b>61</b>
<b>16</b>	Schematic drawing of needle, sheath, dilator& guide wire set.	<b>62</b>
<b>17</b>	Schematic drawing of uterine artery catheterization.	<b>63</b>
<b>18</b>	Flush pelvic aortogram shows internal iliac artery.	<b>64</b>
<b>19</b>	(A) Digital subtraction angiogram anterior obtained with selective injection via the left internal iliac artery. (B) The origin of the uterine artery (UA).	<b>65</b>
<b>20</b>	Schematic drawing of different cobra catheter.	<b>66</b>
<b>21</b>	Robert's catheter.	<b>68</b>

<b>22</b>	UFE procedure.	<b>69</b>
<b>23</b>	<p>(A) CEUS shows contrast enhancing rim surrounding a low-echogenic center.</p> <p>(B) MRI 3 month's post-embolization shows an increased size of the vascularized zone.</p>	<b>70</b>
<b>24</b>	<p>(A) Selective catheterization (cross-over) of right uterine artery with filling of fibroid.</p> <p>(B) After embolization the contrast injection demonstrates ceased flow in the fibroid.</p>	<b>72</b>
<b>25</b>	<p>(A) Coronal T1-weighted gradient-echo MR angiography in 43-year-old woman before UFE.</p> <p>(B) Completion conventional aortogram after UFE with pigtail catheter at level of renal arteries confirms enlarged left ovarian artery (arrow).</p> <p>(C) Angulated targeted MIP from coronal T1-weighted gradient echo MR angiography depicts left lower pole renal artery (arrow) giving off the ovarian artery.</p>	<b>76</b>

<b>26</b>	Pulmonary embolism, (A, B, C) Contrast-enhanced CT scans of the thorax obtained 4 days after UFE.	<b>81</b>
<b>27</b>	(A,B) Uterine necrosis. (C) Gross pathological specimen following urgent hysterectomy.	<b>83</b>
<b>28</b>	Fibroid expulsion in a 48-year-old woman.	<b>87,88</b>
<b>29</b>	Twelve months after UFE, CT reveals the hyperdense ring to be calcium.	<b>89</b>
<b>30</b>	(A) Sagittal T2-weighted MRI reveals an enlarged uterus due to an 8 cm intramural fibroid (arrowheads). (B) Axial T2-weighted MRI depicts the relation of the fibroid to the distorted uterine cavity	<b>93</b>
<b>31</b>	Bilateral uterine artery embolization procedure.	<b>94,95</b>
<b>32</b>	(A) Sagittal T2-weighted MRI obtained 7 months after uterine fibroid embolization shows a normal-sized uterus and restoration of the uterine architecture as well as the remnant (arrow) of the embolized fibroid.	<b>96</b>

	(B) Axial T2- weighted MRI depicts the relation of the remnant (arrow) to the uterine cavity.	
33	Longitudinal axis transabdominal US image obtained prior to UFE shows an 11-cm posterior intramural uterine fibroid.	98
34	<p>(A) Mapping arteriographic image obtained after aortic injection of contrast material.</p> <p>(B) Arteriographic image obtained after selective injection of contrast material into the right uterine artery.</p> <p>(C) Arteriographic image obtained after selective injection of contrast material into the left uterine artery.</p> <p>(D) Postprocedural arteriographic image obtained after aortic injection of contrast material shows no visible flow into either uterine artery.</p> <p>(E) Follow-up sagittal transabdominal US image obtained 3 months after UAE.</p> <p>(F) Follow-up axial transabdominal US image obtained 7 months after UAE.</p> <p>(G) Color Doppler US usually shows a</p>	98,99, 100,101

	marked decrease in the blood flow to fibroids after embolization.	
<b>35</b>	<p>(A) Enhanced T1-weighted MR image obtained before uterine artery embolization shows large submucosal fibroid.</p> <p>(B) Enhanced T1-weighted MR image obtained 7 days after embolization.</p> <p>(C) Enhanced T1-weighted MR image obtained 4 months after uterine artery embolization.</p> <p>(D) Enhanced T1-weighted MR image obtained 1 year after uterine artery embolization.</p>	<b>102,103</b>
<b>36</b>	Longitudinal section of a uterine fibroid that measures 78 × 53 mm. The uterus is easily seen with the MRI view.	<b>106</b>
<b>37</b>	AP projection of the uterine fibroid with five cryoprobes that have been activated for ~2 minutes. Each MRI probe is easily seen.	<b>106</b>
<b>38</b>	A transverse section of the uterine fibroid that shows two of the five cryoprobes. They have each been activated for ~3 minutes.	<b>107</b>

<b>39</b>	Entire consumption of the uterine fibroid after ~40 minutes of treatment.	<b>107</b>
<b>40</b>	(A) An extracorporeal (FEP-BY02). (B) A transrectal (Ablatherm) high intensity focused ultrasound transducer.	<b>124</b>
<b>41</b>	Schematic of patient positioning for MRgFUS.	<b>128</b>
<b>42</b>	(A) T2-weighted axial MR image of a hypo intense fibroid in a 42-year-old patient before MRgFUS (B) T1-weighted gadolinium-enhanced axial MR image of the same fibroid immediately after treatment.	<b>129</b>
<b>43</b>	(A) T2-weighted axial MR image of a hypo intense fibroid in a 46-year-old patient before MRgFUS. (B) T1-weighted gadolinium-enhanced axial image of the same fibroid immediately after treatment.	<b>130</b>
<b>44</b>	(A) Planning coronal T2-weighted fast Spin Echo MR image shows the treatment plan. (B) Gross specimen shows central necrosis	<b>135</b>