





ثبكة المعلومات الجامعية





جامعة عين شمس

التوثيق الالكتروني والميكروفيلم



نقسم بللله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأفلام قد اعدت دون آية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15-20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of 15-25c and relative humidity 20-40 %



ثبكة المعلومات الجامعية







Study of Catecholamines Levels in Patients with Psychogenic Erectile Dysfunction

Thesis

Submitted for partial fulfillment of master degree In Drematology and Andrology

<u>By</u>

Rana Mohamed Ezz-Eldin Fareed

M.B.,B.CH
Faculty of Medicine, Cairo University

Supervised by Prof. Dr. Hosni Awad

Professor and chairman of Andrology and STD's Faculty of Medicine, Cairo University

Dr. Hamed Abdalla Hamed

Assistant Professor of Andrology and STD's Faculty of Medicine, Cairo University

Dr. Shedeed Ashour Shedeed

Lecturer of Andrology and STD's Faculty of Medicine, Cairo University

Faculty of Medicine Cairo University 2001

B .



لقاه حسرة	جارحة ا		الراب	كلية
-----------	---------	--	-------	------

" بسم الله الرحمن الرحم "

وحضر اجتماع لجنه الحكم ملى الرساله المقدمة وبالدابيب/ ريا محمد عمر المري مُوريدٍ

(١) تودائه ل. اوله لدت ان الدكتور لعفل

(٢) كروز من الدور الثاني الاحتجان الدائر سيستير في والمل مرا من المنابع مرا لمنا سوار

اجتمعت لزنه الحكم على الرسالة التكونه عن الساده الم الاستاذ الدانتور/حسى احمد كرم ما المالاررت كم مع وها مدارات لالورة (على ملائيم)

الاستاذ الدكتور الجميرين الدافل _ اسادين وما مه ايرام، لدكورة السال (حمد دامل) الاستاذ الدكتور الى صعدته والما والله مرالتاس (مري بيل ممر ها رجم)

وذلان في م المعلق لمرس م المراس على في السياعة المار د في ولمه عانيه بعدن في مم المؤكرات (١)

واستهل الباحث المناقدة بصرتر بنود الرسسساله

م ناقمه السادة لهذا الجنة الحكم في

وقررت اللينشية وسرك السالا

استاذ دكتور ر مدت درا ما

16/45

استاذ د تور حبسى كوش

ces coli Til

استاذ دكتسرر



ABSTRACT

This work aimed to study role of sympathetic over activity as a possible mechanism of Psychogenic Erectile Dysfunction. Fifty subjects were divided into two groups, group A included 30 patients with Psychogenic Erectile Dysfunction as estimated by normal clinical response to Intracavernosal injection test and RigiScan monitoring, group B included 20 subjects with normal erectile function.


The 24 hour urine have been collected for each subject in both groups. The VMA levewas measured. There was no significant difference in the level of VMA between both groups, P value >0.05.

KEY WORDS

Catecholamines

Psychogenic

Erectile Dysfunction



AKNOWLEDGMENT

I would like to express my deepest gratitude and sincere appreciation to Prof. Dr. Hosny Awad, Professor and chairman of Andrology and STD's, Cairo Universit, for his wise guidance and detailed supervision, and for his encouragement to complete this work.

I am deeply indebted to Dr. Hamed Abdalla Hamed, Assistant Professor of Andrology and STD's, Cairo University, who was very helpful, reassuring and encouraging. I am greatly thankful to him for his expert help, valuable advices and kind supervision; for his patience, trustful help and sincerely effort to pursue this work.

I wish to express my deepest thanks and appreciation to Dr. Shedeed Ashour Shedeed, Lecturer of Andrology and STD's, Cairo University for his cooperation to create a good work.

My deepest gratitude to Dr. Amal El-Saied, Lecturer of Clinical Pathology, Cairo University for her precious assistance in performing the lab work.

Cairo University, for their help and cooperation throughout the work.

Finally, I'm grateful to the patients for whom all our efforts are devoted.

* * * * * *