



Screening of Developmental Dysplasia of the Hip in Neonates with Breech Presentation

Thesis

Submitted For Fulfillment of
Ph. Degree in Childhood Studies Medical Department
(Child Health and Nutrition)

By

Samar Mohamad Ezz Al-Din Almakrawy

(M.B.; B.Ch., M.Sc. Pediatrics)

Supervised By

Dr. Gamal Samy Ali

Professor of Pediatrics

Institute of Postgraduate Childhood Studies
Ain-Shams University

Dr. Hanan Mohamad Aissa

Professor of Radiodiagnosis

Faculty of medicine - Ain-Shams University

Dr. Ossama A. El Shazly

Assist. Professor of Orthopedic surgery

Faculty of medicine - Ain-Shams University

***Institute of Postgraduate Childhood Studies
Ain Shams University***

2015



معهد الدراسات العليا للطفولة
قسم الدراسات الطبية

مسح تشخيصى لتشوه نمو مفصل الفخذ فى الأطفال حديثى الولادة ذوي وضع المقعدة

خطة دراسة توطئه للحصول على درجة الدكتوراة الفلسفية فى دراسات الطفولة
القسم الطبى (صحة و تغذية الطفل) معهد الدراسات العليا للطفولة
مقدمة من

الطبيبة / سمر محمد عز الدين محمود المكاوى
ماجستير طب الأطفال - جامعة عين شمس
تحت إشراف

الأستاذ الدكتور / جمال سامى على

أستاذ طب الأطفال - قسم الدراسات الطبية
معهد الدراسات العليا للطفولة - جامعة عين شمس

الأستاذة الدكتورة / حنان محمد عيسى


أستاذ الأشعة التشخيصية
كلية الطب - جامعة عين شمس

الأستاذ الدكتور / أسامة عبد الرؤوف الشاذلى

أستاذ مساعد جراحة العظام
كلية الطب - جامعة عين شمس

معهد الدراسات العليا للطفولة - جامعة عين شمس

2015



TO

My Mother Soul ♥

My Father ♥

My Dear Uncle Dr. Saied ♥

My Brother ♥

My Sisters ♥


My Niece Salma ♥

My Nephews ♥

And My Dear Friend

Amany ♥

Acknowledgments



بسم الله الرحمن الرحيم

In the name of ALLAH, the most beneficent, the most merciful.

*It is a pleasure to express my deepest thanks and profound gratitude to **Prof. Gamal Samy** Professor of Pediatrics, Institute of Postgraduate Childhood Studies, Ain Shams University for helping me to choose the subject of this work and for his endless support and sincere advices. It has been an honor and a privilege to work under his generous supervision.*

*My deep appreciation and deepest sense of obligation to **Prof. Hanan Aissa** Professor of Radio diagnosis, Faculty of Medicine, Ain Shams University for her valuable advices, enriching observations and great effort in the practical portion which was very essential for this work.*

*My utmost thanks and deepest gratitude and appreciation to **Ass. Prof. Ossama Elshazly** Associate Professor of Orthopedic surgery, Faculty of Medicine, Ain Shams University for his meticulous scientific supervision and valuable remarks during every step of this work.*

*Words cannot suffice to express my thanks and sense of obligation to my dear friend and my senior in neonatology and ultrasound **Dr. Amany Reda** fellow pediatrics, NICU, Ain Shams university for*

her great help in the practical part of this work without which this work could never have seen the light.

To all my patients and their families who were patience with us.

To my colleagues in the NICU, Ain Shams University and to all who shared in a way or another in the successfulness of this work,

*Last but not least, my thanks to **my father** and **all my family** for their great assistance and support.*



Contents

	Page
List of Abbreviations	i
List of Tables	ii
List of Figures	iv
Introduction	1
Aim of the work	4
Review of literature	
I. Chapter (1) <i>Developmental dysplasia of the hip</i>	5
Definitions of DDH.....	5
Anatomy and Embrology of DDH	9
Aetiology and Risk Factors of DDH.....	13
Frequency of DDH.....	17
Pathophysiology of DDH	
Pathoanatomy of DDH.....	18
Natural history of DDH	19
Breech Presentation	20
II. Chapter (2)	22
Clinical and Ultrasonographic Screening of DDH.	22
Clinical examination (Screening) of DDH.....	24
The Neonate.....	28
The infant.....	30
The Walking child	34
Ultrasonographic examination (Screening) of DDH	36
Types of Ultrasonographic Screening	38
Time of Ultrasonographic Screening.....	40
Equipment Specifications	42
Examination Specifications	43
Methods of Examination.....	47
Static Method.....	47
Dynamic Method	56

III. Chapter (3)	60
Treatment of DDH	60
Newborns and Infants younger than 6 months...	62
Hip Splints	63
Pavlic Harness.....	65
Children 6 months to 2 years of age	71
Closed Reduction.....	72
Children older than 2 years.....	73
Open Reduction	75
Complications	78
Subjects and Methods	80
Results	88
Discussion	109
Summary	126
Conclusion	130
Recommendations	131
References	132
Arabic summary	

List of Abbreviations

AAP	: American Academy of Paediatrics.
ACR	: American College of Radiology
AAOS	: American Academy of Orthopedic Surgeons
AIUM	: American Institute of Ultrasound in Medicine
AVN	: Avascular Necrosis.
CDH	: Congenital dislocation of the hip.
C.S.	: Caesarean section.
DCI	: Dynamic coverage index
DDH	: Developmental Dysplasia of the hip.
Exam.	: Examination
FH	: Family history.
Fig.	: Figure
GA	: Gestational age
mos.	: Months
NICU	: Neonatal Intensive Care Unit.
No.	: Number
NVD	: Normal Vaginal Delivery.
PG	: Primigravida.
SMAC	: Standing Medical Advisory Committee
USG	: Ultrasonography
US	: Ultrasound
USPSTF	: United States Preventive Services Task Force

List of Tables

Table No.	Title	Page
1	Risk factors associated with DDH that used to inform selective DDH screening policies	13
2	Screening programme ideals and difficulties encountered with DDH	24
3	Early management for suspected DDH	42
4	Graf's morphologic hip classification (simplified)	52
5	Relationship of the Graf stage and hip morphology in DDH	54
6	Graf's Classification System of DDH, Based on sonographic angles of the Hip	86
7	Base line qualitative and quantitative descriptive demographic data of the included populaion	89
8	Base line qualitative and quantitative descriptive demographic data of the DDH diagnosed cases	91
9	The numbers of hips defined as normal or pathological on clinical examination versus ultrasonography.	92
10	Relation between clinical examination and 1 st USG screening results or (effect of presence of positive clinical examination and 1 st USG screening results).	94
11	Relation between clinical examination and 2nd USG screening results	95
12	1 st USG screening results	96

Table No.	Title	Page
13	2 nd USG screening results	97
14	Distribution of hips by Graf type and gender	97
15	Effect of the amount of amniotic fluid on occurrence of DDH	99
16	Relation between Sex and occurrence of DDH.	100
17	Effect of prematurity on occurrence of DDH	100
18	Effect of LBW on occurrence of DDH	101
19	Effect of pregnancy status on occurrence of DDH	101
20	Effect of maternal parity on occurrence of DDH.	102
21	Effect of mode of delivery on occurrence of DDH.	102
22	Effect of presence of positive family history on occurrence of DDH.	103
23	Effect of presence of postural molding conditions on occurrence of DDH.	103
24	Comparing between patients with DDH and patients with physiologically immature hip as regard risk factors using Chi-square tests.	104

List of Figures

Figure No.	Title	Page
1	Illustrative Figures of anatomy of normal hip joint and DDH.	8
2	The spectrum of anatomic abnormalities of the hip that can occur with DDH.	8
3a, 3b	Normal anatomy of the Hip Joint.	10
4a, 4b	Types and variations of Breech Presentation	21
5	Tests to evaluate hip stability	27
6	Barlow provocative test	28
7	The Ortolani maneuver	29
8	Limitation of abduction	31
9	Asymmetric thigh skin folds	33
10	positive Galeazzi sign	33
11	Kliscic test	34
12 a b c	Coronal view of the hip in the standard plane Coronal anatomic illustration Probe position in coronal view Coronal ultrasound image	45
13 a b c	Transverse view of the hip flexed 90° Transverse anatomic illustration Transverse flexion view. Coronal ultrasound image.	46
14 a b	Special positioning apparatus. USG exam. of the right hip of a baby.	48
15	Coronal ultrasound section through the acetabulum	49
16	Standard coronal ultrasound section through the acetabulum of a normal hip	51

Figure No.	Title	Page
17	Coronal US image of the Lt hip. The femoral head is sublimed with approximately 10% coverage of the femoral head	56
18	Technique for evaluating the neonatal hip using U/S	58
19	Screening for DDH; clinical algorithm	59
20	The Pavlik harness	67
21	Short leg hip spica cast	71
22	Diagrammatic illustration of the safe zone of Ramsey.	72
23	Arthrogram of a reduced hip for evaluating the stability of reduction.	73
24	Treatment algorithm for DDH according to different age groups	77
25	Barlow test and Ortolani test	83
26 a	Normal hip ultrasound demonstrating alpha and beta angles.	84
26 b	US machine	85
c	Hip US exam. of breech neonate	
27	Sex distribution in the studied neonates.	90
28	Sensitivity and Specificity of Clinical Examination in the prediction of 1 st ultrasound screening results.	93
29	Relation between clinical examination and 1 st USG screening results.	94
30	Relation between clinical examination and 2nd USG screening results.	95
31	Relation between oligohydramnios and DDH	99

Figure No.	Title	Page
32	Sonogram of (a) unilateral left sided grade III DDH case and (b) normal right side	106
33	Sonogram shows grade IV DDH case	107
34	Sonogram shows bilateral grade III case	108

Introduction

The term Developmental dysplasia of the hip (**DDH**) describes a range of hip abnormalities affecting the newborn in which the femoral head and acetabulum are in improper alignment or grow abnormally, or both (*Shipman, 2006*).

Clinical instability of the hip is the traditional hallmark of the disorder, but the definition of DDH also includes hips with radiological abnormalities of the femoral head or acetabulum that may or may not be associated with joint instability (*Dezateux, 2007*).

DDH is one of the most common congenital malformations and it is an important cause of childhood disability (*Gelfer and Kennedy, 2008*). Uncorrected DDH is associated with long term morbidity such as gait abnormalities, chronic pain and degenerative arthritis (*Shorter et al, 2011*).

The exact incidence of DDH is difficult to determine because of a discrepancy in definition of the condition, type of examination used and different levels of skills of clinicians. The international incidence ranges from *as low as 1 per 1,000 to as high as 34 per 1,000*. Higher incidences are reported when ultrasonography is also used in addition to clinical examination (*Noordin et al, 2010*).

The precise cause of DDH is unknown, with a combination of genetic and environmental influences associated with DDH and hip dislocation including family

history, oligohydramnios, vaginal delivery, breech presentation and female gender (*Sewell et al, 2009*).

Among these, breech presentation was found to be one of the most important (*Yau et al, 2012*). Breech presentation has been explained as a risk factor for DDH in terms of the intrauterine posture involved, in which leg movement is restricted and in which knee extension can stretch the hamstring and thereby increase the possibility of hip dislocation (*Dogruel et al, 2008*). The American Academy of paediatrics (AAP) now recommends ultrasound DDH screening of all female breech babies (*POSNA 2013*).

The early diagnosis of DDH is vital as timely initiation of appropriate treatment minimizes the associated risk of early osteoarthritis (*Clarke et al, 2012*). Early screening has the potential to prevent long term hip dysplasia and arthritis requiring hip replacement (*Shorter et al, 2011*). No first-line method exists for diagnosing DDH during the newborn period. However, a careful physical examination is recommended as a screening tool, particularly for high-risk infants (*Storer et al, 2006*).

Ultrasound scanning is the investigation of choice to evaluate DDH in infants younger than six months of age and it is useful to diagnose more subtle forms of the disorder when clinical examination is equivocal. It is also the only imaging modality that enables a three dimensional real-time image of a neonate's hip (*Noordin et al, 2010*).

There is general consensus that early detection of DDH improves outcome. When detected early the soft tissues are