



# **Morphological Plasticity of Some Moss Species in Response to Various Environmental Conditions via *In vitro* Culture**

**A thesis**

Submitted for the degree of Master of Science as a partial fulfillment for the requirements for the Master of Science

(Botany)

*By*

**Mohamed Farag Abu Elhamd Ali**

B. Sc. - Botany (2010)

*Supervisors*

**Prof. Wagieh El-Sayed El-Saadawi**

Professor of Botany

Faculty of Science

Ain Shams University

**Prof. Hanaa Mostafa Shabbara**

Professor of Bryology

Faculty of Science

Ain Shams University

**Dr. Hanan Farid Mahmoud Shaban**

Lecturer in Department of Botany

Faculty of Science

Ain Shams University

**Botany Department**

**Faculty of Science**

**Ain Shams University**

**(2015)**



**Student Name:** Mohamed Farag Abu Elhamd Ali

**Thesis Title:** “Morphological Plasticity of Some Moss Species in Response to Various Environmental Conditions via *In vitro* Culture”

**Degree:** Master of Science (Botany)

**This thesis has been supervised by:**

- **Prof. Wagieh El-Sayed El-Saadawi**

Professor of Botany

Botany Department - Faculty of Science

Ain Shams University

- **Prof. Hanaa Mostafa Shabbara**

Professor of Bryology

Botany Department - Faculty of Science

Ain Shams University

- **Dr. Hanan Farid Mahmoud Shaban**

Lecturer in Department of Botany

Botany Department - Faculty of Science

Ain Shams University



## Approval Sheet

### Title of Thesis:

Morphological Plasticity of Some Moss Species in Response to Various Environmental Conditions via *In vitro* Culture

**Name of Student:** Mohamed Farag Abu Elhamd Ali

**Degree:** Master of Science (Botany)

**This thesis has been approved by:**

- **Prof. Nady A. El-Bassuoni Ghanem**  
Professor of Ecology and Cryptogamic plants  
Faculty of Science (Boys) – El-Azhar University
- **Prof. Maged Mahmoud Abo Elenein**  
Professor of Plant Taxonomy  
Faculty of Education - Ain Shams University
- **Prof. Wagieh El-Sayed El-Saadawi**  
Professor of Botany - Faculty of Science  
Ain Shams University
- **Prof. Hanaa Mostafa Shabbara**  
Professor of Bryology - Faculty of Science  
Ain Shams University

**Prof. Maher Mohamed Shehata**

Head of Botany Department  
Faculty of Science  
Ain Shams University



﴿رَبِّ أَوْزِعْنِي أَنْ أَشْكُرَ نِعْمَتَكَ الَّتِي

أَنْعَمْتَ عَلَيَّ وَعَلَىٰ وَالِدَيَّ وَأَنْ أَعْمَلَ

صَالِحًا تَرْضَاهُ وَأَدْخِلْنِي بِرَحْمَتِكَ فِي

عِبَادِكَ الصَّالِحِينَ﴾

النمل: ١٩

## *Declaration*

This thesis has not been previously submitted for any degree at this or any other university. The references in the text will show specifically the extent to which I have availed myself of the work of other authors.

*Mohamed Farag*

# Acknowledgements

*First of all, the ultimate praise gratitude and thanks be to **Allah** for granting me the ability to accomplish this work.*

*I would like to express my deep thanks and gratitude to **Prof. Wageih El-Saadawi** for supervising this work, effort to facilitate its production and critical revision of the manuscript. I am deeply grateful to **Prof. Hanaa Shabbara**, for supervising this work, valuable suggestions encouragement during this work, kind help and patience. My appreciation is for **Dr. Hanan Shaban**, for her experience to facilitate the production of this work, kindness and friendship.*

*"I will always be in debt for their great help and support".*

*Thanks are also to **Dr. Usama Abou-Salama** "ex-supervisor" for his advice at the beginning of the work.*

*Thanks and gratitude are to **Prof. Maher Mohamed Shehata**, Prof. of Genetic Engineering and Head of Botany Department, Faculty of Science, Ain Shams University, and to **Prof. Mohamed Sayed Tantawy** ex-head of the Botany Department for their support and continuous help.*

*Special thanks are offered to my lab mates **Dr. Manal Ibrahim**, **Dr. Sahar Ibrahim** and **Dr. Mai Taha**. They have taught me all principles of research work, how to deal with mosses, as well their useful assistance and continuous guidance throughout this work.*

*I wish to express my gratitude to the following people who generously contributed relevant literature and for their open discussions and help in*

*this work: **Dr. Hesham Abd El Fat`tah** (Botany Department, Faculty of Science, Ain Shams University) for his help in the identification of algal strains, **Dr. Mahmoud Magdy** (Genetics Department, Faculty of Agriculture, Ain Shams University) for his great efforts in teaching me phylogentic programs and how to identify mosses based on molecular analysis. Special thanks to **Dr. Olaf Werner** (Murcia University, Spain) for explaining the basics of molecular systematics and repeating the analysis of two sequences in his lab, as well as to **Dr. Sherif Zaki** (Microbiology Department, Faculty of Science, Ain Shams University) for his help in finding out the source of contamination in culture media at beginning of the work.*

*Thanks to all members of the Botany Department for their support help and encouragement.*

*All my respectful gratitude is to my family; my mother, father, brother and sisters for their support through my life steps and to the recent source of happiness in my life, my wife Randa.*

*Mohamed Farag*

# List of Contents

Item	Page
List of Figures	iv
List of Plates	v
List of Tables	xi
Abbreviations	xii
Preface	xiii
Abstract	1
I. Introduction	2
II. Aim of the Work	8
III. Materials and Methods	10
A. Materials	10
1. Plant materials	10
a. Fresh sample	10
b. Herbium sample	11
2. Soil	11
3. Nutrient media	13
a. Solid media	14
b. liquid media	15
B. Methods	16
1. Developing of <i>in vitro</i> cultivated gametophores and investigating the morphological plasticity in response to SWHC and/or nutrient concentrations	16



a. <i>B. argenteum</i> and <i>B. subapiculatum</i>	16
b. <i>F. hygrometrica</i>	19
c. <i>P. hastata</i>	20
<b>2. Biomass Production</b>	<b>21</b>
<b>3. Molecular analysis for identification</b>	<b>22</b>
a. DNA extraction	22
b. PCR reaction	23
c. Sequencing of PCR products	24
d. Sequences alignment	24
e. Tree construction	25

#### **IV. Results** **26**

<b>A. Evaluation of the developed <i>in vitro</i> cultivation technique</b>	<b>26</b>
<b>B. Morphological Plasticity of the studied species</b>	<b>28</b>
1. <u><i>Bryum argenteum</i></u>	<b>28</b>
a. Response to different percentages of SWHC	28
b. Response to nutrient concentrations of solid media	35
2. <u><i>Bryum subapiculatum</i></u>	<b>49</b>
a. Response to different percentages of SWHC	49
b. Response to nutrient concentrations of solid media	54
3. <u><i>Funaria hygrometrica</i></u>	<b>70</b>
Response to different percentages of SWHC	70
4. <u><i>Philonotis hastata</i></u>	<b>81</b>
Response to nutrient concentrations of solid media	81
<b>C. Biomass Production</b>	<b>95</b>
<b>D. Molecular Analysis</b>	<b>103</b>

#### **V. Discussion** **108**

<b>A. The <i>in vitro</i> cultivation of mosses</b>	<b>108</b>
<b>B. Growth of mosses on media</b>	<b>112</b>
<b>C. The Plasticity of Protonema</b>	<b>113</b>

<b>D. The Plasticity of gametophores</b>	<b>116</b>
<b>VI. Conclusions and Recommendations</b>	<b>122</b>
<b>VII. Summary</b>	<b>123</b>
<b>VIII. References</b>	<b>126</b>
<b>Appendices (1-6)</b>	<b>140</b>
<b>Glossary</b>	<b>146</b>
الملخص	٢
المستخلص	١

# List of Figures

No.	Title	Page
1	The gel of genome DNA for the samples under study. The samples in wells as follow; 1. <i>Funaria hygrometrica</i> , 2. <i>Bryum subapiculatum</i> , 3. <i>Philonotis hastata</i> and 4. <i>Bryum argenteum</i> .	103
2	Neighbour-joining phylogenetic reconstruction of the sample (Ba mf 2014) with 21 <i>Bryum</i> taxa, 3 <i>Brachymenium</i> taxa, <i>Mielichhoferia macrocarpa</i> and <i>Pohlia flexuosa</i> –as an out group- based on chloroplast <i>rbcL</i> sequences.	104
3	Neighbour-joining phylogenetic reconstruction of the sample (Ba mf 2014) with 12 <i>Bryum</i> taxa (multi-sequences for some taxa) and <i>Pohlia proligera</i> as an out group based on nuclear ribosomal ITS sequences.	105
4	The chromatograms of sequences of <i>B. subapiculatum</i> , the upper one of <i>rbcL</i> marker and the lower one of ITS marker.	106
5	Neighbour-joining phylogenetic reconstruction of the sample (Ph mf 2014) with 6 <i>Philonotis</i> taxa and <i>Breutelia chrysocoma</i> as an out group based on chloroplast <i>rbcL</i> sequences.	106
6	Neighbor-joining phylogenetic reconstruction of the sample (Ph mf 2014) with 14 <i>Philonotis</i> taxa (multi-sequences for some taxa) and <i>Breutelia chrysea</i> as an out group based on nuclear ribosomal ITS sequences.	107

# List of Plates

No.	Title	Page
1	The growth of <i>Bryum argenteum</i> under three different percentages (33%, 66% and 99%) of soil water holding capacity through different times of the experiment (i.e. after 1, 2, 4 and 6 weeks). (a-d) 33% SWHC: the covering percentage increased gradually to be about 15%, 30%, 50% and 75% -80%; (e-h) 66% SWHC: the covering percentage increased gradually to be about 15%, 35%, 70% and 85% -90%; (i-l) 99% SWHC: the covering percentage increased gradually to be about 15%, 35%, 65% and 85%.	31-32
2	Differences in the branching pattern of gametophores of <i>Bryum argenteum</i> after six weeks. (a-b) under half maximum percentage (33%) of SWHC showing numerous short young and long old branches; (c-d) under maximum percentage (66%) of SWHC showing a few short young and numerous long old branches; (e-g) under flooding percentage (99%) of SWHC showing long old branches and etiolated solitary gametophore.	34
3	Protonema morphogenesis of <i>Bryum argenteum</i> grown on nutrient free (NFA) medium. (a) protonema; (b) two rhizoids different in diameters; (c) pigmented multicellular chloronema gemma; (d) two rhizoids showing different peculiar structures on them.	37
4	Gametophores of <i>Bryum argenteum</i> grown on nutrient free (NFA) medium after 10 weeks. (a) numerous developed gametophores in the media from the single inoculated gametophore; (b) six gametophores attached at base; (c) apices of gametophores, arrows refer to leaves different in morphology; (d) parts of two magnified gametophores having different colours, arrow refers to two leaf apices, having different shapes.	38
5	Morphology of leaves of numerous gametophores of <i>Bryum argenteum</i> grown on nutrient free (NFA) medium, showing variation in leaf shape, leaf apex and costa between them.	39

<b>6</b>	Protonema morphogenesis of <i>Bryum argenteum</i> grown on half strength MS (HsMS) medium. (a) normal chloronema filament; (b) gemmiferous chloronema filament; (c) flask-shaped cell in chloronema filament (act as detaching point); (d) division of flask-shaped cell; (e) another form for detaching point of gemmiferous filaments by tmema cells; (f) two scars after separation at tmema cells.	<b>42</b>
<b>7</b>	Morphology of leaves of numerous gametophores of <i>Bryum argenteum</i> grown on half strength MS (HsMS) medium, showing leaves of similar morphology.	<b>43</b>
<b>8</b>	Protonema morphogenesis of <i>Bryum argenteum</i> grown on full strength MS (FsMS) media. (a) cluster of protonema, black arrow refers to chloronema filament, red arrow refers to gemmiferous filament; (b) cluster of protonema, arrow refers to tmema cells; (c) protonemal filament showing transition from caulonema to chloronema; (d) magnified chloronema filament; (e) magnified caulonemal filament; (f) division of chloronema cells to form brachyctes; (g) four brachyctes.	<b>46</b>
<b>9</b>	Morphology of leaves of numerous gametophores of <i>Bryum argenteum</i> grown on full strength MS (FsMS) medium, showing variation of one gametophore from the other in leaf size, leaf apex and costa.	<b>47</b>
<b>10</b>	Surface view of jars of <i>Bryum argenteum</i> of each replicate of the NFA, HsMS and FsMS media showing the pattern of growth after 10 weeks. Red point refers to cultures contaminated with bacteria in HsMS medium and with fungi in FsMS medium.	<b>48</b>
<b>11</b>	The growth of <i>B. subapiculatum</i> under three different percentages (33%, 66% and 99%) of soil water holding capacity through different times of the experiment (i.e. after 20, 30, 40 and 50 days). (a-d) 33% SWHC; showing the weak growth of protonema after 40 days and onwards; (e-h) 66% SWHC; showing vigours growth of protonema after 20 days and a white and a yellowish green destroyed protonema after 30 days and onward as a result of contamination by either cyanobacteria or diatoms; (i-l) 99% SWHC; showing moderate growth of protonema after 20 days and	<b>51-52</b>

	a yellowish green protonema, which is partially or completely covered by diatoms colonies after 30 days and onward.	
<b>12</b>	Protonema morphogenesis of <i>Bryum subapiculatum</i> grown on nutrient free (NFA) medium. (a) chloronema filament; (b) caulonema filament; (c) caulonema filament showing a branching initiation point before septum; (d) pigmented rhizoids, showing a branching initiation point before septum and the reduction in a diameter of new branches; (e) caulonema filament carrying branched chloronema filaments.	<b>56</b>
<b>13</b>	Gametophores of <i>Bryum subapiculatum</i> grown on nutrient free (NFA) media after ten weeks. (a) numerous gametophores developed from the single inoculated gametophore, arrows refer to gametophore developed inside medium; (b-c) stunted gametophores with axillary red rhizoids, (d) etiolated gametophore.	<b>57</b>
<b>14</b>	Morphology of leaves of numerous gametophores of <i>Bryum subapiculatum</i> grown on nutrient free (NFA) medium, showing variation in leaf morphology between the gametophores.	<b>58</b>
<b>15</b>	Protonema morphogenesis of <i>Bryum subapiculatum</i> grown on half strength MS (HsMS) medium. (a) chloronema filament; (b) a cluster of protonemal gemmae borne terminally on chloronema filament, arrow refers to point of detaching; (c) magnified multicellular chloronema gemma.	<b>61</b>
<b>16</b>	Gametophores of <i>Bryum subapiculatum</i> grown on half strength MS (HsMS) medium after 10 weeks. (a) the protonema filaments developed from the single inoculated gametophore, arrows refer to protonema gemmae; (b-d) the developed gametophores.	<b>62</b>
<b>17</b>	Morphology of leaves of numerous gametophores of <i>Bryum subapiculatum</i> grown on half strength MS (HsMS) medium, showing similarty in leaf shape and differences in leaf length between the gametophores.	<b>63</b>
<b>18</b>	Protonema morphogenesis of <i>Bryum subapiculatum</i> grown on full strength MS (FsMS) medium. (a) two chloronema filaments with different diameters; (b) caulonema filament branched into chloronema filaments; (c) two clusters of protonemal gemmae	<b>66</b>

	borne terminally on chloronema filaments; (d-e) branching of chloronema filaments with chloronema gemmae.	
<b>19</b>	Gametophores of <i>Bryum subapiculatum</i> grown on full strength MS (FsMS) medium after 10 weeks. (a) the protonema filaments developed from the single inoculated gametophore, arrows refer to protonema gemmae; (b-d) the developed gametophores.	<b>67</b>
<b>20</b>	Morphology of leaves of numerous gametophores of <i>Bryum subapiculatum</i> grown on full strength MS (FsMS) medium, showing high degree of similarity in leaf morphology.	<b>68</b>
<b>21</b>	Surface view of jars of <i>Bryum subapiculatum</i> of each replicate of the NFA, HsMS and FsMS media showing the pattern of growth after 10 weeks. Red points refer to cultures contaminated with bacteria in NFA media, with algae in HsMS medium and with fungi in FsMS medium.	<b>69</b>
<b>22</b>	The growth of <i>Funaria hygrometrica</i> grown under three different percentages (33%, 66% and 99%) of soil water holding capacity through different times of the experiment (i.e. after 20, 30, 40 and 50 days). (a-d) 33% SWHC: showing the absence of any contamination and the increase of new developed gametophores; (e-h) 66% SWHC: showing the appearance of cyanobacterial contamination (red arrows) which increased from (e) to cover 100% of the soil surface in (h) and delayed the development of new gametophores; (i-l) 99% SWHC; i) showing a flourishing of cyanobacteria growth on all soil surface and its high effect in the retardation of the development of new gametophores, (j, k, l) submerging the protonema by <i>Nostoc</i> filaments and stopping the development of new gametophores.	<b>71-72</b>
<b>23</b>	Morphology of leaves of numerous gametophores of <i>Funaria hygrometrica</i> grown under half maximum percentage (33%) of SWHC, showing variation in leaf morphology.	<b>74</b>
<b>24</b>	Morphology of leaves of numerous gametophores of <i>Funaria hygrometrica</i> grown under maximum percentage (66%) of SWHC, showing some similarity in leaf morphology.	<b>77</b>
<b>25</b>	Morphology of leaves of numerous gametophores of <i>Funaria hygrometrica</i> grown under flooding percentage (99%) of SWHC	<b>80</b>