

Creatine Kinase as a Biochemical Marker in Diagnosis of Placenta Increta and Percreta

Thesis

Submitted as Partial Fulfillment of MD Degree
in Obstetrics and Gynecology

By

Sarah Safwat Moawad Amin

M.B.B.Ch – Ain Shams University – 2007;

Msc. Ain Shams University 2012

Assistant Lecturer of Obstetrics and Gynecology

Under Supervision of

Prof. Dr. Sherif Abdel Khalek Akl

Professor of Obstetrics and Gynecology

Faculty of Medicine – Ain Shams University

Prof. Dr. Hassan Tawfik Khairy

Professor of Obstetrics and Gynecology

Faculty of Medicine – Ain Shams University

Prof. Dr. Sherif Fekry Hendawy

Professor of Obstetrics and Gynecology

Faculty of Medicine – Ain Shams University

Dr. Sherif Hanafi Hussain

Assistant Professor of Obstetrics and Gynecology

Faculty of Medicine – Ain Shams University

**Faculty of Medicine
Ain Shams University
2015**



Acknowledgments

First and foremost, I feel always indebted to **Allah**, the Most Merciful, Who gives me power to accomplish this work,

I would like to express my deepest appreciation and sincere gratitude to **Prof. Dr. Sherif Abdel Khalek Akl**, Professor of Obstetrics and Gynecology, Faculty of Medicine – Ain Shams University, for his sincere help, constant encouragement, constructive criticism, and valuable guidance; I was truly honored to work under his supervision.

I wish also to express my great gratitude and utmost appreciation to **Prof. Dr. Hassan Tawfik Khairy**, Professor of Obstetrics and Gynecology, Faculty of Medicine – Ain Shams University, for his valuable suggestions and instructions during the progress of this work.

Special thanks go to **Prof. Dr. Sherif Fekry Hendawy**, Professor of Obstetrics and Gynecology, Faculty of Medicine – Ain Shams University, for his guidance, continuous support and encouragement through the whole work,

Special thanks go to **Dr. Sherif Hanafi Hussain**, Assistant Professor of Obstetrics and Gynecology, Faculty of Medicine – Ain Shams University, for his great efforts and time she has devoted to accomplish this work,

I owe special thanks to **Family**, especially my Parents for pushing me forward in every step in the journey of my life.

Special thanks to my husband dr.Mohammed Mahmoud Samy for his contribution in collecting & revising data.

 Sarah Safwat Moawad Amin



Contents

Subject	Page No.
List of Abbreviations	i
List of Tables.....	iii
List of Figures.....	v
Protocol.....	--
Introduction.....	1
Aim of the Work.....	4
Review of Literature	
- Placenta Accreta.....	5
- Management of Placenta Accreta	18
- Role of Imaging in Placenta Accreta	33
- Creatine kinase in placenta accreta.....	46
Patients and Methods.....	50
Results.....	70
Discussion	97
Summary.....	111
Conclusions and Recommendations	117
References.....	120
Arabic Summary	—

List of Abbreviations

Abbr.	Full term
2DUS	: Two-dimensional ultrasound
3D	: Three-dimensional
3DPD	: 3 dimensional power doppler
3DUS	: Three-dimensional ultrasound
ACOG	: American College of Obstetricians and Gynecologist
AJOG	: American Journal of Obstetrics and Gynecology
AUC	: Area under the curve
CK	: Creatine kinase
CS	: Cesarean section
EGFR	: Epidermal growth factor receptor
FFP	: Fresh frozen plasma
IIA	: Internal iliac artery
MAP	: Morbidly adherent placenta
MNGCs	: Number of multinucleated giant cells magnetic
MRI	: Magnetic resonance imaging
MS US	: Multislice ultrasound
MSV	: Multi-Slice View
NPV	: Negative predictive value
OMD	: Orthogonal display mode
PAD	: Placental adhesion disorders
PPV	: Positive predictive value
PRBCs	: Packed red blood cells.

List of Abbreviations *(Cont.)*

Abbr.	Full term
RCOG	: Royal College of Obstetricians and Gynecologist
ROC	: Receiver operating characteristic (ROC)
ROI	: Region of interest
SD	: Standard deviation
SPSS	: Statistical package for social sciences
UA	: Uterine artery
US	: Ultrasound
VEGFR-2	: Vascular endothelial growth factor receptor-2

List of Tables

Table No.	Title	Page No.
Table (1):	Characteristics of the whole study population	71
Table (2):	Characteristics of patients with normal placenta and those with abnormal placentation	72
Table (3):	Results of 2D grayscale US with 3DPD and 3D power Doppler with MS US examination in patients with normal placenta and those with abnormal placentation	75
Table (4):	Results of quantitative and qualitative CK assay in patients with normal placenta and those with abnormal placentation.....	77
Table (5):	Operative details in patients with normal placenta and those with abnormal placentation	79
Table (6):	Diagnostic value of 2D grayscale US with 3DPD for identification of abnormal placentation	82
Table (7):	Diagnostic value of 3D power Doppler with MS US for identification of abnormal placentation	84
Table (8):	Diagnostic value of elevated (>160 U/l) CK for identification of abnormal placentation	86
Table (9):	Receiver-operating characteristic (ROC) curve analysis for the value of quantitative CK assay in the diagnosis of abnormal placentation	88

List of Tables *(Cont.)*

Table No.	Title	Page No.
Table (10):	Diagnostic value of a CK cut-off level of >60 U/l for identification of abnormal placentation	90
Table (11):	Comparison of the sensitivities* of CK >60 U/l, 2D grayscale US, 3DPD and 3D power Doppler with MS US.....	92
Table (12):	Comparison of the specificities* of CK >60 U/l, 2D grayscale US, 3DPD and 3D power Doppler with MS US.....	93
Table (13):	Multivariable binary logistic regression analysis for the diagnostic value of 2D grayscale USwith 3DPD combined with quantitative CK assay	94
Table(14):	Comparison of the ROC curves associated with quantitative CK assay, 2DUSwith 3DPD, or Quantitative CK assay and 2DUS with 3DPD combined.....	95

List of Figures

Figure No.	Title	Page No.
Figure (1):	Photomicrograph of the utero-placental interface of a placenta accreta at term.	13
Figure (2):	Early pregnancy; sagittal view and low implanted sac in a pregnancy which went to term.	16
Figure (3):	Pregnancy complicated with placenta percreta.....	17
Figure (4):	Perioperative view of a placenta percreta left in situ after the closing of the vertical fundal uterine incision	22
Figure (5):	Placenta percreta with bladder invasion at cesarean delivery	32
Figure (6):	Gray-scale and color Doppler images showing sonographic diagnostic criteria for morbidly adherent placenta	35
Figure (7):	Three-dimensional power Doppler used to map intraplacental vascularization and vascularization of uterine serosa–bladder interface.....	37
Figure (8):	MRI: Abnormal uterine bulging	39
Figure (9):	MRI: heterogeneity of signal intensity within the placenta.....	40
Figure (10):	MRI: presence of dark intraplacental bands on T2-weighted images	40
Figure (11):	General electric Voluson E6.....	54

List of Figures (Cont.)

Figure No.	Title	Page No.
Figure (12):	2D grayscale showing focal exophytic mass in case of percreta	56
Figure (13):	2 D gray scale showing multiple abnormal placental lacunae with loss and disruption of the retroplacenta l sonolucent zone.....	56
Figure (14):	Placental invasion.....	59
Figure (15):	Normal placentation	61
Figure (16):	3D multislice showing normal placentation.....	62
Figure (17):	Abnormal placentation	63
Figure (18):	Mean age of patients with normal placenta and those with abnormal placentation.	73
Figure (19):	Parity in patients with normal placenta and those with abnormal placentation.	73
Figure (20):	Number of previous cesarean deliveries in patients with normal placenta and those with abnormal placentation.....	74
Figure (21):	Percentage of patients with evidence of placental invasion by 2D grayscale US or 3D power Doppler with MS US examination and those with elevated	76
Figure (22):	Box plot showing the CK level in patients with normal placenta and those with abnormal placentation.....	78

List of Figures (Cont.)

Figure No.	Title	Page No.
Figure (23):	Mean gestational age at delivery in patients with normal placenta and those with abnormal placentation.....	80
Figure (24):	Rate of performance of additional surgical procedures in patients with normal placenta and those with abnormal placentation.	81
Figure (25):	Incidence of unwanted intraoperative outcomes in patients with normal placenta and those with abnormal placentation.	81
Figure (26):	Performance of 2D grayscale US with 3DPD for identification of abnormal placentation.	83
Figure (27):	Performance of 3D power Doppler with MS US for identification of abnormal placentation.	85
Figure (28):	Performance of elevated (>160 U/l) CK for identification of abnormal placentation.	87
Figure (29):	Receiver-operating characteristic (ROC) curve for the value of quantitative CK assay in the diagnosis of abnormal placentation.	89
Figure (30):	Performance of a CK cut-off level of >60 U/l for identification of abnormal placentation.	91

List of Figures *(Cont.)*

Figure No.	Title	Page No.
Figure(31):	Comparison of the ROC curves associated with quantitative CK assay, 2DUS with 3DPD , or quantitative CK and 2DUS with 3DPD combined.....	96

Creatine Kinase as a Biochemical Marker in Diagnosis of Placenta Increta and Percreta

Protocol of Thesis

Submitted as Partial Fulfillment of MD Degree
in Obstetrics and Gynecology

By

Sarah Safwat Moawad Amin

M.B.B.Ch – Ain Shams University – 2007;
Msc. Ain Shams University 2012
Assistant Lecturer of Obstetrics and Gynecology

Under Supervision of

Prof. Dr. Sherif Abdel Khalek Akl

Professor of Obstetrics and Gynecology
Faculty of Medicine – Ain Shams University

Prof. Dr. Hassan Tawfik Khairy

Professor of Obstetrics and Gynecology
Faculty of Medicine – Ain Shams University

Prof. Dr. Sherif Fekry Hendawy

Professor of Obstetrics and Gynecology
Faculty of Medicine – Ain Shams University

Dr. Sherif Hanafi Hussain

Lecturer of Obstetrics and Gynecology
Faculty of Medicine – Ain Shams University

**Faculty of Medicine
Ain Shams University
2013**

Introduction