



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ





شبكة المعلومات الجامعية



شبكة المعلومات الجامعية

التوثيق الالكتروني والميكرو فيلم

جامعة عين شمس

التوثيق الالكتروني والميكرو فيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأفلام قد اعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15 – 20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of
15 – 25c and relative humidity 20-40 %



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



شبكة المعلومات الجامعية



بالرسالة صفحات
لم ترد بالأصل

Instrumental Methods for Determination of Some Antihistaminics

THESIS

**Presented by
Hanaa Karam Y. Mikael
(B. Pharmaceutical Sciences, 1988)
Cairo University**

**Presented for the Partial Fulfillment of
Master Degree
In
Pharmaceutical Sciences
(Analytical Chemistry)**

Supervized by

Prof.Dr. MOUSTAFA S. TAWAKKOL
Vice President and Dean of Faculty of Pharmacy
Misr University for Science and Technology

Prof. Dr. NAGLAA M. ELKOUSY
Professor Dr. of Pharmaceutical Chemistry
National Organization for Drug
Control and Research

Prof. Dr. AZZA A. M. MOUSTAFA
Professor Dr. of Analytical Chemistry
Faculty of Pharmacy Cairo
University

2001


B

9105


Approval Sheet

This thesis has been approved by the committee in charge :


1. Prof. Dr. M. Abd El Hady.
Professor of Analytical Chemistry,
Dean of Faculty of Pharmacy,
Alexandria University.




2. Prof. Dr. L. E. Abd El – Fattah.
Professor of Analytical Chemistry,
Faculty of Pharmacy,
Cairo University.



3. Prof. Dr. M. S. Towakkol
Professor of Analytical Chemistry,
Vice President and Dean of
Faculty of Pharmacy,
Misr University for Science and Technology. (supervisor)



4. Prof. Dr. A. A. M. Moustafa
Professor of Analytical Chemistry,
Faculty of Pharmacy,
Cairo University. (supervisor)



Note

Beside the work carried in this thesis, the candidate has attended and successfully passed the postgraduate courses examination with the grades mentioned in the following:

Subject	Grade
- Quality Control	Excellent
- Stability Indicating Methods	Very Good
- Bioavailability	Excellent
- Instrumental Analysis	Very Good
- Computer and its Application	Very Good
- Searching in Literature	Very Good
- Statistics	Excellent
- Higher Mathematics	Very Good

Prof. Dr. B. E. El-Zeany

Prof. and Head of Analytical Chemistry Department

Faculty of Pharmacy

Cairo University

ACKNOWLEDGEMENT

I am deeply grateful to Prof. Dr M. S. Tawakkol, Professor of Analytical Chemistry, Vice President and Dean of Faculty of Pharmacy Misr University for Science and Technology (MUST), for his kind supervision, valuable advices, continuous encouragement and guidance, as well as for his constructive comments and kind help during all stages of this thesis.

I would like to convey my deep appreciation to Prof. Dr. N. El Kousy, Professor of Pharmaceutical Chemistry, NODCAR, for her sincere help, beneficial discussions and fruitful criticism.

I am also indebted to Prof. Dr. A.A. Moustafa Professor of Analytical Chemistry, Faculty of Pharmacy, Cairo University, for her kind guidance and sincere help.

Thanks to Professor Dr. A. M. Moloukia, chairman of the National Organization for Drug Control and Research for his encouragement and for all colleagues and assistants in the National Organization for Drug Control and Research, for their kind help.

I am greatly grateful to my parents and husband for their help, support, patience, assistance and encouragement.

Hanaa Karam

CONTENTS

-The Aim of this Work.....	xviii
-Preface.....	xix

PART I: General Introduction and Literature Review

I.1	General Introduction.....	1
I.1.1	Definition, History and Classification	1
I.2	Literature Review.....	14
I.2.1	Acrivastine.....	14
I.2.2	Astemizole.....	16
I.2.3	Terfinadine.....	19

PART II: Spectrophotometric and Spectrofluorimetric Methods for the Determination of Acrivastine, Astemizole and Terfinadine Using Acid

Dye Technique:

II.1	Introduction.....	25
II.2	Experimental.....	26
II.2.1	Apparatus.....	26
II.2.2	Materials.....	26
II.2.2.1	Reagents.....	26
II.2.2.2	Pure Samples and Pharmaceutical Preparations.....	26
II.2.3	Standard Solutions.....	27
II.2.4	Test Solutions for Pharmaceutical Preparations.....	28
II.2.5	Colorimetric Methods Using Eosin B and TOO.....	28
II.2.5.1	Procedure.....	28
II.2.5.2	Optimizaion of Reactions Conditions.....	29
II.2.5.2.1	Determination of Wavelength of Maximum	

	Absorption of the Formed Ion Pair.....	29
II.2.5.2.2	Effect of pH and Volume of Buffer on the Reaction...	32
II.2.5.2.3	Effect of Extracting Solvent.....	32
II.2.5.2.4	Effect of Volume of Dye Solution.....	32
II.2.5.2.5	Effect of Phase Volume Ratio.....	36
II.2.5.2.6	Effect of Time and Number of Extractions.....	37
II.2.5.2.7	Stability of the Color	37
II.2.5.2.8	Assessment of the Stoichiometry of the Reactions.....	37
II.2.5.2.9	Linearity of Absorbance to Concentration of Drugs and Reproducibility of the Results.....	43
II.2.5.2.10	Application of the Proposed Procedures for the Determination of the Investigated Drugs in their Dosage Forms.....	50
II.2.6.	Spectrofluorimetric Method Using Eosin B.....	55
II.2.6.1	Determination of Excitation and Emission Maxima for the Reaction Products of Antihistaminic Drugs with Eosin B.....	55
II.2.6.2.	Procedures.....	55
II.2.6.2.1	Linearity of Fluorescence Intensity to Concentration of Drugs and Reproducibility of the Result.....	55
II.2.6.2.2	Application of the Proposed Spectrofluorimetric Procedure for the Determination of the Investigated Drugs in their Dosage Forms.....	61
II.2.7	Statistical Analysis of the Results.....	66
II.2.8.	Discussion.....	69

**PART III: Spectrophotometric Methods for the Determination of
Acrivastine and Astemizole Using Charge Transfer Technique Using
Iodine, DDQ and TCNE:**

III.1	Introduction.....	72
III.2	Experimental.....	73
III.2.1	Apparatus.....	73
III.2.2	Materials.....	74
III.2.2.1	Reagents	74
III.2.2.2	Pure Samples and Pharmaceutical Preparations.....	74
III.2.3	Standard Solutions	74
III.2.4	Test Solutions for Pharmaceutical Preparations.....	74
III.2.5	Procedures.....	75
III.2.6.	Optimization of Reaction Conditions.....	76
III.2.6.1	Determination of Wavelength of Maximum Absorption.....	76
III.2.6.2	Effect of Solvents.....	78
III.2.6.3	Effect of Volume of Reagent Solution.....	78
III.2.6.4	Effect of Time and Heating Temperature.....	78
III.2.6.5	Assessment of the Stoichiometry of the Reaction	82
III.2.6.6	Linearity of Absorbance to Concentration of Drugs and Reproducibility of the Results.....	82
III.2.6.7	Application of the Proposed Procedures for the Determination of the Investigated Drugs in their Pharmaceutical Dosage Forms.....	95
III.2.7	Statistical Analysis of the Results.....	102
III.2.8.	Discussion.....	105

PART IV: TLC and HPLC Methods for the Determination and Stability

Testing of Acrivastine:

IV.1	Introduction	108
	<u>Section A: TLC Method for the Determination and</u>	
	<u>Stability Testing of Acrivastine</u>	
IV.A.1	Experimental.....	110
IV.A.1.1	Apparatus.....	110
IV.A.1.2	Chemicals and Solvents.....	110
IV.A.1.3	Pure Samples and Pharmaceutical Preparations.....	110
IV.A.1.4.	Standard Solutions.....	111
IV.A.1.5.	Test Solutions for Pharmaceutical Preparations.....	111
IV.A.1.6.	Preparation of the Degradation Products.....	111
IV.A.1.7.	Preparation of the Laboratory Prepared Mixtures.....	112
IV.A.2.	Chromatography.....	112
IV.A.2.1	Optimization of the Separation.....	112
IV.A.2.2	Procedures for Construction of Calibration Curve and Reproducibility of the Results	114
IV.A.2.3.	Application of the Proposed Procedures for the Determination of Acrivastine in Semprex Capsules.....	115
IV.A.2.4.	Determination of Acrivastine in Presence of its Degradation Products.....	115
IV.A.3.	Statistical Analysis of the Results.....	121
IV.A.4.	Discussion.....	122
	<u>Section B: HPLC Method for the Determination and</u>	
	<u>Stability Testing of Acrivastine</u>	
IV.B.1	Experimental.....	123
IV.B.1.1	Apparatus.....	123
IV.B.1.2	Chemicals and Solvents.....	123

IV.B.1.3	Pure Samples and Pharmaceutical Preparations	123
IV.B.1.4.	Standard Solutions.....	124
IV.B.1.5.	Test Solutions for Pharmaceutical Preparations.....	124
IV.B.1.6.	Preparation of Laboratory Prepared Mixtures.....	124
IV.B.2.	Chromatography.....	124
IV.B.2.1	Optimization of the Separation.....	124
IV.B.2.2	Procedures for Construction of Calibration Curve and Reproducibility of the Results	125
IV.B.2.3.	Application of the Proposed Procedures for the Determination of Acrivastine in Semprex Capsules.....	125
IV.B.2.4.	Determination of Acrivastine in Presence of its Degradation Products.....	126
IV.B.3.	Statistical Analysis of the Results.....	132
IV.B.4.	Discussion.....	133
<u>References</u>	135

List of Figures

Figure	Page
(1) Absorption Spectra of Eosin B Complexes in Chloroform with Acrivastine ($7 \mu\text{g.ml}^{-1}$), Astemizole ($3 \mu\text{g.ml}^{-1}$) and Terfinadine ($10 \mu\text{g.ml}^{-1}$)	30
(2) Absorption Spectra of TOO Complexes in Chloroform with Acrivastine ($3.5 \mu\text{g.ml}^{-1}$), Astemizole ($6 \mu\text{g.ml}^{-1}$) and Terfinadine ($2.5 \mu\text{g.ml}^{-1}$).	31
(3) Effect of pH on the Absorbance of the Reaction Product of the Drugs with Eosin B. Acrivastine ($10 \mu\text{g.ml}^{-1}$), Astemizole ($10 \mu\text{g.ml}^{-1}$) and Terfinadine ($20 \mu\text{g.ml}^{-1}$)	33
(4) Effect of pH on the Absorbance of the Reaction Product of the Drugs with TOO. Acrivastine ($10 \mu\text{g.ml}^{-1}$), Astemizole ($10 \mu\text{g.ml}^{-1}$) and Terfinadine ($20 \mu\text{g.ml}^{-1}$).	33
(5) Effect of Volume of McIlvaine's Buffer on the Reaction with EosinB.Acrivastine ($10 \mu\text{g.ml}^{-1}$), Astemizole ($10 \mu\text{g.ml}^{-1}$) and Terfinadine ($10 \mu\text{g.ml}^{-1}$).	34
(6) Effect of Volume of McIlvaine's Buffer on the Reaction with TOO. Acrivastine ($10 \mu\text{g.ml}^{-1}$), Astemizole ($10 \mu\text{g.ml}^{-1}$) and Terfinadine ($20 \mu\text{g.ml}^{-1}$).	34
(7) Effect of Volume of 0.5% w/v Eosin B on the Reaction. Acrivastine ($10 \mu\text{g.ml}^{-1}$), Astemizole ($10 \mu\text{g.ml}^{-1}$) and Terfinadine ($20 \mu\text{g.ml}^{-1}$).	35
(8) Effect of Volume of 0.1% w/v TOO on the Reaction.Acrivastine ($10 \mu\text{g.ml}^{-1}$), Astemizole ($10 \mu\text{g.ml}^{-1}$) and Terfinadine ($20 \mu\text{g.ml}^{-1}$).	35
(9) Effect of Time of Extraction in Eosin B Method. Acrivastine	