

**EFFECT OF SOME TREATMENTS ON GROWTH
AND YIELD OF WASHINGTON NAVEL ORANGE
UNDER SANDY SOIL**

By

WAFAA ABD EL – AZIZ MOHAMED SHEHATA

B.Sc. Agric. Sci. (Pomology), Fac. Agric., Cairo Univ., Egypt, 2001

M.Sc. Agric. Sci. (Pomology), Fac. Agric., Cairo Univ., Egypt, 2008

THESIS

**Submitted in Partial Fulfillment of the
Requirements for the Degree of**

DOCTOR OF PHILOSOPHY

In

**Agricultural Sciences
(Pomology)**

**Department of Pomology
Faculty of Agriculture
Cairo University
EGYPT**

2017

APPROVAL SHEET

EFFECT OF SOME TREATMENTS ON GROWTH AND YIELD OF WASHINGTON NAVEL ORANGE UNDER SANDY SOIL

Ph.D. Thesis

In

Agric. Sci. (pomology)

By

WAFAA ABD EL – AZIZ MOHAMED SHEHATA

B.Sc. Agric. Sci. (Pomology), Fac. Agric., Cairo Univ., Egypt, 2001

M.Sc. Agric. Sci. (Pomology), Fac. Agric., Cairo Univ., Egypt, 2008

APPROVAL COMMITTEE

DR. SALAMA EID SALEM SHER.....

Research Professor of Pomology, Agriculture Research Center

Dr. RAMZY GEORGE STI.....

Prof. of Agric Pomology, Fac. Agric., Cairo University

Dr. MOHAMED REDA BARAKAT.....

Prof. of Agric Pomology, Fac. Agric., Cairo University

Dr. ABEER TAHSEEN MOHSEN.....

Prof. of Agric Pomology, Fac. Agric., Cairo University

Date: / / 2017

SUPERVISION SHEET

EFFECT OF SOME TREATMENTS ON GROWTH AND YIELD OF WASHINGTON NAVEL ORANGE UNDER SANDY SOIL

Ph.D. Thesis

In

Agric. Sci. (pomology)

By

WAFAA ABD EL – AZIZ MOHAMED SHEHATA

B.Sc. Agric. Sci. (Pomology), Fac. Agric., Cairo Univ., Egypt, 2001

M.Sc. Agric. Sci. (Pomology), Fac. Agric., Cairo Univ., Egypt, 2008

SUPERVISION COMMITTEE

Dr. MOHAMED REDA BARAKAT

Professor of Pomology, Fac. Agric., Cairo University

Dr. ABEER TAHSEEN MOHSEN

Professor of Pomology, Fac. Agric., Cairo University

Dr. RAMADAN ABOU SERIE SAYED

Research Professor of Pomology, Agriculture Research Center

ACKNOWLEDGEMENT

First of all, I do thank Allah for the gifts he has given me.

*There are no sufficient words to thank **prof. Dr. Mohammed Reda Barakat** Professor of Pomology, Faculty of Agriculture, Cairo University for suggesting the point of the work, fruitful supervision, encouragement and sincere guidance through out the practical work and revision the manuscript of this thesis.*

*I wish to express my sincere thanks, deepest gratitude and appreciation to **prof. Dr. Abeer Tahssen mohsen** professor of Pomology, Faculty of Agriculture, Cairo University for her guidance through the course of study, supervision, continued assistance and revision the manuscript of this thesis.*

*I wish to express my greatest gratitude to **prof. Dr. Ramadan Abou Serie Sayed** research professor of Citrus Research Department., Horticulture Research Institute, ARC, Giza for his valuable advices, continuous support and his guidance throughout this work,*

*Grateful appreciation is also extended to my friends and all staff members of Citrus Research Department and **Dr. Salama Eid Salem** for his careful revision of the manuscript.*

Special deep appreciation is given to my father, mother, brother, sisters and my husband Ahmed El-Sayed for his continuous help, encouragement and giving me his whole hearted support.

Thank you all

Contents

INTRODUCTION.....	1
REVIEW OF LITERATUR	4
1. Morphological and physiological characteristics.....	5
a. Effect of urea on morphological and physiological characteristics.....	5
b. Effect of petruscine on morphological and physiological characteristics.....	5
c. Effect of Gibberellins (GA ₃) on morphological and physiological characteristics.....	7
d. Effect of algae on morphological and physiological characteristics.....	8
2. Flowering and fruitlets parameters.....	11
a. Effect of urea on flowering and fruitlets parameters.....	11
b. Effect of petruscine on flowering and fruitlets parameters.....	12
c. Effect of Gibberellins (GA ₃) on flowering and fruitlets parameters.....	13
d. Effect of algae on flowering and fruitlets parameters.....	14
3. Yield and physical characteristics.....	15
a. Effect of urea on yield and physical characteristics.....	15
b. Effect of petruscine on yield and physical characteristics.....	17
c. Effect Gibberellins (GA ₃) on yield and physical characteristics.....	18
d. Effect of algae on yield and physical characteristics.....	19
4. Fruit quality.....	21
a. Effect of urea on fruit quality.....	21
b. Effect of petruscine on fruit quality.....	22
c. Effect of Gibberellins (GA ₃) on fruit quality.....	22
d. Effect of algae on fruit quality.....	24
5. Chemical Parameters.....	26
a. Effect of urea on chemical parameters:.....	26
b. Effect of petruscine on chemical parameters.....	27
c. Effect of Gibberellins (GA ₃) on chemical parameters.....	28
d. Effect of algae on chemical parameters.....	29
MATERIALS AND METHODS.....	32
RESULTS AND DISCUSSION.....	41
1. Morphological and physiological parameters.....	41
a. Shoot length "cm".....	41
b. Shoot diameter "mm".....	43
c. Leaf area "cm ² ".....	45
d. Dry matter ratio (%).....	47

2. Flowering and fruitlets parameters	50
a. Leafy inflorescence "% "	50
b. Woody inflorescence "% "	53
c. Fruit retention "% "	55
3. Yield and Fruit quality	58
a. Yield "kg"	58
b. Fruit weight "gm"	60
c. Fruit size "cm ³ "	63
d. Fruit length "mm"	64
e. Fruit diameter "mm"	66
f. Fruit juice percentage (%)	68
g. Ascorbic acid (vitamin C) "mg/100 ml"	70
h. Total acidity percentage "% "	72
i. Total soluble solids (T.S.S. %)	74
j. T.S.S/ Acid ratio:	76
4. Chemical Parameters	80
a. Leaf nitrogen content "gm/100gm"	80
b. Leaf phosphorus content "gm/100gm"	82
c. Leaf potassium content "gm/100gm"	84
d. Total carbohydrates "gm/100gm dried weight"	86
e. Total free amino acids	88
f. Plant pigments	90
SUMMARY	97
REFERENCE	112
الملخص العربي	1

INTRODUCTION

Citrus (*Citrus* spp.) is one of the most important fruit crops in Egypt. It is the first popular fruit in Egypt and the most important fruit crops regarding both production and consumption because of the high nutritional value of their licenses and the length of displayed markets. (El- Kassas, 1984). Moreover, it occupies about 530415 feddans which represent 34.7% of total fruit cultivated area, the fruiting area of citrus occupies 440706 feddan produced 4402180 tons in Egypt during 2014 according to the yearly Bull. Agric. Economic and Statistics- Ministry of Agriculture and Land Reclamation of Egypt.

Washington naval orange considered as reported by many workers the first crop among the different citrus cultivars grown in Egypt. Leading for local market and especially export (El-Shobaky and Mohamed, 2000). So, the improvement of fruit yield and quality are very important objective to be achieved by different management practices in Egypt, especially in poor sandy soils. (Sayed *et al*, 2007)

Flowering induction and flower number are main factors for yield and fruit setting production in citrus crops. This event usually is achievable when source of growth stimulation sustains within the environment. Availability of essential minerals during morphological and physiological process can play an important role in growth and fruit setting in fruit trees.

Citrus yield can be closely related to production of total flowers in spring bloom even though most of the flowers do not set fruit (Chermahini *et al*, 2011). Applications of various nutrients applied to tree also affect fruit growth development and overall final quality.

Nitrogen could influence the number of flowers and fruit set (Albrigo, 1999). The pre-bloom urea spray increases the reserve N since it is readily accessible and metabolizable (Rabe, 1994).

Polyamines a beneficial sub-group of amines. Polyamines possess diverse biological functions and are formed during metabolic processes in living organisms. Polyamines in plants are involved in many steps of protein synthesis, embryogenesis, transcription of genes, cell division, organ development, fruit ripening, leaf senescence, tuber dormancy and stress minimization of plant organs. Polyamines, natural compounds involved in many plant growth and development processes, are ubiquitously present in plant cells (Valero et al., 2002). Putrescine concentration increases in plants in response to a number of stress factors including water stress, acid treatment, mineral deficiency, and osmotic shock or CO₂ treatment. (Kosson and Prang, 2005).

The application of plant growth regulator (PGR) can provide significant economic advantages to citrus growers when used in appropriate situations as these have proven effective in stimulating a number of desired responses such as increase in fruit size and delay in fruit maturity (Basharat *et al*, 2008). Fruit development is thought to be triggered by hormones as it is evident from the report by (Talon et al., 1990) that the endogenous gibberellin status of the developing citrus ovaries is the limiting factor for the initiation of fruit development.

Plant hormones play an integral role in controlling the growth, development, metabolism and morphogenesis of higher plants (Taiz and Zeiger, 1991). Auxins, gibberellins, cytokinins, ethylene and abscisic acid are well known plant hormones. However, growth

hormones differ considerably in their mode of actions (Najma and Aisha, 2006).

Gibberellins are the most powerful of the growth promoters because they increase internode spacing, induce and promote flowering in many plants, and modify the flower sex expression in some plants (Davies, 1995).

Seaweed extract is a new generation of natural organic fertilizers containing highly effective nutritious and increase yield and resistant ability of many crops. Recently, macroalgae found to be viable alternative to chemical fertilization due to its high level of organic matter, micro and macro elements, vitamins, fatty acids and rich in growth regulators. (Crouch and Staden, 2005).

The main objective of this study was to investigate the possibility of pre-bloom foliar application of urea and Putrescine and full-bloom exogenous application of GA3, green algae extract and red algae extract for improve, morphological characteristics, flowering and fruitlets parameters, fruit quality, nutritional status and increase total yield of Washington navel orange. In addition to determine the optimum time of urea, Putrescine and GA3, green and red algae applied as a foliar spray to improve marketable yield of Washington navel orange for competitiveness of citrus fruit products in local and international markets.