



شبكة المعلومات الجامعية

بسم الله الرحمن الرحيم



شبكة المعلومات الجامعية
@ ASUNET



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الالكتروني والميكرو فيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأفلام قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيدا عن الغبار

في درجة حرارة من ١٥-٢٥ مئوية ورطوبة نسبية من ٢٠-٤٠%

To be Kept away from Dust in Dry Cool place of
15-25- c and relative humidity 20-40%

بعض الوثائق الأصلية تالفة

بالرسالة صفحات لم ترد بالاصل

**THE VALUE OF TRIPHASIC SPIRAL CT
IN COMPARISON TO OTHER CONVENTIONAL
MODALITIES IN DIAGNOSIS OF
HEPATOCELLULAR CARCINOMA**

Thesis

Submitted for Partial Fulfillment of
Master Degree in Tropical Medicine

By

ABEER IBRAHIM SAYD

(M.B.B.Ch.)

Faculty of Medicine
Ain Shams University

Supervised By

DR. ALY ZAKI HELMY

Ass. Professor of Tropical Medicine
Faculty of Medicine
Ain Shams University

DR. HESHAM KHALIL DABBOUS

Ass. Professor of Tropical Medicine
Faculty of Medicine
Ain Shams University

DR. OSAMA MOHAMED HETTA

Lecturer of Radiodiagnosis
Faculty of Medicine
Ain Shams University

Faculty of Medicine
Ain Shams University

2002

ACKNOWLEDGEMENT

*First and foremost thanks to **ALLAH**, the most merciful.*

*I would like to express my deep gratitude to **Prof. Dr. Afaf Massoud** Head of Tropical Medicine department, Ain Shams University, for her continuous support.*

*I wish to express my deep appreciation and sincere gratitude to **Dr. Aly Zaki Helmy** Ass. Prof. Of Tropical Medicine, Ain Shams University, for his close supervision, valuable instructions, continuous help, patience and guidance.*

*My deepest gratitude to **Dr. Hesham Khalil Dabbous** Ass. Prof. of Tropical medicine, Ain Shams University, for his encouragement and revision of the work.*

*I wish to express my supreme gratitude to **Dr. Osama Mohamed Hetta** Lecturer of Radiodiagnosis, Ain Shams University, for his immense effort in the practical part of the work.*

Last but not least, I want to thank all my colleagues, friends and my patients without their help, this work could not have been completed.

CONTENTS

	Page
Introduction	1
Aim of the Work	3
Review of Literature	4
Benign Tumors of the Liver	5
Malignant Tumor of the Liver	31
Patients and Methods	112
Results	118
Discussion	152
Conclusions and Recommendations	171
Summary	172
References	175
Arabic Summary	--

LIST OF TABLES

No.	Title	Page
1	Age distribution in the two studied groups	128
2	Sex and special habits distribution in the two studied groups	128
3	Symptomatology of the two studied groups	129
4	Comparison of the bleeding tendency between the two studied groups	130
5	Data of the past history of the two studied groups	131
6	Manifestations of liver cell failure in the two studied groups	132
7	Comparison of the abdominal examination findings between the two studied groups	133
8	Distribution of hepatitis C and B among the two studied groups	134
9	Comparison between the two studied groups as regards the liver function tests, PT, CBC and ESR	135
10	Detection of ascites by spiral CT in the two studied groups	136
11	Detection of portal vein thrombosis by spiral CT in the two studied groups	136
12	Detection of enlarged abdominal lymph nodes by spiral CT in the two studied groups	136
13	Comparison between the ultrasound and spiral CT abilities in detection of hepatic focal lesions number in group I	137

No.	Title	Page
14	Comparison between the ultrasound and spiral CT abilities in detection of hepatic focal lesions number in group II	138
15	Comparison between the ultrasound and spiral CT abilities in detection of ascites in all studied patients	139
16	Comparison between the ultrasound and spiral CT abilities in detection of enlarged abdominal lymph nodes in all studied patients	139
17	Comparison between the ultrasound and spiral CT abilities in detection of portal vein thrombosis in all studied patients	139
18	Comparison between spiral CT enhancement pattern of hepatic focal lesion(s) suggestive of HCC and histopathological examination of hepatic focal lesion(s) biopsies in group I	140
19	Comparison between spiral CT enhancement pattern of hepatic focal lesion(s) suggestive of HCC and histopathological examination of hepatic focal lesion(s) biopsies in group II	141
20	Correlation between the hepatic focal lesion(s) size detected by spiral CT and the serum level of alpha-fetoprotein in all studied patients	142

LIST OF FIGURES

No.	Title	Page
1	Triphasic spiral CT during the arterial phase shows an ill defined heterogeneously hyperattenuating mass (against the faintly enhanced liver parenchyma) in the posterior segment of the right lobe invading the posterior branch of right portal vein and eliciting neoplastic thrombosis (a). Rapid wash out of the contrast material from the mass and the portal vein thrombus is observed during the porto-venous phase (b). The lesion is hypoattenuating in delayed phase (c).	144
2	Triphasic spiral CT during the arterial phase shows a well defined nearly rounded hyperattenuating focal lesion arising from the left hepatic lobe with opacified branch of left hepatic artery that is supplying it (a). Feeding out of the contrast material from the lesion during the porto-venous phase (b). This picture is diagnostic for HCC.	145
3	Heterogeneously hyperattenuating hepatic focal lesion arising from the posterior segment of right hepatic lobe is observed during the arterial phase of spiral CT with a small hyperattenuating satellite lies anterior to it (a). Rapid wash out of the contrast material from the two lesions is observed during the porto-venous phase (b).	146
4a	Triphasic spiral CT during the arterial phase shows a hyperattenuating mass rising from the caudate lobe and another hyperattenuating smaller mass arising from the posterior segment of the right hepatic lobe at a lower level.	147
4b	Rapid wash out of the contrast material from the two lesions is observed during the porto-venous phase that is diagnostic of HCC.	148
5	Hepatocellular carcinoma, trabecular type, grade I. Well-differentiated malignant hepatocytes arranged in trabeculae and lined by endothelial cells. H & E, X 200.	149
6	Hepatocellular carcinoma, trabecular and acinar type, grade II. Malignant hepatocytes are arranged in trabecular and pseudoglandular pattern and lined by endothelial cells. Some acini contain colloid-like material. H & E, X 200.	149

No.	Title	Page
7	Hepatocellular carcinoma, grade III. The nuclei of anaplastic cells are markedly hyperchromatic and these cells are pleomorphic. Multinucleated giant tumor cells are evident. H & E, X 200.	150
8	Hepatocellular carcinoma, solid type, grade III. The anaplastic hepatocytes are arranged in solid pattern. Multinucleated giant tumor cells are also seen. H & E, X 200.	150
9	Hepatocellular carcinoma, grade IV. Anaplastic bizarre shaped spindle cells are seen. This case is proved to be HCC by immunostain. H & E, X 200.	151