### Role of Mixed Venous Oxygen Saturation in I.C.U Patients

#### Essay

Submitted for Partial Fulfillment for Master Degree in **Intensive Care** 

### Presented By

Sherien Mohammed Ibrahim El Gazzar

(M.B.B.,Ch.)

Faculty of Medicine - Benha University

### Supervised By

#### Prof. Dr. Bahaa El Din Ewees Hassan

Professor of Anesthesia and Intensive Care Faculty of Medicine - Ain Shams University

#### **Dr. Hadeel Magdy Abdel Hameed**

Assistant Professor of Anesthesia and Intensive Care Faculty of Medicine - Ain Shams University

#### Dr. Mai Mohsen Abdel Aziz

Lecturer of Anesthesia and Intensive Care Faculty of Medicine - Ain Shams University

Faculty Of Medicine
Ain Shams University
2013



سورة البقرة الآية: ٣٢



First of all, thanks to ALLAH for helping and guiding me in accomplishing this work and for everything else I have.

I would like to express deep thanks and regards to my supervisor **Prof. Bahaa El Din Ewees Hassan** Professor of Anesthesia And Intensive Care, Faculty of Medicine, Ain Shams University; for his generously guidance of my work. He shows a high capability to deal logically with everything and pays attention to tiny details to accomplish this study in a good manner.

I would like to express my deepest regards and appreciation to **Dr. Hadeel**Magdy Abdel Hameed, Assistant Professor of Anesthesia And Intensive Care,

Faculty of Medicine, Ain Shams University; for her sincere guidance, continuous

encouragement, advice, offering every possible help in all occasions and for her

valuable notes in this study.

I would like to express my deepest regards and appreciation to **Dr. Mai**Mohsen Abdel Aziz, Lecturer of Anesthesia And Intensive Care, Faculty of Medicine, Ain Shams University; for her patience, support, great help in all occasions and for her valuable notes in this study.

I would also like to give all thanks and gratitude to My parents for their unlimited support and help they offered me throughout my life.

Last but not least I would like to express my gratitude and greetings to **Those who** inspired us by sacrificing their lives for their dignity.

Sherin El-gazzar



#### **Contents**

Title	Page
♦ List of Figures	I
♦ List of Tables	VI
♦ List of Abbreviations	VII
♦ Introduction	1
♦ Aim of the Work	3
♦ Chapter (1): The Physiology of Oxyg Transport	
♦ Chapter (2): Monitoring of Mixed Ve Oxygen Saturation	
♦ Chapter (3): Clinical Applications of Mixed Venous Oxygen Saturation	
• Summary	120
♦ References	123
♦ Arabic Summary	······

Figures No.	Title	Page No.
Figure (1)	The oxygen cascade. The effects of hypoventilation are shown as the grey line and the effects of a pathological shunt are shown as a dashed line (Richardson et al., 1995).	5
Figure (2)	Graph to show perfusion and ventilation in different segments moving up the lung. Arrow indicates area of lung with perfect matching of ventilation to perfusion (West and Wagner, 1998).	8
Figure (3)	a): Normal diffusion from alveolus to capillary during passage of blood along the capillary. By 0.25s the red cell hemoglobin is completely saturated and the partial pressure of oxygen in the blood equilibrates with that in the alveolus and therefore diffusion stops. (b): Diffusion of oxygen with increased cardiac output (note the shorter timescale on the x-axis). The red cell may only be in contact with alveolar gas for 0.25s; however this will still allow sufficient time to achieve full saturation (West and Wagner, 1998).	9
Figure (4)	Oxyhemoglobin equilibrium (dissociation) curve of hemoglobin. The normal P50 value is indicated by the dashed lines. The changes in position of the curve associated with various effector molecules are indicated by the dashed arrows (Guyton and hall, 2010).	13
Figure (5)	Specifies the normal partial pressures and contents (or concentrations) for oxygen. It also describes the overall transfer of gas between the lungs and blood and between the blood and tissue (Pittman, 2011).	17
Figure (6)	Gas transport in a healthy 70-kg adult breathing room air (FIO2 of 0.21) at standard barometric pressure (10 kPa) (Archie, 1981).	19
Figure (7)	Clinical measures of pulmonary gas exchange, oxygen delivery, and oxygen consumption (Taylor and Weibel, 1981).	22

Figures No.	Title	Page No.
Figure (8)	Graph showing the balance between oxygen delivery and oxygen consumption. The horizontal solid line shows the extent to which oxygen delivery can be reduced and compensated by an increase in oxygen extraction (normally between 20-30%, between A and B). Point C shows the point beyond which compensation is insufficient and oxygen consumption is limited by delivery (supply-dependent), and anaerobic metabolism, producing lactic acid, results (Law and Bukwirwa, 1999).	38
Figure (9)	Schematic representation of the four types of acute circulatory failure. Importantly, several types of shock may coexist (Lim et al, 2003).	39
Figure (10)	Pulmonary artery catheter (Steingrub et al., 1991).	44
Figure (11)	Features of the pulmonary artery catheter (Karahan et al., 2008).	44
Figure (12)	Features of the pulmonary artery catheter (Cruz and Franklin, 2001).	45
Figure (13)	Reflection spectrophotometry (Reinhart et al., 2004).	47
Figure (14)	Technique of insertion (Kern, 2011).	49
Figure (15)	Choice of the site of insertion (Summerhill and Baram, 2005).	52
Figure (16)	Shows the proper orientation of the pulmonary artery catheter when inserted through the left subclavian vein, the curvature of the catheter is oriented so that it will facilitate passage of the catheter through the cardiac chambers and into the pulmonary artery(Lefrant et al., 2000)	56
Figure (17)	Modified Seldinger technique for central venous access (Rudy, 1993).	57

Figures No.	Title	Page No.
Figure (18)	Shows the appearance of a pulmonary artery (Swan-Ganz) catheter tip with the balloon at full inflation (top) and at less than full inflation (bottom), the tip of the catheter protrudes beyond the balloon when the balloon is less than fully inflated (Johnston et al., 1986).	60
Figure (19)	Waveforms by location of the pulmonary artery (PA) catheter tip, tracings obtained in the right atrium or pulmonary capillary wedge position share similar morphology, the transition from the right ventricle to the pulmonary artery tracing can be identified by the increase in diastolic pressure and the presence of a dicrotic notch, the diastolic "step-up" results from the transducer crossing the pulmonic valve, the dicrotic notch reflects closing of the pulmonic valve. (Marino, 1991).	62
Figure (20)	Shows the different components of the right atrial pressure tracing, a simultaneous ECG is shown to demonstrate the timing of the different components (Mark, 1998b).	63
Figure (21)	Trans-bronchial monitoring of (SvO2) (Brimacombe et al., 2000).	73
Figure (22)	Minimally invasive algorithm breaking down components of oxygen delivery and consumption followed by sub-components investigating root cause of imbalance (Zaja, 2007).	92
Figure (23)	SvO2 /hemoglobin (Hb) relation for VO2 at 200 ml O/min. The variations in cardiac output (Q) are accompanied by large variations in SvO2 (Sadaka et al., 2011).	99
Figure (24)	SvO2 recording after surgery, the effects and the treatment of shivering and the relationship between SvO2, cardiac output (CO) and metabolic rate (V $\times$ O2) (Richard et al., 2011).	100

Figures No.	Title	Page No.
Figure (25)	Effect of changing frequency of counter-pulsation from1:1(counter-pulsation with each heart beat) to 1:2(counter-pulsation with every other heart beat) on SvO2 (Gore and Sloan, 1984).	107
Figure (26)	Abrupt change in pulmonary arterial SvO2 in dobutamine-dependent patients whose intravenous line accidently became disconnected and response to reinstituting treatment (Gore and Sloan, 1984).	109
Figure (27)	Sodium nitroprusside infusion was started at the time marker. SvO2 and cardiac output rose as blood concentration increased (Divertie and McMichlan, 1984).	110
Figure (28)	Falling levels of SvO2, with increased PEEP. "Best PEEP" was 5cm $H2O^{=}\pi r^{2}O$ . Note influence of raised intrathoracic pressure with cough (Divertie and McMichan, 1984).	111
Figure (29)	SvO2/cardiac index (CI) relation according to the equation of Fick. The SvO2/CI is curvilinear. Thus, for constant VO2, the variations in CI induce large variations in SvO2if the initial CI value is low. By contrast, if the initial CI value is high, the variations have only slight impact on SvO2, these relations are altered if the variations in CI are associated with variations in oxygen consumption (VO2) (Jubran et al., 1998).	112
Figure (30)	A small and then a larger drop in SvO2, accompanied turning and tracheobronchial suctioning respectively. Level remained depressed and preceded cardiac arrest by 15 to 20 minutes (Divertie et al., 1983).	113

Figures No.	Title	Page No.
Figure (31)	The contribution of SvO2 to arterial hypoxemia in the setting of significant shunt (or V/Q mismatch) is illustrated. The effect of a large change in SvO2 is shown to highlight the impact of SvO2. In this example, there will be a change in SaO2 equivalent to half the increase in SvO2 (Walley et al., 1988).	115
Figure (32)	Arterial and venous oxygen saturations in various vascular regions, arterial oxygen saturations are shown on the right, and venous ones on the left. The numbers shown are percentages (Reinhart et al., 1994).	118

### **List of Tables**

Tables No.	Title	Pages No.
Table (1)	Normal blood oxygen values (Hett and Jonas, 2004).	12
Table (2)	Equipment needed for catheter insertion and pressure monitoring (Lefrant et al., 2000).	50
Table (3)	Mixed venous oxygen saturation (SvO2) and oxygen metabolism (Van Der Hoven et al., 1997).	75
Table (4)	Some causes of SvO2 fluctuation (Maddirala and Khan, 2010).	76

a/A	Ratio PaO2/PAO2
A-a	Gradient PAO2 - PaO2Alveolar-arterial Partial Pressure Oxygen Gradient
ATP	Adenosine triphosphate
CaO2	Content of oxygen in arterial blood
CCO	Continuous cardiac output
CCo2	Oxygen concentration in maximally saturated pulmonary end-capillary blood
CCU	Coronary care unit
CI	Cardiac index
cmH2O	Centimeters of water
CO	Cardiac output
CO2	Carbon dioxide;
СРАР	Continuous positive airway pressure
CvO2	Content of oxygen in venous blood
CVP	Central venous pressure
DO2	Oxygen delivery
DO2crit	Critical oxygen delivery
DO2I	Oxygen Delivery Index
2,3-DPG	Diphosphoglycerate
EGDT	Early Goal Directed Therapy
EMD	Electromechanical dissociation
ERO2	Oxygen extraction ratio
ERO2crit	Critical oxygen extraction ratio
FiO2	Fraction of inspired oxygen;
Hb	Hemoglobin

TII 02	011-1
HbO2	Oxyhemoglobin
ICU	Intensive care unit
Kpa	Kilopascal
$\mathbf{L}\mathbf{V}$	Left ventricle
LVEDP	Left ventricular end diastolic pressure
LVSWI	Left ventricular stroke work index
MAP	Mean arterial pressure
ml	Milliliter
mmHg	Millimeter of mercury
O2	Oxygen
OEC	Oxyhemoglobin dissociation curve
P50	At which 50% of the hemoglobin is saturated
PA	Pulmonary artery
PAC	Pulmonary artery catheter
PaCO2	Partial pressure of carbon dioxide in arterial blood
PAO2	Partial pressure of oxygen in alveoli
PaO2	Partial pressure of oxygen in arterial blood
PAP	Pulmonary artery pressure I.J.V.
PAWP	Pulmonary artery wedge pressure
PCO2	Partial pressure of carbon dioxide
PCWP	Pulmonary capillary wedge pressure
PeCO2	Mixed expired PCO2
PEEP	Positive end-expiratory pressure
PeO2	Mixed inspired PO2
PetCO2	End-tidal carbon dioxide

Pulmonary function tests
Hesymbol relating the hydrogen ion concentration or activity of a solution to that of a given standard solution.
Gastric pH
Inspired PO2
Partial pressure of oxygen (O2) in the blood
Partial pressure of oxygen in venous blood.
Pulmonary vascular resistance index
Intrapulmonary shunt
Shunt fraction
Cardiac Output
Right atrium
Red blood cells
Oxygen reserve
Return of spontaneous circulation
Respiratory Quotient
Right ventricle
Right ventricular end diastolic volume
Right ventricular ejection fraction
Right ventricular stroke work index
Arterial oxygen saturation
Oxygen saturation of central venous blood
Stroke volume index
Mixed venous oxygen saturation
Systemic vascular resistance index
Temperature

TB	Blood Temperature
TI	Injectate temperature
V/Q	Ventilation-perfusion
VIP	Variable infusion port
VO2	Oxygen consumption
vol%	The amount of O2 in milliliters that is in 100 mL of blood (vol% = mL O2/ 100 mL blood).



# Introduction

