

RECENT TRENDS IN LAPAROSCOPIC MANAGEMENT OF COLORECTAL CARCINOMA

Eassy

***Submitted for Partial Fulfillment of Master
Degree
In General Surgery***

By

Mina Fayez Abd-elsayed Youssef
*M.B.,B.Ch,
Faculty of Medicine, Tanta Univercity*

Under Supervision Of

Prof. Dr. Mohammed Emad Saleh
*Professor of general surgery
Faculty of medicine, Ain Shams university*

Dr. Amr Mohammed Mahmoud El-Hefny
*Lecturer of general surgery
Faculty of Medicine, Ain Shams University*

Dr. Fady Makram Benjamine
*Lecturer of general surgery
Faculty of Medicine, Ain Shams University*

Faculty of medicine

List of Contents

Title	Page No.
Abbreviations	ii
List of Tables	v
List of Figures	vi
Introduction	1
Aim of the work.....	5
<u>Review of Literature</u>	
Chapter (1): Surgical and laparoscopic anatomy of the colon and rectum	6
Chapter (2): Surgical pathology of colorectal carcinoma	39
Chapter (3): Diagnosis and investigations for colorectal carcinoma.....	52
Chapter (4): Conventional treatment of colorectal carcinoma.....	66
Chapter (5): Laparoscopic management of colorectal carcinoma.....	74
Chapter (6): Advantages, disadvantages and complications of laparoscopic management.....	172
Summary	192
Conclusions.....	194
References	195
Arabic Summary	--

List of Abbreviations

ACL	Anococcygeal ligament
ACRS	American Society of Colon and Rectal Surgeons
AJCC	The American Joint Committee on Cancer
APC	Adenomatous polyposis coli
APR	Abdominoperineal resection
APR	Abdominoperineal resection
AS	Anal sphincter
BL	Broad ligament
CD	Chron's disease
CDS	Complicated Diverticulosis of the Sigmoid
CEA	Carcinoembryonic antigen
COLOR	Colon Carcinoma Laparoscopic or Open Resection
DCBE	Double contrast barium enema
DRE	Digital rectal exam
EAES	European Association of Endoscopic Surgeons
EIA	External iliac artery
EIV	External iliac vein
FAP	Familial adenomatous polyposis
FOBT	Blood Fecal occult test
FT	Fallopian tube
GFN	Genitofemoral nerve
GV	Gonadal vessels
HALS	Hand-assisted Laparoscopic Surgery
HALS	Hand-assisted laparoscopic surgery
HF	hepatic flexure

List of Abbreviations_(Cont...)

HLA	Human leucocytes antigen
HN	Hypogastric nerve
HNPCC	Hereditary nonpolyposis colorectal cancer
ICA	Ileocolic artery
ICV	Ileocolic vein
IIA	Internal iliac artery
IIR	Internal inguinal ring
IMA	Inferior mesenteric artery
IMV	Inferior mesenteric vein
L bMCA	Left branch of the middle colic artery
LAC	Laparoscopic-assisted Colectomy
LCA	Left colic artery
LIQ	The Left lower Quadrant
LUL	Latral umbilical ligament
LUQ	The Left Upper Quadrant
MCA	Middle colic artery
MCV	Middle colic vein
MUL	Medial umbilical ligament
NHMRC	National Health and Medical Research Council
OC	Open Colectomy
Pb	Pancreatic body
PET	Positron emission tomography
PF	Pelvic floor
RA	Renal artery
RALC	Right-angled linear cutter

List of Abbreviations_(Cont...)

RIQ	The Right lower Quadrant
RUQ	The Right Upper Quadrant
SC	Sigmoid colon
SF	Splenic flexure
SLC	Straight linear cutter stapler
SMA	Superior mesenteric artery
SMV	Superior mesenteric vein
TC	Transverse colon
TME	Total Mesorectal Excision
UC	Ulcerative Colitis
UF	Uterine fundus

List of Tables

Table No.	Title	Page No.
Table (1)	Stages and prognosis for colorectal cancers.....	47

List of Figures

Figures No.	Title	Page No.
Fig. (1):	Blood supply of the colon	13
Fig. (2):	The portal venous circulation.....	17
Fig. (3):	Lymphatic Drainage of the colon	18
Fig. (4):	Lymphatic drainage of rectum and anal canal	20
Fig. (5):	Just below the liver in a thin patient, the hepatic flexure, duodenum, and pancreatic head may be seen. HF, hepatic flexure.....	22
Fig. (6):	The splenic flexure may be seen by lifting the omentum cephalad. In this thin patient, many of the left colonic vessels and retroperitoneal structures are seen. SF, splenic flexure; Pb, pancreatic body; L bMCA, left branch of the middle colic artery; RV, renal vein; RA, renal artery; K, kidney; LCA, left colic artery.....	24
Fig. (7):	Just inferior to the splenic flexure, the ligament of Treitz and the main vessels of the left colon are seen. LT, ligament of Treitz; IMV, inferior mesenteric vein; IMA, inferior mesenteric artery.....	24
Fig. (8):	By retracting the small bowel to the right side of the abdomen, the attachments of the sigmoid colon and the main vessels of the left colon may be seen	25
Fig. (9):	During the surgical mobilization of the sigmoid colon, the relationships of the gonadal vessels and the ureter are appreciated. SC, sigmoid colon; GV, gonadal vessels; U, ureter.....	25
Fig. (10):	During a surgical dissection of the origin of the inferior mesenteric artery, the relationships of the hypogastric nerves and the aorta are appreciated. Note how the two branches (left and right) are straddling the aorta. IMA, inferior mesenteric artery; A, aorta; HN, left branch of the hypogastric nerve plexus.....	26
Fig. (11):	With a patient in the Trendelenburg position and the right side tilted upward, the terminal ileum, cecum, and ligament of Treitz may all be visualized. I, terminal ileum; LT, ligament of Treitz; C, cecum.....	27
Fig. (12):	The major vascular structures of the right colon may be appreciated through the mesocolon, along with the right kidney and duodenum, with the small bowel retracted inferiorly and to the left. ICA, ileocolic artery; ICV, ileocolic vein; K, right kidney; D, duodenum; SMV, superior mesenteric vein; SMA, superior mesenteric artery.....	27
Fig. (13):	As the right colon is mobilized, the retroperitoneal structures are well seen. HF, hepatic flexure; Ps, psoas major muscle; GV, gonadal vessels; U, ureter; D, duodenum; P, pancreatic head; Ap, appendix.....	28
Fig. (14):	In thin patients, the vessels of the transverse colon and major structures in this region maybe seen. L, inferior edge of liver; TC, transverse colon; MCV, middle colic vein; MCA, middle colic artery; D, duodenum; P, head of pancreas	28

List of Figures (Cont...)

Figures No.	Title	Page No.
Fig. (15):	In the left inguinal region, the relationships of the gonadal vessels, vas deferens, and the major vessels exiting into the left leg are well appreciated during laparoscopy. GV, gonadal vessels; IIR, internal inguinal ring; EIA, external iliac artery; EIV, external iliac vein; Vd, vas deferens.....	29
Fig. (16):	A broad view of the pelvis is seen during laparoscopy in women. U, uterus; DP, Douglas pouch; O, left ovary; R, rectum; SC, sigmoid colon; U, right ureter	30
Fig. (17):	Lifting up on the right uterine adnexa permits appreciation of the relationships of these structures to the pelvis. UF, uterine fundus; O, ovary; FT, Fallopian tube; BL, broad ligament; EIA, external iliac artery	30
Fig. (18):	After complete mobilization of the rectum, the laparoscopic view affords excellent appreciation of some of the deep pelvic structures. R, rectum; AS, anal sphincter; PF, pelvic floor; ACL, anococcygeal ligament; C, coccyx	31
Fig. (19):	Umbilical ligament (transperitoneal view of right hemipelvis). MUL= medial umbilical ligament, MUL= medial umbilical ligament, LUL= lateral umbilical ligament, B= bladder, V = vas deferens, SPV= spermatic vessels, EPI= external iliac vein, U= ureter.....	33
Fig. (20):	Pouch of douglas, B = bladder, V = vas deferens, U = ureter, IIA = internal iliac artery, EIV = external iliac artery, R = rectum, Si =sigmoid colon.....	34
Fig. (21):	Base of vasculodeferential triangle composed of vas deferens (V) anteriorly, of medial umbilical ligament(MUL) medially, and of medial aspect of external iliac vein (EIV) laterally; triangle contains obturator lymphnodes along with obturator nerve and vessels (not visible). SpV = spermatic vessels, EIA = external iliac artery, GFN= genitofemoral nerve, B = bladder, U = ureter, IIA = internal iliac artery	35
Fig. (22):	Diagrammatic representation of two forms of sessile polyp (hyperplastic polyp and adenoma) and of two types of adenoma (pedunculated and sessile). There is only a loose association between the tubular architecture for pedunculated adenomas and the villous architecture for sessile adenomas.....	40
Fig. (23):	Polyps of colon as revealed by colonoscopy. A. Sessile, B. Pedunculated, C. Tubulovillous (<i>Chen et al. 2005</i>).	40
Fig. (24):	A, Pedunculated adenoma showing a fibrovascular stalk lined by normal colonic mucosa and a head that contains abundant dysplastic epithelial glands, hence the blue color with the H & E stain. B, A small focus of adenomatous epithelium in an otherwise normal (mucin-secreting, clear) colonic mucosa, showing how the dysplastic columnar epithelium (deeply stained) can populate a colonic crypt and create a tubular architecture.....	44

List of Figures (Cont...)

Figures No.	Title	Page No.
Fig. (25):	A- Familial adenomatous polyposis in an 18-year-old woman. The mucosal surface is carpeted by innumerable polypoid adenomas. B- Endoscopic image of sigmoid colon of patient with familial adenomatous polyposis	44
Fig. (26):	Carcinoma of the caecum. The fungating carcinoma projects into the lumen but has not caused obstruction.....	48
Fig. (27):	Carcinoma of the descending colon. This circumferential tumor has heaped-up edges and an ulcerated central portion. The arrows identify separate mucosal polyps.....	48
Fig. (28):	Positions of the surgical team and equipment for the diagnostic laparoscopy at the beginning of the procedure.....	59
Fig. (29):	Cannula positions for the diagnostic laparoscopy. If liver ultrasound is used, the right cannula should be a 10-mm size (asterisk)	60
Fig. (30):	Alaparoscopic ultrasound probe can be readily used to assess the liver and retroperitoneal structures during diagnostic laparoscopy.....	62
Fig. (31):	“Rtmning” of the small bowel begins with appropriate positioning of the patient (inset: patient in left side up with intestines retracted to the right) and starting the evaluation at the ligament of Treitz. Note that the inferior mesenteric vein is readily seen even in moderately obese patients.....	63
Fig. (32):	Running the bowel using the “hand-over-hand” technique: The right-handed grasper (1) releases the bowel and prepares to move from point A on the bowel to point C, whereas the left-handed grasper (2), at point B, prepares to slide underneath the other grasper	64
Fig. (33):	Diagnostic laparoscopy nearly always affords an excellent view of the uterus, Fallopian tubes, and ovaries	65
Fig. (34):	Extent of surgical resection is defined by the arterial-associated nodal distribution. The normal arterial anatomy is shown here. The superior hemorrhoidal(superior rectal) artery is the continuation of the inferior mesenteric artery.....	66
Fig. (35):	Extent of resection for a cecal or ascending colon cancer; the ileocolic, right colic, and right branch of the middle colic vessels are divided for adequate lymphadenectomy.	68
Fig. (36):	Cecal-sparing resection in selected patients with hepatic-flexure cancers.	68
Fig. (37):	Right and transverse colectomy for cancers in the mid-transverse colon or splenic flexure.	70
Fig. (38):	Left hemicolectomy resection for a proximal sigmoid or descending colon cancer. If the inferior mesenteric artery (with superior rectal) is included in the resection, the anastomosis must be to the upper rectum to allow adequate vascularization from pelvic branches.....	72

List of Figures (Cont...)

Figures No.	Title	Page No.
Fig. (39):	Positions of the equipment and the surgical team for the laparoscopic right colectomy. Initially the surgeon is at the left side of the patient, and returns to this position after lymphovascular pedicle ligation.....	75
Fig. (40):	Positions of the cannulae for the right colectomy. The suprapubic cannula is used for the laparoscope while the lymphovascular pedicles are dissected and divided.....	77
Fig. (41):	Good visualization of the right mesocolon is achieved by proper positioning of the patient and by placement of the omentum above the colon.(<i>Jacobs et al., 1991</i>).....	77
Fig. (42):	Definitive identification of the ileocolic pedicle is achieved by retraction at the ileocecal junction.....	78
Fig. (43):	Various approaches to the right colon mobilization have been described. A: Lateral to medial (“classic” open approach); B: Medial to lateral (preferred approach); C: Retroperitoneal approach	78
Fig. (44):	The surgeon’s first step in the dissection is to mark the inferior border of the ileocolic pedicle.....	80
Fig. (45):	From between the legs, the surgeon dissects the peritoneum overlying the ileocolic vascular pedicle over to the superior mesenteric vessels	80
Fig. (46):	The origins of the ileocolic artery and vein are identified, clipped, and divided.....	81
Fig. (47):	Anatomic variations of the origin of the ileocolic vessels. A The ileocolic artery runs in front of the superior mesenteric vein. B The ileocolic artery runs behind the superior mesenteric vein.....	81
Fig. (48):	Dissection of the ventral side of the superior mesenteric vein permits a complete dissection of the root of the middle colic artery and vein.....	82
Fig. (49):	Accessory middle colic or right colic veins are clipped and divided. These are common	83
Fig. (50):	The peritoneum is incised along the base of the ileal mesentery upward to the duodenum.....	84
Fig. (51):	The right mesocolon is dissected away from the retroperitoneal structures from medial to lateral	84
Fig. (52):	The venous anatomy between the hepatic flexure and the middle colic vessels has many variations, including Henle’s trunk, the fusion between the gastroepiploic vein, and a branch of the right or middle colic vein	86
Fig. (53):	With earlier steps accomplished, the hepatocolic ligament is easily divided, freeing up the proximal transverse and hepatic flexure of the right colon	86
Fig. (54):	Finally, the tumor-bearing segment of the right colon, with its lateral attachments, are freed up, completing the right colon mobilization	87

List of Figures (Cont...)

Figures No.	Title	Page No.
Fig. (55):	After drawing out the right colon using a wound protector, an anastomosis is accomplished extracorporeally. A A functional end-to-end anastomosis is created with a linear-cutter stapler. Note that the colon is occluded using a large Kocher clamp. B The anastomosis is completed with a rightangled firing of the linear-cutter stapler, completely sealing off the bowel. C The completed anastomosis before returning it to the abdomen.....	87
Fig. (56):	Positions of the equipment and the surgical team for the laparoscopic sigmoid colectomy.....	90
Fig. (57):	Positions of the cannulae for laparoscopic sigmoid colectomy	90
Fig. (58):	Active positioning using gravity produces optimum exposure. The greater omentum and the transverse colon are placed in the left subphrenic region and maintained in this position by the Trendelenburg tilt (inset). Subsequently, the proximal small bowel loops are placed in the right upper quadrant.....	92
Fig. (59):	The uterus can be suspended to the abdominal wall using a suture placed through its fundus	94
Fig. (60):	Initial dissection starts with an incision of the sigmoid mesentery at the sacral promontory with dissection cephalad posterior to the IMA.....	95
Fig. (61):	The dissection behind the IMA involves preservation of the main hypogastric nerve trunks, but also division of the branches traveling to the colon	95
Fig. (62):	Radical lymphadenectomy involves exposure of the main trunk of the IMA and skeletonization, but preservation of the hypogastric nerve trunks.....	96
Fig. (63):	The IMA is divided 1—2cm distal to its origin, or just distal to the left colic branch.....	97
Fig. (64):	The IMV is divided in a safe area between the pancreas and the left colic vessels	99
Fig. (65):	An avascular plane exists between Toldt's fascia and the mesocolon, which is bluntly dissected medial to lateral after IMA and IMV ligation. (Inset: Cross-sectional drawing illustrating the correct surgical plane indicated by arrow)	99
Fig. (66):	Lateral dissection then proceeds after the previous medial dissection	101
Fig. (67):	The dissection of the upper rectum should proceed with caution because the hypogastric nerves are tented upward and may be inadvertently injured. These nerves may be swept posteriorly before dividing the soft tissues in the area	102
Fig. (68):	After upper rectal mobilization, area of mesorectal division is chosen	103
Fig. (69):	Distal bowel division is performed through the right lower quadrant cannula using an endoscopic stapler	103

List of Figures (Cont...)

Figures No.	Title	Page No.
Fig. (70):	Proximal bowel division is performed after dividing the mesocolon up to the chosen site	104
Fig. (71):	Medial to lateral dissection beneath the left mesocolon provides excellent views of the distal pancreas, the base of the left transverse mesocolon, and retroperitoneum	106
Fig. (72):	Specimen extraction at the suprapubic site involves double protection: 1) a wound protector; and 2) an impermeable retrieval sac	107
Fig. (73):	After specimen extraction, the proximal colon is drawn out through this site, keeping the wound protector in place	109
Fig. (74):	The anvil and center rod of the circular stapler are introduced into the bowel lumen and secured with a purse string suture.....	109
Fig. (75):	The bowel is reintroduced into the abdominal cavity, checking for adequate length for anastomosis. The bowel should comfortably reach the pelvis without tension	110
Fig. (76):	Reestablishment of the pneumoperitoneum can be achieved quickly by twisting the wound protector, then clamping it at the skin level with a Kocher clamp(<i>Balli et al., 2000</i>).	110
Fig. (77):	The anastomosis is then done under laparoscopic guidance, perforating the proximal rectal stump with the sharp spike of the circular stapler, then performing a standard double-stapled anastomosis	111
Fig. (78):	After firing the stapler, the anastomosis is checked by filling the pelvis with saline, then insufflating the rectum with air using a rectoscope. The bowel upstream of the anastomosis is gently occluded during this test.....	111
Fig. (79):	Positions of the equipment and the surgical team for phase I of the laparoscopic total abdominal colectomy.....	116
Fig. (80):	Positions of the equipment and the surgical team for phase II of the laparoscopic total abdominal colectomy.....	116
Fig. (81):	Positions of the equipment and the surgical team for phase III of the laparoscopic total abdominal colectomy.....	117
Fig. (82):	Positions of the cannulae for laparoscopic total abdominal colectomy.....	118
Fig. (83):	Dissection is commenced at the sacral promontory posterior to the inferior mesenteric vessels.....	120
Fig. (84):	Dissection is continued superiorly beneath the IMA, protecting the hypogastric neural plexus.....	120
Fig. (85):	After creating a peritoneal window to the left of the inferior mesenteric vein, the pedicle is ligated using an endoscopic stapler. Note that the ureter is clearly isolated posterior to the pedicle.....	120
Fig. (86):	The left colic artery and vein are ligated separately	122
Fig. (87):	The left mesocolon is dissected away from the retroperitoneal structures using medial to lateral blunt dissection	122

List of Figures (Cont...)

Figures No.	Title	Page No.
Fig. (88):	The mesorectum is divided sharply, starting on the right side, using a harmonic scalpel or bipolar device	123
Fig. (89):	Therectum is next divided from the right side using one or two applications of an endoscopic stapler	123
Fig. (90):	The colon is reflected medially and dissection of the lateral attachments of the left colon proceed up to the splenic flexure.....	125
Fig. (91):	Separation of the omentum from the colon is required for accurate mobilization of the splenic flexure.....	126
Fig. (92):	Splenic flexure mobilization may be expedited with the dissection proceeding from medial to lateral on the transverse colon.....	126
Fig. (93):	Phase III begins with an incision just below the ileocolic pedicle.....	128
Fig. (94):	After mobilizing the pedicle, it is ligated well away from its origin using a bipolar coagulation device.....	128
Fig. (95):	The ileal and right colonic mesenteric attachments are completely freed retroperitoneally, bluntly dissecting a tunnel dorsal to the ileal mesentery.....	129
Fig. (96):	Just cephalad to the ligated ileocolic pedicle, the middle colic pedicle is identified.....	130
Fig. (97):	The ligation of the middle colic vessels may be safer if performed on the right and left branches, rather than on the main trunk	131
Fig. (98):	Attachments of the ileum just medial to the base of the appendix are incised, carrying the incision cephalad toward the root of the mesentery.....	133
Fig. (99):	The last lateral adhesions of the right colon are incised up to the hepatic flexure.....	133
Fig. (100):	The hepatocolic ligament is divided from medial to lateral, completely freeing up the right colon	134
Fig. (101):	The entire colon may then be pulled out through the suprapubic incision.....	134
Fig. (102):	After removing the entire colon, the center rod and anvil is inserted into the end of the ileum and secured using a purse-string suture	135
Fig. (103):	After passing the circular stapler up to the top of the rectum, the spike is protruded through the rectal wall adjacent to the rectal staple line.....	136
Fig. (104):	A standard double-stapled technique is used for the anastomosis of ileum to rectum.....	136
Fig. (105):	Positions of the equipment and the surgical team for the laparoscopic anterior resection for rectal cancer	139
Fig. (106):	Positions of the cannulae for the laparoscopic anterior resection. Note that the surgeon works mainly through the right lower quadrant and the suprapubic cannulae	140
Fig. (107):	Dissecting plane from the medial or lateral sides of the sigmoid colon optimally involve sweeping the Toldt's fascia posteriorly (thick gray line). This safely isolates and preserves the ureter and gonadal vessels	141

List of Figures (Cont...)

Figures No.	Title	Page No.
Fig. (108):	In the medial approach, the superior rectal (or inferior mesenteric) vessels are tented anteriorly and the plane is dissected between the vessels and Toldt's fascia.....	142
Fig. (109):	It is always an option to perform dissection laterally, verifying the location of the ureter and gonadal vessels.....	142
Fig. (110):	Once the adventitious tunica of the inferior mesenteric artery is exposed, it is clipped then transected. Hypogastric nerves are exposed and preserved.....	146
Fig. (111):	Next, the inferior mesenteric vein and the left colic artery can be simultaneously divided with an endoscopic stapler from the right side. Note that the ureter and gonadal vessels are clear of the stapler	147
Fig. (112):	The rectal dissection starts from the right side, carefully identifying and sweeping down the hypogastric nerves, which can be tented upward with traction	147
Fig. (113):	With careful traction and countertraction by the surgeon and the assistant, the boundary of the left side of the rectum between peritoneum and mesorectum is dissected (arrow)	149
Fig. (114):	Next, the peritoneal reflection is incised, exposing Denonvilliers' fascia and protecting the seminal vesicles or vaginal wall.....	149
Fig. (115):	The lateral ligaments are placed under tension by drawing the rectum to the right side of the pelvis, then this area is dissected, carefully preserving the nerve trunks heading distally	150
Fig. (116):	With tension applied to the left side of the rectum at the proposed transaction line, the mesorectum is divided using the laparoscopic coagulation shears.....	152
Fig. (117):	An endoscopic linear stapler is introduced through the suprapubic cannula and fired across the distal resection line at right angles to the bowel.	152
Fig. (118):	A transanally introduced circular stapler is placed at the top of the rectum and the spike is protruded through the wall posterior to linear staple line	153
Fig. (119):	Left. The double-stapled anastomosis is performed with all surrounding tissues clear of the two bowel ends.....	154
Fig. (120):	Right. After resection, the staples should be evaluated to be sure there has been good "B" formation of the staples [both from laparoscopic and intraluminal (endoscopic) evaluation]	154
Fig. (121):	Positions of the equipment and the surgical team for the laparoscopic APR.....	157
Fig. (122):	Positions of the cannulae for the laparoscopic APR. Note that the proposed colostomy site is <i>not</i> used as a cannula site, because it is usually too close to the (optical) port through which the laparoscope is used	158
Fig. (123):	Initial dissection involves incising the peritoneum just to the right of the superior rectal artery just at the sacral promontory and working cephalad.	160

List of Figures (Cont...)

Figures No.	Title	Page No.
Fig. (124):	The IMA is divided using a LigaSure 5mm device, dividing small visceral branches of the hypogastric nerves, but preserving the main trunks leading into the pelvis	160
Fig. (125):	Dissection is then continued <i>medial</i> to <i>lateral</i> beneath the divided IMA, identifying and sweeping the ureter and gonadal vessels posteriorly	161
Fig. (126):	When the ureter cannot be easily identified on the medial side, dissection should then proceed laterally, identifying and placing a cotton gauze over the ureter. Returning to the medial side, vessel ligation may proceed with the firm knowledge that the ureter is protected beneath the gauze	162
Fig. (127):	The IMV is ligated only if the ureter is identified and protected.....	163
Fig. (128):	Lateral attachments of the sigmoid colon are next incised sharply.....	164
Fig. (129):	Using triangulating tension, the sigmoid mesocolon is incised up to the bowel edge.....	164
Fig. (130):	Proximal resection line is next incised with an endoscopic stapler	164
Fig. (131):	Posterior mobilization is initiated next at the sacral promontory, carefully sweeping off the hypogastric nerve branches which may be tented upward in the line of dissection (arrows).....	165
Fig. (132):	As the posterior rectal mobilization proceeds, the hypogastric nerves may be well visualized and protected.....	166
Fig. (133):	At the level of the lateral stalks, the pelvic plexus can be preserved unless there is direct tumor infiltration	167
Fig. (134):	It may be highly useful to use the surgeon's doubly-gloved index finger to sound out the rectum or vagina in the distal rectal dissection, confirming the proper plane of dissection.....	168
Fig. (135):	Just as in the rectal mobilization, the perineal phase of the operation is most safely performed in a pattern of 1) posterior, 2) lateral, and 3) anterior dissection. Inset: Anatomic view of the coccyx posteriorly and the pelvic floor and anal sphincter muscles with the perineal dissection.....	170