



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ





شبكة المعلومات الجامعية



شبكة المعلومات الجامعية

التوثيق الالكتروني والميكرو فيلم

جامعة عين شمس

التوثيق الالكتروني والميكرو فيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأفلام قد اعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15 – 20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of
15 – 25c and relative humidity 20-40 %



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



شبكة المعلومات الجامعية



بالرسالة صفحات
لم ترد بالأصل

SEMINAL NEUTRAL ALPHA-1,4 GLUCOSIDASE ACTIVITY IN INFERTILE MALE PATIENTS

Thesis Presented

By

ADEL AWAD NAGUIB AWAD

(M.B.B.Ch)

As a partial Fulfilment of requirements of

Master Degree

In

Dermatology & Venereology

SUPERVISORS

Prof. Dr. MOSTAFA MOHAMMED SOLIMAN

Prof., and Chairman of Dermatology

& Venereology Department

Faculty of Medicine, Menoufia University

Prof. Dr. SAMIA HASSAN KANDIEL

Prof., and Chairman of Clinical

pathology Department

Faculty of Medicine, Menoufia University

Dr. MOHAMMED AHMED BASHA

Lecturer of

Dermatology & Venereology

Faculty of Medicine, Menoufia University

FACULTY OF MEDICINE
MENOUFIA UNIVERSITY

1994

B 9 C 11

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

To My Family

ACKNOWLEDGEMENT

Firstly and lastly thanks to God who helps me for accomplishment of this work .

Wording is not enough to express my sincere gratitude and deepest appreciation to prof . **Dr. MOSTAFA MOHAMMED SOLIMAN** , prof . and chairman of Department of Dermatology & Venereology .Faculty of Medicine , Menoufia University . His fatherly attitude , enthusiastic encouragement , and continuous support made the accomplishment of this work .

I would like to express may thanks and gratitude to prof . **Dr. SAMIA HASSAN KANDIEL** , Prof. and chairman of Clinical pathology Department Faculty of Medicine , Menoufia University . For continous help untiring devotion , and participation in preparing this work .

I wish to express my deep thanks and sincere gratitude to prof . **Dr . MOHAMMED AHMED BASHA** , Lecturer of Dermatology & Venereology faculty of Medicine , Menoufia University . For his unforgettable help , valuable advice , appreciable encouragement , and careful follow up of this work .

I wish to acknowledge the assistance of all members of the staff of Dermatology & Venerology , and Clinical pathology Departments in all phases of this work .

Lastly , but not the least , I feel deeply thankful to all those who extended to me a helping hand during preparation of this work .

Adel Awad Naguib

CONTENTS

	Page
Introduction and Aim of the work	1
Review of Literature :-	
- Seminal plasma	3
- Enzymes of seminal plasma	8
- Seminal Neutral α - 1,4 glucosidase enzyme	11
- Functions of the epididymis	16
- Hormonal control of the epididymal functions	24
- Sperm Density	26
- Sperm Motility	30
Material and Methods	37
- Results	47
- Discussion	91
- Summary and conclusion	99
- References	101
- Arabic summary	—

List of Tables

Table No.		Page
(1)	Age and seminogram of control group.	53
(2)	Comparison of the mean seminal parameters in the normal control group .	54
(3)	Age and seminogram of oligoathenozoospermic with varicocele .	55
(4)	Comparison of the mean seminal parameters in oligoathenozoospermic group with varicocele .	56
(5)	Age and seminogram of oligoathenozoospermic group without varicocele .	57
(6)	Comparison of the mean seminal parameters in oligoathenozoospermic group without varicocele .	58
(7)	Seminal neutral alpha - 1,4 glucosidase enzyme activity n - mol / ml in control , oligoathenozoosperic with varicocele , and oligoathenozoospermic without varicocele .	65
(8)	comparison of the mean seminal neutral alpha - 1,4 glucosidase enzyme activity in (control - oligoathenozoospermic with varicocele and oligoathenozoospermic with out varicocele) groups .	66
(9)	Comparison of the mean different seminal parameters in the normal control group and oligoathenozoospermic group with varicocele (Student T test) .	68

- (10) Comparison of the mean different seminal parameters in the normal control group and oligoathenozoospermic group without varicocele (student T test) . 69
- (11) Comparison of the mean different seminal parameters in the oligoathenozoospermic group with varicocele and oligoathenozoospermic group without varicocele (student T test) . 70
- (12) Comparison of the mean seminal neutral α - 1,4 glucosidase enzyme activity n.mol / ml in the normal control group and oligoathenozoospermic group with varicocele (student T test) . 71
- (13) Comparison of the mean seminal neutral α - 1,4 glucosidase activity n.mol / ml in the normal control group and oligoathenozoospermic group without varicocele (student T test) . 72
- (14) Comparison of the mean seminal neutral α - 1,4 glucosidase activity n.mol / ml in oligoathenozoospermic with varicocele and oligoathenozoospermic without varicocele (student T test) . 73
- (15) Correlation between seminal neutral α - 1,4 glucosidase activity (n.mol / ml) and sperm count in normal control group and different groups. 74

(16)	Correlation between seminal neutral α - 1,4 glucosidase activity (n. mol /ml) and sperm motility (%) in normal control group and different groups .	77
(17)	Correlation between seminal neutral α - 1,4 glucosidase activity (n.mol / ml) and abnormal forms (%) in normal control group and different groups .	80
(18)	Correlation between seminal neutral α - 1,4 glucosidase activity (n.mol / ml) and different seminal parameters in normal control group and different groups .	83
(19)	Correlation between different seminal parameters in normal control group and different groups .	87

List of Figures

Figure No.		Page
(1)	The mean volume of ejaculate (ml) in normal control subjects and different groups .	59
(2)	The mean liquification time (min) in normal control subjects and different groups .	60
(3)	The mean P.H. of semen in normal control subjects and different groups .	61
(4)	The mean sperm count (mill / ml) in normal control subjects and different groups .	62
(5)	The mean sperm motility (%) in normal control subjects and different groups .	63
(6)	The mean abnormal forms (%) in normal control subjects and different groups .	64
(7)	The mean seminal neutral α - 1,4 glucosidase activity (n.mol /ml) control subjects and different groups .	67
(8)	Correlation between seminal plasma neutral α - 1,4 glucosidase activity (n.mol / ml) and sperm count (mill /ml) (control * oligoathenozoospermic with varicocele [□])	75
(9)	Correlation between seminal plasma neutral α - 1,4 glucosidase activity (n.mol / ml) and sperm count (mill /ml) .(control * , oligoathenozoospermic without varicocele [⊗]) .	76

- (10) Correlation between seminal plasma neutral α -1,4 glucosidase activity (n.mol / ml) and sperm motility (%) after liquification . (control *, oligoathenozoospermic with varicocele [*]). 78
- (11) Correlation between seminal plasma neutral α - 1,4 glucosidase activity (n.mol / ml) and motility (%) after liquification . (control * , oligoathenozoospermic without varicocele [*]) . 79
- (12) Correlation between seminal plasma neutral α -1,4 glucosidase activity (n.mol / ml) and abnormal forms (%) . (control * , oligoathenozoospermic with varicocele [*]). 81
- (13) Correlation between seminal plasma neutral α - 1,4 glucosidase activity (n.mol / ml) and abnormal forms (%) .(control * , oligoathenozoospermic without varicocele [*]). 82
- (14) Correlation between seminal plasma neutral α - 1,4 glucosidase activity (n.mol / ml) and volume of ejaculate (ml). (control *,oligoathenozoospermic with varicocele [*] , oligoathenozoospermic without varicocele [*]) . 84
- (15) Correlation between seminal plasma neutral α -1,4 glucosidase activity (n . mol / ml) and liquification time (min) .(control*,oligoathenozoospermic with varicocele [*] , oligoathenozoospermic without varicocele [*]). 85