



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ





شبكة المعلومات الجامعية



شبكة المعلومات الجامعية

التوثيق الالكتروني والميكرو فيلم

جامعة عين شمس

التوثيق الالكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
على هذه الأفلام قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15 – 20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of
15 – 25c and relative humidity 20-40 %



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



شبكة المعلومات الجامعية



بالرسالة صفحات

لم ترد بالأصل

ECOLOGICAL STUDIES ON SOME INSECT PESTS ATTACKING COTTON PLANTS

B 11541

By

Dalia Mohamed Bahgat Mahmoud Shawer

B.Sc. Agriculture Entomology

Tanta University, 1997

Thesis

**Submitted in Partial Fulfillment
of the Requirements for the Degree
of Master Science**

In

ECONOMIC ENTOMOLOGY

Department of Economic Entomology

Faculty of Agriculture

Kafr El-Sheikh

Tanta University

2000

APPROVAL SHEET

Name of Student : Dalia Mohamed Bahgat Mahmoud Shawer

Title of Thesis : Ecological studies on some insect pests attacking cotton plants.

Degree : M.Sc. in Economic Entomology

Supervisory Committee:

1- Prof. Dr. Shawky Mohamed Ibrahim Metwally

Prof. of Economic Entomology and Dean of Fac. of Agric., Kafr El-Sheikh, Tanta Univ.

2- Prof. Dr. Ahmed Saad El-Deen El-Khodary

Prof. of Economic Entomology and Head of Economic Entomology Department, Fac. of Agric., Kafr El-Sheikh, Tanta Univ.

3-Dr. Ibrahim Ibrahim Mesbah

Assistant Professor of Economic Entomology, Fac. of Agric., Kafr El-Sheikh, Tanta Univ.

Approved by:


1- Prof. Dr. *A. El Khodary*

2- Prof. Dr. *R. El Khodary*

3- Prof. Dr. *S. M. I. Metwally*

4- Prof. Dr. *M. A. S. Taha*

Date: May 28, 2000



To

My Dear Father

My Lovely Mother

My Dear Husband

My Brothers and Sister

ACKNOWLEDGMENT

Firstly, ultimated thanks to God.

The present study has been suggested and supervised by **Dr. Shawky Mohamed Ibrahim Metwally**, Professor of IPM, Department of Economic Entomology and Dean of the Faculty of Agriculture, Kafr El-Sheikh, Tanta University to whom the author wishes to express her deep gratitude and sincere thanks for his valuable supervision and help, continuous encouragement and advice and constructive criticism.

Sincere thanks are also due to **Dr. Ahmed Saad El-Deen El-Khodary** Professor of Economic Entomology and Head of Department of Economic Entomology in the same Faculty, for his valuable supervision, encouragement and help throughout the whole course of that study.

Deep appreciation is extended to **Dr. Ibrahim Ibrahim Mesbah**, Assistant Professor of Economic Entomology, in the same Faculty for the time and effort he helpfully offered throughout the whole course of that study.

Deep appreciation to **Prof. Dr. Ramzy Sherif**, Head of Research, Agric. Res. Cent., Sakha, Egypt for his great help in analyzing data.

Thanks are also due to all staff members of Economic Entomology Department, Fac. of Agric., Kafr El-Sheikh, Tanta University for their help and offering facilities during that study.

CONTENTS

	Page
INTRODUCTION	1
REVIEW OF LITERATURE	3
1. The relative susceptibility of some cotton varieties to infestation with cotton bollworms.....	3
2. The predaceous arthropods in cotton fields.....	5
2.1. Survey and seasonal abundance of the most common predaceous arthropods associated with the common pests attacking cotton fields in Egypt.....	5
2.2. Effect of insecticides on some predators in cotton fields.....	7
3. The relationship between the catch of pheromone traps and boll infestation with the cotton bollworms	9
4. Utilization of the parasitoid, <i>Trichogramma evanescens</i> Westwood to control the bollworm in cotton fields	11
MATERIALS AND METHODS.....	17
1. The relative susceptibility of certain varieties of cotton to infestation with bollworms.....	17
1.1. Experimental design.....	17
1.2. Infestation rate and population densities of bollworms.....	17
2. Population densities of the most common predaceous arthropods in untreated and treated cotton fields.....	18
3. Effect of pheromones	20
3.1. The relationship between pheromone traps catch and boll infestation with cotton bollworms	20
3.2. Estimating the percent of infested bolls and number of bollworm larvae.....	20
4. The parasitoid, <i>T. evanescens</i>	21
4.1. The effectiveness of the -parasitoid <i>T. evanescens</i> in reducing damage to cotton infested by bollworm insect pests.....	21
4.2. Evaluation of the release	22
RESULTS AND DISCUSSIONS.....	23
1. The relative susceptibility of seven Egyptian cotton varieties to infestation with bollworms, <i>Pectinophora gossypiella</i> (Saund.) and <i>Earias insulana</i> (Boisd.) under field conditions.....	23
1.1. Percentage of infestation of cotton varieties with <i>P. gossypiella</i> and <i>E. insulana</i> during 1998 season.....	23

1.2.	Percentage of infestation in seven cotton varieties with <i>P. gossypiella</i> and <i>E. insulana</i> during 1999 season	28
1.3.	Average percentage of total infestation in seven cotton varieties with <i>P. gossypiella</i> and <i>E. insulana</i> as one count during 1998 and 1999 seasons.....	33
2.	Population densities of the most common predaceous arthropods in untreated and treated cotton fields.....	38
2.1.	Seasonal abundance of the most predators	38
2.1.1.	<i>Coccinella undecimpunctata</i> (Reiche)	43
2.1.2.	<i>Scymnus spp.</i>	45
2.1.3.	<i>Paederus alfieri</i> Koch.....	47
2.1.4.	<i>Chrysoperla carnea</i> Steph.	49
2.1.5.	<i>Orius spp.</i>	51
2.1.6.	<i>True spiders</i>	53
2.2.	Effect of some weather factors on seasonal abundance of predators in cotton field.....	56
3.	Effect of pheromone.....	59
3.1.	Relationship between number of the male pink bollworm moths caught in pheromone trap and boll infestation rate and number of larvae in green bolls.	59
3.2.	Relationship between number of the male spiny bollworm moths caught in pheromone trap and the boll infestation rate and number of larvae in green bolls	66
3.3.	Statistical analysis	73
4.	Utilization of the parasitoid, <i>T. evanescens</i> for the control of bollworms in cotton field.....	75
	SUMMARY	78
	REFERENCES	83
	ARABIC SUMMARY	

LIST OF TABLES

Table (1): Summary of studies on the predaceous arthropods attacking the most common insect pests which present in cotton fields in Egypt.....	6
Table (2): Conventional insecticide applications for the control of <i>P. gossypiella</i> and <i>E. insulana</i> during 1998 and 1999 cotton seasons.....	19
Table (3): Percentage of boll infestation for the tested Egyptian cotton varieties by the pink bollworm, <i>Pectinophora gossypiella</i> , at Kafr El-Sheikh region during 1998 season.....	24
Table (4): Percentage of boll infestation for the tested Egyptian cotton varieties by the spiny bollworm, <i>Earias insulana</i> , at Kafr El-Sheikh region during 1998 season.....	25
Table (5): Percentage of boll infestation for the tested Egyptian cotton varieties by <i>P. gossypiella</i> and <i>E. insulana</i> , at Kafr El-Sheikh region during 1998 season.....	26
Table (6): Number of both <i>P. gossypiella</i> and <i>E. insulana</i> larvae per 40 bolls for seven Egyptian cotton varieties, at Kafr El-Sheikh region, during 1998 season.....	27
Table (7): Percentage of boll infestation for the tested Egyptian cotton varieties by the pink bollworm, <i>Pectinophora gossypiella</i> , at Kafr El-Sheikh region during 1999 season.....	29
Table (8): Percentage of boll infestation for the tested Egyptian cotton varieties by the spiny bollworm, <i>Earias insulana</i> , at Kafr El-Sheikh region during 1999 season.....	30
Table (9): Percentage of boll infestation for the tested Egyptian cotton varieties by <i>P. gossypiella</i> and <i>E. insulana</i> , at Kafr El-Sheikh region during 1999 season.....	31
Table (10): Number of both <i>P. gossypiella</i> and <i>E. insulana</i> larvae per 40 bolls for seven Egyptian cotton varieties, at Kafr El-Sheikh region, during 1999 season.....	32
Table (11): Average percentage of boll infestation for the tested Egyptian cotton varieties by <i>P. gossypiella</i> and <i>E. insulana</i> , at Kafr El-Sheikh region during 1998 and 1999 seasons.....	35
Table (12): Average number of larvae for both <i>P. gossypiella</i> and <i>E. insulana</i> per 40 bolls for seven Egyptian cotton varieties, at Kafr El-Sheikh region, during 1998 and 1999 seasons.....	36
Table (13): Weekly numbers of the most common predaceous species on untreated cotton plants at Kafr El-Sheikh region, during 1998 season.....	38
Table (14): Weekly numbers of the most common predaceous species on untreated cotton plants at Kafr El-Sheikh region, during 1999 season.....	39

Table (15): Weekly numbers of the most common predaceous species on cotton plants received four conventional insecticidal sprayings at Kafr El-Sheikh region, during 1998 season.	40
Table (16): Weekly numbers of the most common predaceous species on cotton plants received five conventional insecticidal spraying at Kafr El-Sheikh region, during 1999 season.	41
Table (17): Total counts of the six recorded predators from cotton fields in two years study 1998 and 1999 in untreated and treated areas, reduction is presented as percentage.	42
Table (18): Simple correlation coefficients between weather factors (temperature and relative humidity) and population densities of predaceous arthropods in cotton fields, at Kafr El-Sheikh region, during 1998 and 1999 seasons.	58
Table (19): Relationship between the number of male pink bollworm moths caught in sex attractant trap and both of the percent of infested bolls and the number of pink bollworm larvae in green bolls at Kafr El-Sheikh region, during 1998 season.	60
Table (20): Relationship between the number of male pink bollworm moths caught in pheromone trap and both of the percent of infested bolls and the number of pink bollworm larvae in green bolls at Kafr El-Sheikh region, during 1999 season.	61
Table (21): Relationship between the number of male spiny bollworm moths caught in pheromone trap and both of the percent of infested bolls and the number of spiny bollworm larvae in green bolls at Kafr El-Sheikh region, during 1998 season.	67
Table (22): Relationship between the number of male spiny bollworm moths caught in pheromone trap and both of the percent of infested bolls and number of spiny bollworm larvae in green bolls at Kafr El-Sheikh region, during 1999 season.	68
Table (23): Simple correlation coefficients between the male moth numbers caught in pheromone traps and the percent of infested bolls and number of bollworm larvae per 75 bolls & the relationship between the percent of infested bolls and numbers of bollworm larvae, in cotton fields.	74
Table (24): Percent of reduction in numbers of the pink and spiny bollworm larvae in green bolls in the cotton plot treated with one-wave release technique of the parasitoid, <i>T. evanescens</i> , at Kafr El-Sheikh region, in 1999 season.	76
Table (25): Rates of infestation in green bolls in cotton plot treated with one-wave release technique of the parasitoid, <i>T. evanescens</i> and in an untreated plot, at Kafr El-Sheikh region, in 1999 season.	76

LIST OF FIGURES

Fig. 1: Average percentage of boll infestation and average number of larvae for both <i>P. gossypiella</i> and <i>E. insulana</i> per 40 bolls in seven Egyptian cotton varieties, at Kafr El-Sheikh region, during 1998 and 1999 seasons.....	37
Fig. 2: Population fluctuation of <i>Coccinella undecimpunctata</i> on untreated cotton plants, and the recorded Temp. & R.H. at the same period of study at Kafr El-Sheikh region, during 1998 and 1999 seasons.....	44
Fig. 3: Population fluctuation of <i>Scymnus</i> spp on untreated cotton plants, and the recorded Temp. & R.H. at the same period of study at Kafr El-Sheikh region, during 1998 and 1999 seasons	46
Fig. 4: Population fluctuation of <i>P. alferii</i> on untreated cotton plants, and the recorded Temp. & R.H. at the same period of study at Kafr El-Sheikh region, during 1998 and 1999 seasons.....	48
Fig. 5: Population fluctuation of <i>Chrysoperla carnea</i> on untreated cotton plants, and the recorded Temp. & R.H. at the same period of study at Kafr El-Sheikh region, during 1998 and 1999 seasons	50
Fig. 6: Population fluctuation of <i>Orius</i> spp. on untreated cotton plants, and the recorded Temp. & R.H. at the same period of study at Kafr El-Sheikh region, during 1998 and 1999 seasons.....	52
Fig. 7: Population fluctuation of spider mites on untreated cotton plants, and the recorded Temp. & R.H. at the same period of study at Kafr El-Sheikh region, during 1998 and 1999 seasons	54
Fig. 8: A comparison between the total recorded numbers of the predators during 1998 and 1999 seasons in treated and untreated areas of cotton plantations.....	55
Fig. 9: Relationship between number of male pink bollworm moths caught in sex pheromone trap and the percent of infested green bolls, at Kafr El-Sheikh region, during 1998 season.....	62
Fig. 10: Relationship between number of male pink bollworm moths caught in pheromone trap and number of pink bollworm larvae in green bolls, at Kafr El-Sheikh region, during 1998 season.....	63
Fig. 11: Relationship between number of male pink bollworm moths caught in pheromone trap and the percent of infested green bolls, at Kafr El-Sheikh region, during 1999 season.....	64
Fig. 12: Relationship between number of male pink bollworm moths caught in pheromone trap and number of pink bollworm larvae in green bolls, at Kafr El-Sheikh region, during 1999 season.....	65
Fig. 13: Relationship between number of male spiny bollworm moths caught in pheromone trap and the percent of infested green bolls, at Kafr El-Sheikh region, during 1998 season.....	69
Fig. 14: Relationship between number of male spiny bollworm moths caught in pheromone trap and number of spiny bollworm larvae in green bolls, at Kafr El-Sheikh region, during 1998 season.....	70
Fig. 15: Relationship between number of male spiny bollworm moths caught in pheromone trap and the percent of infested green bolls, at Kafr El-Sheikh region, during 1999 season.....	71

Fig. 16: Relationship between number of male spiny bollworm moths caught in pheromone trap and number of spiny bollworm larvae in green bolls, at Kafr El-Sheikh region, during 1999 season..... 72

Fig. 17: Percentage of infested green bolls in untreated and treated plots with the parasitoid, *T. evanescence* for one-wave release technique during the period extending from the last ten days of August to mid-September, at Kafr El-Sheikh region in 1999 season..... 77