THE ROLE OF TIMING IN SUCCESS OF OSSEOINTEGRATION OF IMMEDIATE IMPLANT

Thesis

Submitted to the Faculty of Oral and Dental Medicine,
Cairo University in Partial Fulfillment of Requirement for the
Doctor Degree in
Oral Surgery

By

SANAA SAMY ABD EL-AZIM

B.D.S., M.D.S. Cairo University

Faculty of Oral and Dental Medicine Cairo University

2008

Supervisors

Prof Dr. Hussein Mohammed Ali El-Tanany

Professor of Oral Surgery,
Oral Surgery Department,
Faculty of Oral and Dental Medicine,
Cairo University.

Prof. Dr. Ibrahim Abd Allah Mohammed

Professor of Oral Surgery, Oral Surgery Department, Faculty of Oral and Dental Medicine, Cairo University.

Prof. Dr. Insan Mohammed Moustafa Ashoor

Consultant Teaching Hospital Organization

Ministry of Health

/ حسين محمد علي الطناني

كلية طب الفم والأسنان جامعة القاهرة

/ إبراهيم عبدالله محمد

كلية طب الفم والأسنان جامعة القاهرة

ستشارى بالمستشفيات التعليمية

أهمية توقيت الغرس في نجاح السنية الفورية

إلى كلية طب الفم والأسنان جامعة القاهرة

الطبيبه/ سناع سامي عبدالعظيم الوريوس طب وجراحة الفم والأسنان ـ جامعة القاهرة ماجستير جراحة الفم والأسنان ـ جامعة القاهرة

> كلية طب القاهرة

ACKNOWLEDGEMENT

I would like to express my deep appreciation to **Prof Dr. Hussein Mohammed Ali El-Tanany**, Professor of Oral Surgery, Oral Surgery Department, Faculty of Oral and Dental Medicine, Cairo University, for his encouraging supervision, kind and sincere efforts, and valuable comments.

I also wish to express my sincere gratitude and thanks to **Prof. Dr. Ibrahim Abd Allah Mohammed,** Professor of Oral Surgery, Oral Surgery Department, Faculty of Oral and Dental Medicine, Cairo University, for the great efforts and helpful suggestion to accomplish this work.

I would like to express my deep appreciation and thanks to **Prof. Dr. Insan Mohammed Moustafa Ashoor,**Consultant Teaching Hospital Organization, Ministry of Health, for her continuous assistance and serious encouragement throughout the period of this study.

I would like to express my deep appreciation to **Dr. Hassan Sadek,** Maxillofacial Consultant Chief of Dental Officers Egyptian Armed Forces, for his kind assistance and great co-operation with me in making this work.

I would like to express my deep appreciation to **Prof. Dr. Mounier Ahmed Shaker**, Professor of Oral Surgery, Oral Surgery Department, Faculty of Oral and Dental Medicine, Cairo University, for his continuous assistance, helpful suggestion, and serious encouragement until this work has been developed.

CONTENTS

	Page
INTRODUCTION	1
REVIEW OF LITERATURE	3
AIM OF THE STUDY	35
PATIENTS AND METHODS	36
RESULTS	57
DISCUSSION	99
SUMMARY AND CONCLUSIONS	111
REFERENCES	114
ARABIC SUMMARY	

List of Tables

		Page
Table (1):	The criteria of Loe-Sillness for gingival index	39
Table (2):	Criteria used in recording the periodontal index according to Ramfijord	40
Table (3):	Periotest ranges group I – Time 0	74
Table (4):	Periotest ranges group I - Time 3 months post loading	75
Table (5):	Periotest ranges group I - Time 6 months post loading	76
Table (6):	Gingival index group I - 3 months post loading	77
Table (7):	Gingival index group I - 6 months post loading	78
Table (8):	Periodontal index group I - 3 months post loading	79
Table (9):	Periodontal index group I - 6 months post loading	79
Table (10):	Periotest ranges group II - Time 0	84
Table (11):	Periotest ranges group II - Time 3 months post loading	85
Table (12):	Periotest ranges group II - Time 6 months post loading	86
Table (13):	Gingival index group II - 3 months post loading	87
Table (14):	Gingival index group II - 6 months post loading	88
Table (15):	Periodontal index group II - 3 months post loading	89
Table (16):	Periodontal index group II - 6 months post loading	89

Table (17):	Periotest ranges group III - Time 0 months post loading	93
Table (18):	Periotest ranges group III - Time 3 months post loading	94
Table (19):	Periotest ranges group III - Time 6 months post loading.	95
Table (20):	Gingival index group III - 3 months post loading	96
Table (21):	Gingival index group III - 6 months post loading	97
Table (22):	Periodontal index group III - 3 months post loading	98
Table (23):	Periodontal index group III - 6 months post loading	98

List of Charts

		<u>Page</u>
Chart (1):	Normal periotest value of female maxilla: The green tolerance ranges are representing the inter personal variability.	52
Chart (2):	Normal periotest value of male maxilla: The green tolerance ranges are representing the inter personal variability.	52

List of Figure

		Page
Fig. (1):	Photograph of periapical film showing the socket after immediate extraction of the tooth	38
Fig. (2):	Photograph of panoramic x-ray taken before implantation for general evaluation of teeth and bone	39
Fig. (3):	Photograph showing prepared acrylic surgical template after fabrication	41
Fig. (4):	Photograph showing a case of group III (6 week post extraction) during infiltration anaesthesia	43
Fig. (5):	Photograph showing pilot drilling through acrylic surgical template	44
Fig. (6):	Photograph showing depth gauge used to measuring the depth of pilot site	45
Fig. (7):	Photograph showing different sizes of drills	45
Fig. (8): Fig. (9):	Photograph showing insertion of the implant fixture Photograph showing ratchet and different sizes of	46
Fig. (10):	extender/adaptor	47 47
Fig. (11):	Photograph showing sutures of the wound after implant fixture placement	48
Fig. (12):	Photograph showing porcelain crown covering the abutment which attached to the implant fixture	50
Fig. (13):	Photograph showing periotest apparatus for test implant and teeth mobility.	53

Fig. (14):	Photograph showing testing the implant mobility after exposure of the implant (6 month post insertion of the implant)	53
Fig. (15):	Photograph showing periapical film after placement of implant fixtures	55
Fig. (16):	Photograph showing panoramic x-ray six months after implantation	56
Fig. (17):	Photograph showing colored densometeric analysis of case No. 3 group I six month post implantation	56
Fig. (18):	Photograph showing a case in group I after extraction and immediate preparation of the socket	5658
Fig. (19):	Photograph showing preparation of the socket using different sizes of drills	58
Fig. (20):	Photograph showing post implantation after insertion of the abutment (6 month)	59
Fig. (21):	Photograph showing porcelain crown in place	59
Fig. (22):	Photograph showing periapical film with fracture root of upper central incisor	60
Fig. (23):	Photograph showing the tooth after extraction (Group I)	60
Fig. (24):	Photograph showing periapical film of the same case showing insertion of depth gauging in the socket	61
Fig. (25):	Photograph showing exposure of the socket of the same case	61
Fig. (26):	Photograph showing insertion of the implant in the socket after extraction (group I)	62
Fig. (27):	Photograph showing the same case after covering the implant abutment with porcelain crown	62

Fig. (28):	Photograph showing a case of missing canine after 3 weeks (Group II)	63
Fig. (29):	Photograph showing the same case in group II after incision of 3 lines pyramidal flap	63
Fig. (30):	Photograph showing the same case in group II during pilot drill	64
Fig. (31):	Photograph showing insertion of the implant fixture with bony glass powder in the prepared socket	64
Fig. (32):	Photograph showing final insertion of the implant fixture in the prepared socket	65
Fig. (33):	Photograph showing covering the implant with resorbable guided tissue regeneration membrane	65
Fig. (34):	Photograph showing suturing the wound using black silks suture.	66
Fig. (35):	Photograph showing another case in group I complain of stained and root resorption of upper central incisor	67
Fig. (36):	Photograph showing the tooth after extraction with resorbed root	67
Fig. (37):	Photograph showing the same case after drilling the site for implant fixture insertion	68
Fig. (38):	Photograph showing the same case after insertion of the implant fixture.	68
Fig. (39):	Photograph showing the case after suturing the wound with black silk	69
Fig. (40):	Photograph showing the same case 1 year post implantation with porcelain crown covering the abutment	69
Fig. (41):	Photograph of periapical film showing increase radio- opacities around the implant (6 months post implantation)	70

Fig. (42):	Showing Panoramic view six months post implantation with increase in radio-opacities	71
Fig. (43):	Densometeric analysis of upper central incisor group I of the same color case in patients and methods (Fig. 17) showing increase of bone density on one side of the implant 174 units 6 month post implantation	72
Fig. (44):	Densometeric analysis of he same case showing increase bone density on the other side of the implant to 167 unit 6 months post implantation	72
Fig.(45):	Colored densometeric analysis of upper lateral incisor (group II) showing complete bone formation on one side (green coloration) and newly bone formation on other side of the implant (Red coloration)	81
Fig. (46):	Densometeric analysis of the same case showing increase bone from top to apex (146 unit) on one side of the implant 6 months post implantation	81
Fig. (47):	Densometeric analysis of the same case showing normal bone formation (120 unit) on the other side of the implant and pocket is present after 6 months post implantation period.	82
Fig. (48):	Densometeric analysis of a case of central incisors (group III) showing increase bone density of one side up to 179 unit (3 months post implantation)	91
Fig. (49):	Colored densometeric analysis of the same case (group III) showing increase bone density with green and blue color which indicate that bone formation and increase density of this side	91
Fig. (50):	Densometeric analysis of the same case of central incisors (group III) showing increase bone density from top to apex to 212 unite on one side (6 month post implantation) of the same side	92

AIM OF THE STUDY

The aim of this study is to:

- 1. Evaluate the role of timing of implant placement in extraction site with the aid of guided tissue regeneration at different period of time after extraction.
- 2. Find out the most suitable time after tooth extraction to perform immediate implantation.

INTRODUCTION

The healthy functioning teeth is the target of every living human being, but unfortunately this target may be hard to achieve due to many factors; caries, periodontal diseases, endodontic failure and traumatic injuries⁽¹⁾.

Various attempts to replace missing teeth have presented a treatment challenge throughout the human history. Rehabilitation of the patients suffering from squeals of extracted teeth has been developed greatly^(1,2).

The need of patients increase demand for high quality services with functional and esthetically comfortable prostheses. Now every modern dental practice has to give serious thoughts to implantology⁽³⁾.

Dental implants have become one of the most exciting and rapidly developing aspects of dental practice as they now provide a clear viable treatment alternative to removable or fixed prosthesis. Implants, which are placed into the jaw bones and protrude through mucosal tissues, are used to provide attachment anchorage for replacement of artificial teeth. Advanced periodontal disease or diseases or accident can now elect a treatment option that provides for firm anchoring or bridge work