

Effect of Nicotine on the Olfactory Epithelium and Olfactory Bulb of the Rabbits

Thesis
Submitted For Partial Fulfilment of the
MD Degree In Anatomy
BY

Rima Gawad Mohamed Hamshari
M.Sc. Anatomy.
Ain Shams University.

Under the Supervision of
Prof. Dr. Sawsan Ahmed Abdel Rahman
Professor of Anatomy
Faculty of Medicine, Ain Shams University

Prof. Dr. Kariman Mohamed El-Gohari
Professor of Anatomy
Faculty of Medicine, Ain Shams University

Assist. Prof. Faten Mohamed El-Kholy
Assistant Professor of Anatomy
Faculty of Medicine, Ain Shams University

Assist. Prof. Rania Ahmed Salah El-Din
Assistant Professor of Anatomy
Faculty of Medicine, Ain Shams University

Faculty of Medicine
Ain Shams University
2012

.Acknowledgement

*First and foremost, thanks to **ALLAH** “The most merciful” whose magnificent help was factor in accomplishing this work.*

*I would like to express my sincere thanks and deepest gratitude to **Prof. Dr. Sawsan Ahmed Abd El Rahman**, Professor of Anatomy and Embryology, Faculty of Medicine, Ain Shams University, for her gracious supervision, valuable guidance, generous help, support and continuous encouragement through the whole research. I am deeply affected her noble, character, care and consideration. I am very much privileged and honoured to have her as my supervisor. To her I owe much more than I can express. I wish to be able one day to return to her a part of what she had offered to me.*

*I would like to express my sincere thanks and deepest gratitude to **Prof. Dr. Kariman Mohamed El-Gohari** Professor of Anatomy and Embryology, Faculty of Medicine, Ain Shams University, for her gracious supervision, valuable guidance, generous help, support and continuous encouragement through the whole research. I am deeply affected her noble, character, care and consideration. I am very much privileged and honoured to have her as my supervisor. To her I owe much more than I can express. I wish to be able one day to return to her a part of what she had offered to me.*

*I am deeply indebted to my supervisor **Assist. Prof. Faten Mohamed El-Kholy** Assistant Professor of Anatomy and Embryology, Faculty of Medicine, Ain Shams University for her meticulous supervision, continuous help and valuable support and guidance.*

*I am deeply Thankful to **Assist. Prof. Rania Ahmed Salah El-Din** Assistant Professor of Anatomy and Embryology, Faculty of Medicine, Ain Shams University, for her encouragement, guidance and support.*

*I owe a special acknowledgement to **Prof. Dr. Salwa Lashin and Prof. Dr. Ebtisam Ahmed Bhi El-Din** Professors of Anatomy and Embryology, Faculty of Medicine, Ain Shams University for their great assistance, encouragement, guidance and support.*

*Special and Great thanks for **Prof. Dr. Mohamed Kamal Tawfiq**, Head and Professor of Anatomy and Embryology, Faculty of Medicine, Ain Shams University for his continuous care, constant encouragement, directions, guidance and support. I will be forever thankful.*

I am very grateful to all my professors and colleagues at the Anatomy and Embryology Department, faculty of Medicine, Ain Shams University for their sincere help.

Table of Contents

List of Tables	I
List of Graphs	II
List of Figures	III
Introduction and aim of the work	1
Review of Literature	3
The Olfactory System.....	3
Tobacco Smoking.....	19
Nicotine	22
Effects of Cigarette Smoking and the Nicotine.....	28
Materials and methods	38
Results.....	44
Discussion.....	192
Summary and conclusion	206
Recommendation	210
References.....	211
Annex.....	232
Arabic Summary	

List of Tables

No.	Title	pages
Table 1	Weight of treated mother rabbits (in gm) and the doses of nicotine.	39
Table 2	Number of mothers and number of offspring (Live and Dead) in control group I, group II and group III.	45
Table 3	Gross anomalies in rabbit offspring of the three groups.	45
Table 4	Comparison between groups as regards the weight (in gm.) and the crown rump length (in cm.) and body mass index (BMI) of the offspring.	46
Table 5	Comparison between all studied groups as regards length and width of the olfactory bulb.	54
Table 6	Comparison between all studied groups as regards thickness of the olfactory epithelium.	58
Table 7	Comparison between all studied groups as regards the thickness of the olfactory bulb.	95
Table 8	Comparison between all studied groups as regards the mean width of olfactory nerve fibers layer (ONF).	97
Table 9	Comparison between all studied groups as regards the mean width of layer of glomeruli (G).	98
Table 10	Comparison between all studied groups as regards the mean width of External plexiform layer (EPL).	99
Table 11	Comparison between all studied groups as regards mean of longitudinal diameters of the mitral cells.	100
Table 12	Comparison between all studied groups as regards mean of transverse diameter of mitral cells	101
Table 13	Comparison between all studied groups as regards the mean width of mitral cell layer.	103
Table 14	Comparison between all studied groups as regards the mean width of internal plexiform layer (IPL).	104
Table 15	Comparison between all studied groups as regards width of granular layer (GR)	105

List of Graphs

No.	Title	pages
Graph 1	Comparison between all studied groups as regards weight in gm. (Wt), crown-rump length in cm (C-R) and body mass index (BMI).	47
Graph 2	Comparison between all studied groups as regards length and width in mm. of olfactory bulb.	55
Graph 3	Comparison between all studied groups as regards thickness of the olfactory epithelium in μm .	59
Graph 4	Comparison between all studied groups as regards thickness of olfactory bulb.	96
Graph 5	Comparison between all studied groups as regards the mean width of ONF.	97
Graph 6	Comparison between all studied groups as regards the mean width of layer of glomeruli (G).	99
Graph 7	Comparison between all studied groups as regards the mean width of EPL	100
Graph 8	Comparison between all studied groups as regards longitudinal diameters of the mitral cells	101
Graph 9	Comparison between all studied groups as regards transverse diameter of mitral cells.	102
Graph 10	Comparison between all studied groups as regards the mean width of mitral cell layer.	103
Graph 11	Comparison between all studied groups as regards the mean width of IPL.	104
Graph 12	Comparison between all studied groups as regards the mean width of granular layer.	105

List of Figures

No.	Title	Page
Figure 1a	Anatomy of the olfactory neural pathways, showing the distribution of olfactory receptors in the roof of the nasal cavity	5
Figure 1b	Simplified diagram of cortical regions thought to be involved in the processing of olfactory information as it passes from the olfactory epithelium to the brain	6
Figure 2	Physiological aspect of the olfactory system.	7
Figure 3	Nicotiana tabacum in flower.	20
Figure 4	The chemical structure of nicotine.	23
Figure 5	Cleopatra Cigarette.	39
Figure 6 & 7	Photograph of rabbit offspring at birth, showing the difference in crown-rump length in the three experimental groups.	48
Figure 8	Photograph of two offspring rabbits of treated group II (Tn) showing generalized oedema and hyperaemic skin in one offspring. The other offspring shows a kinked tail	49
Figure 9	Photograph of the top view of the head of rabbit offspring showing dome shaped head of offspring of group II (Tn) as compared with head of control group (Ib)	50
Figure 10	Removal of the skull cap shows the brain atrophy of the brain in the treated group II.	50
Figure 11	Sagittal section of the brain of rabbit offspring showing dilatation of the lateral ventricle of group II (Tn) as compared to control (C) group Ib.	50
Figure 12	Photograph of rabbit offspring of group III (Ts) showing generalized edema and hyperaemic skin vessels.	51
Figure 13	Photograph rabbit offspring of group III (TS).	52
Figure 14	Photograph of the undersurface of the palate showing no anomalies in the hard palate.	53
Figure 15 & 16	Photograph of the ventral aspect and lateral view of the brains of rabbit offspring showing the smaller size of the brain and the olfactory bulb from group II Tn and III Ts compared to the control group (C) Ib.	56
Figure 17	Photograph of the olfactory bulb of rabbit offspring at birth, showing increase of vascularity of the olfactory bulb which is more evident in treated group III (Ts) and apparent decrease in size of the bulb in group III (Ts) as compared to control group (C).	57

Figure 18	Photomicrograph of a section in olfactory mucosa of control rabbit offspring (group1a) showing the pseudostratified epithelium (Ep) and the lamina propria.	61
Figure 19	Photomicrograph of a section in olfactory mucosa of control rabbit offspring (group1b) showing the pseudostratified epithelium (Ep) and the lamina propria (Lp).	62
Figure 20	Photomicrograph of a section in olfactory mucosa of control rabbit offspring (group1b) is showing the epithelium (Ep) and the lamina propria (Lp) containing blood vessels (bv) and nerve fibers (Nf) and glands (G).	63
Figure 21	Photomicrograph of a section in olfactory mucosa of control rabbit offspring (group1a) showing the pseudostratified epithelium with a superficial clear zone, sustentacular cells (S) and nuclei of olfactory receptor neurons (R).	64
Figure 22	Photomicrograph of a semithin section in olfactory mucosa of control rabbit offspring (group1a) is showing the pseudostratified epithelium (Ep) and the lamina propria (LP) containing glands (G), nerve fibers (Nf) and blood vessels (bv).	65
Figure 23	Photomicrograph of a semithin section in olfactory mucosa of control rabbit offspring (group1b) showing the pseudostratified epithelium with sustentacular cells (S) and nuclei of olfactory receptor neurons (R). Few olfactory receptor neurons possess dark nuclei (d) and few show irregular outlines (ir).	66
Figure 24	Photomicrograph of a semithin section in olfactory mucosa of control rabbit offspring (group1a) showing the epithelium with olfactory receptor neurons (R), a globose basal cell (GB) and a mitotic figure (↑). Note the lamina propria containing glands (G), nerve fibers (Nf) and blood vessels (bv).	67
Figure 25	Photomicrograph of a semithin section in olfactory mucosa of control rabbit offspring (group1a) showing a nerve bundle (Nf) in the lamina propria.	68
Figure 26	Photomicrograph of a semithin section in olfactory mucosa of control rabbit offspring (group1b) showing the pseudostratified epithelium.	69
Figure 27	Photomicrograph of a section in olfactory mucosa of control rabbit offspring (group1b) showing the epithelium (Ep) and the lamina propria (Lp).	70

Figure 28	Photomicrograph of a section in olfactory mucosa of control rabbit offspring (group I a) showing the epithelium (Ep) and the lamina propria (Lp) containing glands (G), blood vessels (bv) and nerve bundles (Nf).	71
Figure 29	Photomicrograph of a section in olfactory mucosa of rabbit offspring of group II showing the arrangement of the epithelial cells (Ep) minimally disturbed with the appearance of narrow empty spaces (v) in between.	73
Figure 30	Photomicrograph of a section in olfactory mucosa of rabbit offspring of group II showing wide empty spaces (v) between the epithelial cells.	74
Figure 31	Photomicrograph of a semithin section in olfactory mucosa of rabbit offspring of group II showing the epithelium with many pale swollen olfactory receptor neurons (R) while others appear dark and shrunken (d). Sustentacular cells contain tiny vacuoles (↑) in their cytoplasm near the surface of the epithelium and few of its nuclei appear pale and swollen (S).	75
Figure 32	Photomicrograph of a section in olfactory mucosa of rabbit offspring of group II showing the pseudostratified epithelium with a diminished superficial clear zone, sustentacular cells (S) nuclei occupy upper part of epithelium and nuclei of olfactory receptor neurons (R) the lower two thirds.	76
Figure 33	Photomicrograph of a semithin section in olfactory mucosa of rabbit offspring of group II showing the epithelium with many basal cells (B) resting on a thick basal lamina (BL).	77
Figure 34	Photomicrograph of a semithin section in olfactory mucosa of rabbit offspring of group II showing the epithelium with many basal cells (B) resting on a thick basal lamina (BL). Bowman's gland duct (GD) is seen passing through the basal lamina into the epithelium.	78
Figure 35	Photomicrograph of a semithin section in olfactory mucosa of rabbit offspring of group II showing the olfactory nerve fascicles (Nf) surrounded by numerous ensheathing cells.	79
Figure 36	Photomicrograph of a section in olfactory mucosa of rabbit offspring of group II showing the pseudostratified epithelium (Ep) and lamina propria containing many blood vessels (bv). A nerve bundle (Nf) is seen surrounded by ensheathing cells.	80

Figure 37	Photomicrograph of a section in olfactory mucosa of rabbit offspring of group III showing the pseudostratified epithelium (Ep) with its laminar appearance slightly disturbed.	83
Figure 38	Photomicrograph of a semithin section in olfactory mucosa of rabbit offspring of group III showing the epithelium slightly disturbed with some empty areas (v).	84
Figure 39	Photomicrograph of a section in olfactory mucosa of a premature rabbit offspring of group III (born after 25 days) showing the pseudostratified epithelium (Ep) with its laminar appearance markedly disturbed by many empty spaces (v).	85
Figure 40	Photomicrograph of a section in olfactory mucosa of a premature rabbit offspring of group III (born after 17 days) showing the pseudostratified epithelium (Ep) with its laminar appearance greatly disturbed by many empty spaces (v).	86
Figure 41	Photomicrograph of a section in olfactory mucosa of rabbit offspring of group III showing the pseudostratified epithelium (Ep) with its olfactory receptor neurons (R) having dendrites directed towards the surface of the epithelium and axon directed toward lamina propria.	87
Figure 42	Photomicrograph of a section in olfactory mucosa of rabbit offspring of group III showing the pseudostratified epithelium with its olfactory receptor neurons (S) having dendrites directed towards the surface of the epithelium.	88
Figure 43	Photomicrograph of a section in olfactory mucosa of rabbit offspring of group III showing the pseudostratified epithelium with its olfactory receptor neurons (R) having dendrites directed towards the surface of the epithelium and axons directed towards the lamina propria.	89
Figure 44	Photomicrograph of a semithin section in olfactory mucosa of rabbit offspring of group III showing the pseudostratified epithelium with sustentacular cells (S) nuclei that occupy the upper part of epithelium and nuclei of olfactory sensory neurons (R) in the lower two thirds.	90
Figure 45	Photomicrograph of a semithin section in olfactory mucosa of rabbit offspring of group III showing the pseudostratified epithelium (Ep). Nuclei of olfactory receptor neurons (R) are vesicular with prominent nucleoli.	91

Figure 46	Photomicrograph of a semithin section in olfactory mucosa of rabbit offspring of group III showing fila olfactoria (Nf) surrounded by ensheathing cells (↑) and blood vessels (bv) and glands (G) lying in the lamina propria.	92
Figure 47	Photomicrograph of a semithin section in olfactory mucosa of a premature rabbit offspring of group III (born after 25 days of gestation) showing irregular arrangement of the olfactory epithelium.	93
Figure 48	Photomicrograph of a semithin section in olfactory mucosa of a premature rabbit offspring of group III (born after 17 days of gestation) showing irregular arrangement of the olfactory epithelium.	94
Figure 49	Photomicrograph of a coronal section of the olfactory bulb of rabbit offspring of control group I, showing closely packed cells and columnar arrangement of the neurons in all layers.	107
Figure 50	Photomicrograph of a coronal section of the olfactory bulb of rabbit offspring of control group I, showing closely packed cells and columnar arrangement of the neurons in all layers.	108
Figure 51	Photomicrograph of a coronal section of the olfactory bulb of rabbit offspring of control group I, showing the olfactory nerve fiber (ONF) interwoven in a complicated fashion.	109
Figure 52	Photomicrograph of a coronal section of the olfactory bulb of rabbit offspring of control group I, showing the glomeruli (G), the nerve fibers forming a plexus in the external plexiform layer (EPL).	110
Figure 53	Photomicrograph of a coronal section of the olfactory bulb of control offspring (group I) showing olfactory nerve fibers with insheathing cells.	111
Figure 54	Photomicrograph of a coronal section in olfactory bulb of rabbit offspring of control group (I) showing the olfactory nerve fibers with ensheathing cells entering the glomerulus (G) which contains a central capillary.	112
Figure 55	Photomicrograph of a coronal section of the olfactory bulb of offspring of control (group I) showing GFP, positive reaction (brown colour) in the subpial vessels and in few ensheathing cells along the olfactory nerve fibers layer and periglomerular.	113