ANALYTICAL STUDIES ON MAREVAN IN PURE SAMPLES, PHARMACEUTICAL PREPARATIONS AND BIOLOGICAL FLUIDS.

Presented

BY

HEBA ALI AHMED ALI MOUSSA

B.Sc. Science.

(Chemistry-Physics),

Cairo University,

2001

AThesis submitted for partial fulfillment of Master Degree In Analytical Chemistry

TO
Chemistry Department
Faculty of Science
Cairo University

(2009)

دراسات تطيلية لتقدير عقار الماريفان في الصوره النقيه

والمستحضرات الصيدليه والسوائل الحيويه.

رسالة مقدمة

من الطالبة

هبه على احمد على موسى

بكالوريوس علوم

(كيمياء-فيزياء)

جامعة القاهرة

 $(7 \cdot \cdot 1)$

إلى

قسم الكيمياء

كلية العلوم

جامعة القاهرة

للاستيفاء الجزئي للحصول على درجة التخصص الماجستير في العلوم

في

الكيمياء التحليلية

۲ . . ۹

ABSTRACT

Name: Heba Ali Ahmed Ali Moussa.

<u>Title of thesis</u>: Analytical studies on marevan in pure samples, pharmaceutical

preparation and biological fluids.

Degree: M.Sc, Un published Thesis, Faculty of Science, Cairo University, 2009.

This work has been carried out to determine warfarin anticoagulant drug in pure and

pharmaceutical preparation using new spectrophotometric reagents. This is based on the reaction

of this drug through the charge transfer complex formation reaction with DDQ, TCNQ and pCA

reagents. The sensitivity of these procedures for the spectrophotometric determination of these

drugs has been improved and the optimum practical conditions for the analysis of the drug are

selected.

Keywords: Warfarin, DDQ, TCNQ, pCA, spectrophotometry, charge transfer complex

formation reaction.

Supervisors:

1- Prof. Dr: Barsoum Nashid Barsoum.

2- Prof. Dr: Gehad Genidy Mohammed.

3- Dr: Enas Abdel-Latif.

Prof. Dr. RIFAAT HASSAN HELAL

Chairman of Chemistry Department

Faculty of Science- Cairo University

ACKNOWLEDGEMENT

First thanks to Allah for infinite help and to whom I owe my success

I wish to express my sincere thanks to **Prof. Dr. Barsoum Nashid Barsoum**, prof. of analytical Chemistry, Chemistry department, Faculty of Science, Cairo University, to whom I am deeply indebted for guidance, supervision, encouragement, suggestion, and endless support.

I would like to express my deepest gratitude and sincere appreciation to **Prof. Dr. Gehad G. Mohamed**, Prof. of Analytical Chemistry, Chemistry Department, Faculty of Science, Cairo University, for supervising this work being always available, supporting the technical work and laboratory facilities. Indeed, this work can't be completed without his help and giving me the chance to undertake the subject of this thesis.

Many thanks to **Dr. Inas Abdel-Latif**., Lecreture of biochemistry, Critical care department Cairo University Hospitals for her marvelous guidance, suggestion, and endless support.

Heba Ali

Thankful

I would like also to express my profound gratitude to my parents, wonderful brothers, sisters my fiancé and all my friends for their helpful co-operation and encouragement, through the hard times of this work till it comes to light.

Heba Ali

Table of contents

Subject	Page
Acknowledgement	
Aim of the present work	
CHAPTER I: INTRODUCTION AND LITATURE SURVEY	1
I.1. Introduction on anticoagulant drugs	1
I.1.1.Introduction on marevan (warfarin)	3
I.2.1. Biological activity of warfarin	4
1.2.1. Clinical Pharmacology	4
1.2.2. Mechanism of action:	5
1.2.3. Pharmacokinetics:	5
I.2.4.Uses of warfarin:	7
I.3. Methods of analysis of warfarin	9
I.3.1 Chromatographic methods of analysis.	9
I.3.2. Electrophoresis methods	18
I.3.3. Spectrophotometric methods	20
I.3.4. Fluorometric methods	21
I.3.5. Voltammetric methods	22
I.3.6. Potentiometric method	23
1.4. Charge- transfer complex formation in drug analysis	23
1.4.1. Introduction	23
1.4.2.Identification and structural analysis	24
I.4.3. Examples of CT complexes	25
4.3.1. CT complex formation using (DDQ) reagent	25
4.3.2. CT complex formation using. (TCNQ) reagent	29
4.3.3. CT complex formation using (pCA) reagent.	32
Chapter II: Experimental	38
II.1. Materials	38
II.2. Solutions	38
Stock solutions of pharmaceutical preparations	38



Subject	Page
II.3. Equipments	39
II.4. Procedures	39
II.4.1. Spectrophotometer determination of warfarin via Charge transfer	39
complexation reaction with TCNQ Reagent	
4.1.1. Selection of the suitable wavelength	39
4.1.2. Effect of time	39
4.1.3. Effect of temperature	39
4.1.4. Effect of TCNQ concentration	40
4.1.5. Effect of organic solvents	40
4.1.6. Stoichiometric ratio of the CT- complexes formed	40
(i) The molar ratio method (mrm)	40
(ii) The continuous variation method (cvm)	40
II.4.2. Spectrophotometric determination of warfarin with TCNQ	41
reagent	
4.2.1 Validity of Beer's Law	41
4.2.2. Day - by - day measurements	41
II.4.3. Spectrophotometric determination of warfarin in Some	
pharmaceutical preparations	41
II.4.4. Spectrophotometer determination of warfarin via Charge transfer	41
complexation reaction with DDQ and pCA reagents	
4.4.1. Selection of the suitable wavelength	41
4.4.2. Effect of time	42
4.4.3. Effect of temperature	42
4.4.4. Effect of DDQ and pCA concentration	42
4.4.5. Effect of organic solvents	42
4.4.6. Stoichiometric ratio of the CT- complexes formed	42
(i) The molar ratio method (mrm)	43
(ii) The continuous variation method (cvm)	43
II.4.5. Spectrophotometric determination of warfarin with DDQ and	43
pCA reagents	
4.5.1 Validity of Beer's Law	43
4.5.2. Day - by - day measurements	43
II.4.6. Spectrophotometric determination of warfarin in Some	
pharmaceutical preparations	44
Chapter III: Results And Discussion	45
III.1. Spectrophotometer determination of warfarin via Charge transfer	16
complexation reaction.	46



Subject	Page
1.1. Absorption spectra	47
1.2. Effect of solvent	51
1.3. Effect of reagent concentrations	54
1.4 Effect of time	56
1.6. Stoichiometry of the formed CT	60
1.7. Validity of Beer's law in microdetermination of Warfarin using DDQ,	67
TCNQ and pCA reagent	
1.8. Between–day measurements	74
1.9. Spectrophotometric determination of Warfarin in pharmaceutical	74
preparations.	
Conclusion	84
References	86
Summary	100
Arabic Summary	104



List of Schemes

Scheme (1). Fibrin formation

Scheme (2). Reaction of pCA (H₂O) with compound (B).

Scheme (3). Proton transfer from pCA to the basic center of the drug.

Scheme (4). warfarin-DDQ CT complex.

Scheme (5). warfarin-TCNQ CT complex..

Scheme (6). warfarin–PCA CT complex.



List of figures

Fig. (1) :	Structure of warfarin sodium.
Fig. (2) :	Structure of DDQ reagent.
Fig. (3) :	Structure of TCNQ reagent.
Fig. (4) :	Structure of pCA reagent.
Fig. (5) :	Absorption spectrum of warfarin-DDQ CT complex in acetonitrile.
Fig. (6) :	Absorption spectrum of warfarin-TCNQ CT complex in DMF.
Fig. (7) :	Absorption spectrum of warfarin-pCA CT complex in acetonitrile.
Fig. (8) :	Effect of organic solvents on the absorption spectra of warfarin–DDQ, warfarin-TCNQ and warfarin–pCA CT complexes
Fig. (9) :	Effect of DDQ concentration on the formation of warfarin CT complex in acetonitrile.
Fig. (10) :	Effect of TCNQ concentration on the formation of warfarin CT complex in DMF.
Fig. (11):	Effect of pCA concentration on the formation of warfarin CT complex in acetonitrile.
Fig. (12):	Effect of time on the absorbance of warfarin-DDQ CT complex in acetonitrile.
Fig. (13):	Effect of time on the absorbance of warfarin-TCNQ CT complex in DMF.
Fig. (14):	Effect of time on the absorbance of warfarin-pCA CT complex in acetonitrile
Fig. (15):	Effect of temperature (0-50 °C) on the absorbance of CT complex of warfarin with DDQ
	in acetonitrile.
Fig. (16) :	Effect of temperature (10-50 °C) on the absorbance of CT complex of warfarin with
	TCNQ in DMF.
Fig. (17) :	Effect of temperature (0-50 °C) on the absorbance of CT complex of warfarin with pCA
	in acetonitrile.
Fig. (18) :	Job's method for warfarin- DDQ CT complex in acetonitrile.

Fig. (19) :	Job's method for warfarin-TCNQ CT complex in DMF.
Fig.(20) :	Job's method for warfarin-pCA CT complex in acetonitrile.
Fig. (21):	Molar ratio of warfarin-DDQ CT complex in acetonitrile.
Fig.(22) :	Molar ratio of warfarin-TCNQ CT complex in DMF.
Fig. (23) :	Molar ratio of warfarin-pCA CT complex in acetonitrile.
Fig. (24) :	Validity of Beer's law of the CT reaction between DDQ and warfarin at selected
	optimum conditions
Fig. (25) :	Validity of Beer's law of the CT reaction between TCNQ and warfarin at selected
	optimum conditions
Fig. (26) :	Validity of Beer's law of the CT reaction between pCA and warfarin at selected optimum
	conditions
Fig. (27) :	A comparative study between the official method and the proposed method in the
	determination of warfarin using DDQ reagent.
Fig. (28) :	A comparative study between the official method and the proposed method in the
	determination of warfarin using TCNQ reagent.
Fig. (29) :	A comparative study between the official method and the proposed method in the
	determination of warfarin using pCA reagent



List of tables

Table No.		Page No.
Table (1)	The molar absorptivity values of warfarin-DDQ CT complex in different	52
	solvents	
Table (2)	The molar absorptivity of warfarin-TCNQ CT complex in different	52
	solvents	
Table (3)	The molar absorptivity of warfarin-pCA CT complex in different	53
	solvents	
Table (4)	Microdetermination of warfarin via its reaction with DDQ at $\lambda_{max} = 468$ nm, t =	68
	25 min, T = 35 °C	
Table (5)	Microdetermination of warfarin via its reaction with TCNQ at $\lambda_{max} = 847$ nm, $t =$	69
	25 min, T = 30-40 °C.	
Table (6)	Microdetermination of warfarin via its reaction with PCA at λ_{max} = 5+13 nm, t =	70
	25 min, T = 25 °C	
Table (7)	Spectral characteristics of warfarin CT coloured reaction products and the	
	analytical characteristics (accuracy and precision) of its reaction with DDQ,	
	TCNQ and PCA	73
Table (8)	Between – day precision of the determination of warfarin using DDQ.	75
Table (9)	Between – day precision of the determination of warfarin using TCNQ.	76
Table (10)	Between – day precision of the determination of warfarin using PCA.	77
	Between - day precision of the determination of warrann using FCA.	, ,



Table No.		Page No.
Table (11)	Spectrophotometric microdetermination of warfarin in different pharmaceutical preparations using DDQ and official methods	78
Table (12)	Spectrophotometric microdetermination of warfarin in different pharmaceutical preparations using TCNQ and official methods.	79
Table (13)	Spectrophotometric microdetermination of warfarin in different pharmaceutical preparations using pCA and official methods	80





Abbreviation

2,6-DAPY: 2,6-diaminopyridine

APH: amineptine

ATPH: amitriptyline

BCG: bromocresol green

BCP: bromocresol purple

BDMAENDI: N,N'-bis-[2-N,N-dimethylaminoethyl)]-1,4,6,8-naphthalenediimide

BHENDI: N,N'-bis[2-hydroxyethyl)]-1,4,6,8-naphthalenediimide

CE: capillary electrophoresis

CIP: ciprofloxacin

CPR: chlorophenol red

CQP: chloroquine phosphate

CSPs: chiral stationary phases.

CT: charge-transfer

DDQ: 2, 3-dichloro-5, 6-dicyanobenzoquinon.

DMF: *N*,*N*-dimethylformamide medium

DPP: disopyramide phosphate

β- \Box CD : β-cyclodextrin

EKC: electrokinetic chromatography

FL: 2,7-dinitro-9-fluorenone

FLE: fleroxacin

GC: gas chromatography

HPLC: high performance liquid chromatography

