

Ain Shams University Faculty of Pharmacy Microbiology & Immunology Dept.

Inactivation of *Pseudomonas aeruginosa* quorum sensing signals by some soil bacterial isolates

A Thesis

Submitted in Partial Fulfillment of the Requirments for the

Master's Degree

In

Pharmaceutical Sciences (Microbiology and Immunology)

By

Masarra Mohammad Abo-el-soud Sakr

Bachelor of Pharmaceutical Sciences, Faculty of Pharmacy, Ain Shams University, 2006



Ain Shams University Faculty of Pharmacy Microbiology & Immunology Dept.

Inactivation of *Pseudomonas aeruginosa* quorum sensing signals by some soil bacterial isolates

A Thesis

Submitted in Partial Fulfillment of the Requirments for the Master's Degree

In

Pharmaceutical Sciences (Microbiology and Immunology)

By

Masarra Mohammad Abo-el-soud Sakr

Bachelor of Pharmaceutical Sciences, Faculty of Pharmacy, Ain Shams University, 2006

Under Supervision of

Prof. Dr. Nadia A. El-Haleem Hassouna, PhD

Professor of Microbiology and Immunology, Faculty of Pharmacy, Ain Shams University

Prof. Dr. Mohammad Mabrouk Aboulwafa, PhD

Professor of Microbiology and Immunology Vice Dean for Community Service and Environmental Development,

Faculty of Pharmacy, Ain Shams University

Dr. Khaled Mohamed Anwar Aboshanab, PhD

Assisstant Professor and Acting Head of Microbiology and Immunology Department
Faculty of Pharmacy, Ain Shams University

Acknowledgement

First, I thank "Allah" for granting me the power to accomplish this work.

I would like to express my deepest thanks to **Prof. Dr. Nadia A. El-Haleem Hassouna**, Professor of Microbiology and Immunology, and founder of the Microbiology and Immunology Department, Faculty of Pharmacy, Ain Shams University, for her valuable scientific supervision, constructive advice and continuous guidance throughout the work.

My deepest gratitude and appreciation is expressed to **Prof. Dr. Mohammad Mabrouk Aboulwafa**, Vice Dean for Community Service and Environmental Development and former Head of Microbiology and Immunology Department, Faculty of Pharmacy, Ain Shams University, for suggesting the point, planning the work, providing continuous scientific supervision, kindly supplying laboratory facilities whenever needed and for his constructive criticism throughout this study.

I am also greatly indebted to **Dr. Khaled Mohammed Aboshanab**, Assistant Professor and Acting Head of Microbiology and Immunology Department, Faculty of Pharmacy, Ain Shams University, for his divine support, following up and giving his valuable time and effort. I also thank him for providing assistance throughout the genetics work in this study.

Grateful thanks are extended to my dear **colleagues** and to all the **workers** at the Faculty of Pharmacy, Ain Shams University for their help and support during this work.

My everlasting thanks and appreciation are also directed to my beloved **family** and **friends** for continuous support, encouragement and sincere help throughout my whole life.

Last but not least, I would like to express my deepest and most sincere gratitude to my **husband** for his patience, understanding, motivation and everlasting support.

و آخر دعوانا أن الحمد لله رب العالمين.

Masarra Mohammad Sakr

List of Abbreviations

AHL	A aul homograina lagtona
	Acyl homoersine lactone
Asp	Aspartate
ATCC	American Type Culture Collection
BHL	Butyryl homoserine lactone
C4-HSL	Butyryl homoserine lactone
C6-HSL	Hexanoyl homoserine lactone
C-7 HSL	Heptanoyl homoserine lactone
C-8 HSL	Octanoyl homoserine lactone
Dil 10	10 fold culture dilution
EDTA	Ethylene diamnine terta-acetic acid
HHL	Hexanoyl homoserine lactone
His	Histidine
HSL	Homoserine lactone
LB	Lauria Bertani
ORF	Open Reading Frame
QMEAN	Qualitative model energy analysis
QS	Quorum sensing
Rf	Retardation factor
RNase	Ribonuclease
SDS	Sodium dodecyl sulphate
SET	Sodium chloride-EDTA-Tris buffer
TAE	Tris-Acetic acid-EDTA buffer
TE	Tris-EDTA buffer
TLC	Thin Layer Chromatography
Tyr	Tyrosine
Undil	Undiluted culture
YMG	Yeast extract-Malt extract-Glucose

Contents

Tit	tle	Page
Ab	ostract	
Int	troduction	1
Lit	terature Review	4
1.	Quorum sensing definition	4
2.	Historical Background	
3.	What does Quorum sensing regulate in bacterial cell	6
	3.1. Virulence	
	3.2. Biofilm	7
	3.3. Bioluminescence	8
	3.4. Pigmentation	9
	3.5. Secondary metabilote production	9
	3.6. Motility and clumping	9
	3.7. Sporulation	9
	3.8. Extracellular polysaccharide	10
	3.9. Fruiting body formation	10
	3.10. Effects on Eukaryotic cells	
	3.11. Bacterial cross talk and Symbiosis	11
	3.12. Antimicrobial activity	12
4.	Quorum sensing systems	12
	4.1. Gram negative bacteria	12
	4.2. Gram positive bacteria	
	4.3. Hybrid system	
5.	\mathcal{E}	
	5.1. AHL signaling	15
	5.2. AI-2 signaling	
	5.3. AI-3 signaling	19
6.	Quorum sensing in Pseudomonas aeruginosa	
	6.1. AHL signaling	
	6.2. Quinolone signalling	
7.	Extraction and detection of AHL	24
	7.1. Extraction of AHL	24
	7.2. Detection AHL	
	7.2.1. Biological detection	
	7.2.1.1. Chromobacterium violaceum	
	7.2.1.2. Other commonly used biosensors	25
	7.2.2. Chemical detection	26

8.	Quorum sensing inhibition as a new chemotherapeutic approach	. 26
9.	Strategies to interfere with quorum sensing	. 27
	9.1. Drug target: the signal generator	. 28
	Targeting AHL synthsis	. 28
	Targeting AI-2 synthesis	. 29
	9.2. Drug Target: the signal molecule	. 30
	Chemical degradation	. 30
	Enzymatic degradation	. 30
	AHL lactonasaes	. 32
	AHL acylases	. 34
	AHL metabolism	. 36
	9.3. Drug target: the signal receptor	. 36
	Targeting AHL Receptor	. 36
	Targeting AI-2 Receptor	
Ma	aterials and Methods	. 39
1.		
	1.1. Chromobacterium violaceum CV026	. 39
	1.2. Pseudomonas aeruginosa isolates	
	1.3. Bacterial isolates from soil	. 42
2.	Chemicals	. 43
3.	Instruments	. 45
4.	Media	. 46
	4.1. Ready made media and media components	
	4.2. In-house formulated media	. 47
	4.2.1. Lauria Bertani broth	. 47
	4.2.2. Lauria Bertani agar	. 47
	4.2.3. Lauria Bertani semisolid agar	. 47
	4.2.4. Starch nitrate agar	. 48
	4.2.5. YMG broth	. 48
	4.2.6. Slant medium	
	4.2.7. Medium used for bacterial suspension in lyophilization	. 48
5.	Reagents, Buffers and Solutions	. 49
	5.1. Buffers used for bacterial cell washing and lysis	. 49
	5.1.1. Cell washing buffer	
	5.1.2. Cracking buffer	. 49
	5.2. Protein assay reagents and standard solutions	. 49
	5.2.1. Complex-forming reagent	. 49
	5.2.2. Folin reagent	
	5.2.3. Bovine Serum Albumin Standard Solutions	
	5.3. Buffers used in quorum sensing inhibitory activity characterization	
	5.3.1. Mcilvaine buffer	
	5.3.2. Glycine NaOH buffer	. 50

	5.4. Reagents, buffers and solutions used in DNA manipulation technique	es
		50
	5.4.1. Tris-HCl (1M,pH 7.5)	50
	5.4.2. SET buffer	51
	5.4.3. TE buffer	51
	5.4.4. TAE buffer (50X)	51
	5.4.5. DNA loading buffer	52
	5.4.6. EDTA solution (0.5 M, pH 8)	52
	5.4.7. Sodium dodecyl sulphate (SDS) 10% solution	52
	5.4.8. NaCl 5 M solution	
	5.4.9. Ethidium Bromide Solution (5 mg/ml)	53
	5.4.10. Chloroform/isolamyl alcohol mixture	53
	5.4.11. Agarose gel for electrophoresis	53
	5.4.12. DNA ladder for gel electrophoresis	53
	5.4.13. Master Mix for PCR	54
	5.5. Other buffers, reagents and solutions	54
	5.5.1. Hexanoyl homoserine lactone stock solution	54
	5.5.2. Kanamycin sulphate solution	54
	5.5.3. Acidified ethyl acetate solution	55
	5.5.4. Crystal violet solution	55
	5.5.5. M63 Buffer	55
	5.5.6. Phosphate Buffered Saline	55
6.	Extraction of acyl homoserine lactone signal molecules from Pseudomon	as
	aeruginosa isolates	
7.	TLC analysis of the extraced AHL signals of <i>P. aeruginosa</i>	56
8.	Collection of soil samples	57
9.	Recovery of bacterial isolates from soil	57
10.	Screening of the recovered bacterial soil isolates for quorum sensing	
	inhibitory activity	
	10.1. Screening of the bacterial isolates recovered on R ₂ A	
	10.1.1. Primary screening using the synthetic signal molecule hexano	
	homoserine lactone (HHL)	58
	10.1.2. Secondary screening of the bacterial soil isolates with HHL	
	degrading activity against quorum sensing signals of some <i>P</i> .	
	aeruginosa extracts	59
	10.2. Screening of <i>Streptomyces</i> isolates	
	10.2.1. Primary screening using the synthetic signal molecule HHL	60
	10.2.2. Secondary screening of the <i>Streptomyces</i> isolates with HHL	
	degrading activity against quorum sensing signals of some P .	
	aeruginosa extracts	
11.	Characterization of quorum sensing inhibitory activity of tested bactertial	
	soil isolates	61

11.1. Characterization of quorum sensing inhibitory activity of bactertia	
soil isolates recovered on R ₂ A agar	61
11.1.1. Using whole cell culture	61
11.1.1.1. Quorum sensing inhibitory activity of tested isolates at	
different levels of substrate concentration and culture	
dilutions	61
11.1.1.2. Quorum sensing inhibitory activity of some selectd	
isolates at shorter reaction time	62
11.1.1.3. Quorum sensing inhibitory activity of some selectd	
isolates at different incubation temperatures	62
11.1.1.4. Quorum sensing inhibitory activity spectrum of some	
selectd isolates	62
11.1.1.5. Detection whether the quorum sensing inhibitory	
activity of test isolates is extracellular or cell associated	63
11.1.2. Using crude bacterial cell lysate of some selected isolates	63
11.1.2.1. Preparation of crude cell lysates of the selected isolates	
	63
11.1.2.2. Measurements of the totel protein concentrations of the	
prepared crude enzyme extracts	64
11.1.2.3. Quorum sensing inhibitory activity of the crude enzym	e
extracts as determined by well diffusion method	
11.1.2.4. Studying physical parameters that may affect the crude	<u>;</u>
enzyme stability of the selected isolates	
11.1.2.4.1. Thermal stability	
11.1.2.4.2. Stability at different pH values	66
11.1.2.5. Effect of some divalent metals and EDTA on activity	67
11.1.2.6. Effect of adding the signal molecule HHL to the growth	
medium on the enzyme productivity of the selected isolates . (67
11.1.2.7. Catalytic parameters of crude enzyme extracts of the	
selected isolates	68
11.1.2.7.1. Catalytic activity rates of the crude enzyme	
extracts	
11.1.2.7.2. Effect of temperature on catalytic activity	
11.1.2.7.3. Effect of pH on catalytic activity	
11.1.2.7.4. Kinetic constants, Vmax and Km	
11.1.2.7.5. Enzyme substrate specificity	72
11.2. Characterization of quorum sensing inhibitory activity of	
Streptomyces isolates	
11.2.1. Quorum sensing inhibitory activity of tested isolates at different	
levels of substrate concentration	
11.2.2. Quorum sensing inhibitory activity of the cell free supernatant	
as determined by well diffusion method	
11.2.3. Studying physical parameters that may affect the crude enzyments	
stability of the selected isolate	74

	11.2.3.1. Thermal stability	74
	11.2.3.2. Stability at different pH values	75
	11.2.4. Effect of some divalent metals and EDTA on activity	
	11.2.5. Catalytic parameters of crude enzyme extract of the selected	
	isolate	75
	11.2.5.1. Catalytic activity rate of the crude enzyme extract	75
	11.2.5.2. Effect of temperature on catalytic activity	75
	11.2.5.3. Effect of pH on catalytic activity	76
	11.2.5.4. Kinetic constants, Vmax and Km	76
	11.2.5.5. Enzyme substrate specificity	76
12.	Detection of lactonase gene in some selected isolates recovered onto R ₂ A	1
	agar by PCR	
	12.1. Chromosomal DNA extraction	
	12.2. Design and preparation of primers	78
	12.2.1. Collection of data from the NCBI GenBank database	
	12.2.2. The alignment of genes using clustalW	78
	12.2.3. Design of the primers	78
	12.2.4. Checking the specificity of the primers	
	12.2.5. Preparation of the primers for PCR	78
	12.3. PCR amplification of lactonase gene	79
	12.4. Agarose gel electrophoresis	
13.	Sequencing of the amplified lactonase genes of test isolates	
	13.1. PCR clean up	
	13.2. Sequencing	80
	13.3. Assembling of sequence data and alignment against protein	
	homologues in database	
	Prediction of the tertiary structure of the lactonase enzymes	
15.	Identification of the selected isolates	
	15.1. Identification of the selected isolates recovered on R_2A agar	
	15.1.1. Gram stain	
	15.1.2. 16S ribosomal RNA gene analysis	
	15.2. Identification of the selected <i>Streptomyces</i> isolate	81
ъ.	K	00
	TIC and being falls and the day of AIH along to a fall and the control of the con	82
1.	TLC analysis of the extracted AHL signals of <i>P. aeruginosa</i> isolates	
2.	Recovery of bacterial isolates from soil	83
3.	Screening of the recovered bacterial soil isolates for quorum sensing	02
	inhibitory activity	
	3.1. Screening of the bacterial isolates recovered on R ₂ A	
	3.1.1. Primary screening using the synthetic signal molecule HHL	03
	3.1.2. Secondary screening of the bacterial soil isolates with HHL	
	degrading activity against quorum sensing signals of some <i>P. aeruginosa</i> extracts	Q <i>C</i>
	· ·	
	3.2. Screening of <i>Streptomyces</i> isolates	00

3.2.1. Primary screening using the synthetic signal molecule HHL 88
3.2.2. Secondary screening of the <i>Streptomyces</i> isolates with HHL
degrading activity against quorum sensing signals of some P.
aeruginosa extracts89
4. Characterization of quorum sensing inhibitory activity of tested bactertial
soil isolates90
4.1. Characterization of quorum sensing inhibitory activity of bactertial
soil isolates recovered on R ₂ A agar90
4.1.1. Using whole cell culture
4.1.1.1. Quorum sensing inhibitory activity of tested isolates at
different levels of substrate concentration and culture
dilutions90
4.1.1.2. Quorum sensing inhibitory activity of some selectd
isolates at shorter reaction time
4.1.1.3. Quorum sensing inhibitory activity of some selectd
isolates at different incubation temperatures95
4.1.1.4. Quorum sensing inhibitory activity spectrum of some
selectd isolates
4.1.1.5. Detection whether the quorum sensing inhibitory activity
of test isolates is extracellular or cell associated
4.1.2. Using crude bacterial cell lysate of some selected isolates 100
4.1.2.1. Quorum sensing inhibitory activity of the crude enzyme
extracts as determined by well diffusion method
4.1.2.2. Studying physical parameters that may affect the crude enzyme stability of the selected isolates
4.1.2.2.1. Thermal stability 101
4.1.2.2.1. Thermal stability
4.1.2.3. Effect of some divalent metals and EDTA on activity . 102
4.1.2.4. Effect of adding the signal molecule HHL to the growth
medium on the enzyme productivity of the selected isolates
4.1.2.5. Catalytic parameters of crude enzyme extracts of the
selected isolates
4.1.2.5.1. Catalytic activity rates of the crude enzyme
extracts
4.1.2.5.2. Effect of temperature on catalytic activity 107
4.1.2.5.3. Effect of pH on catalytic activity
4.1.2.5.4. Kinetic constants, Vmax and Km
4.1.2.5.5. Enzyme substrate specificity
4.2. Characterization of quorum sensing inhibitory activity of <i>Streptomyces</i>
isolates
4.2.1. Quorum sensing inhibitory activity of tested isolates at different
levels of substrate concentration

	4.2.2. Quorum sensing inhibitory activity of the cell free supernata	nts
	as determined by well diffusion method	113
	4.2.3. Studying physical parameters that may affect the crude enzy	me
	stability of the selected isolate	114
	4.2.3.1. Thermal stability	114
	4.2.3.2. Stability at different pH values	115
	4.2.4. Effect of some divalent metals and EDTA on activity	116
	4.2.5. Catalytic parameters of crude enzyme extract of the selected	
	isolate	117
	4.2.5.1. Catalytic activity rate of the crude enzyme extract	117
	4.2.5.2. Effect of temperature on catalytic activity	
	4.2.5.3. Effect of pH on catalytic activity	119
	4.2.5.4. Kinetic constants, Vmax and Km	
	4.2.5.5. Enzyme substrate specificity	
5.	\mathcal{C}	
6.	Sequencing of the amplified lactonase genes of test isolates	
7.	J	
8.		
	8.1. Identification of the selected <i>Bacillus</i> isolates	
	8.2. Identification of the selected <i>Streptomyces</i> isolate	
	scussion	_
1.	TLC analysis of the extracted AHL signals of <i>P. aeruginosa</i> isolates	
2.	J	131
3.		
	inhibitory activity	
	3.1. Screening of the bacterial isolates recovered on R ₂ A	
4	3.2. Screening of <i>Streptomyces</i> isolates	
4.	Characterization of quorum sensing inhibitory activity of tested bacterti	
	soil isolates	
	4.1. Characterization of quorum sensing inhibitory activity of bactertial	
	soil isolates recovered on R ₂ A agar	
	4.1.1. Using whole cell culture	
	4.1.2. Using crude bacterial cell lysate of some selected isolates 4.2. Characterization of quorum sensing inhibitory activity of <i>Streptomy</i>	
	isolates	
5.	Detection of lactonase gene in the selected <i>Bacillus</i> isolates by PCR	
<i>5</i> . 6.	Sequencing of the amplified lactonase genes of test isolates	
7.	Prediction of the tertiary structure of the lactonase enzymes	
8.	Identification of the selected isolates	
9.		
	ımmary	
	eferences	
Ap	opendix	197
_		

List of Tables

Table
Table 1: The most commonly identified AHLs produced by different bacterial species and their LuxI/R homologs 17
Table 2: Functions controlled by the quorum sensing circuity in P. aeruginosa 23
Table 3: AHL-degrading enzymes described for various bacterial species 31
Table 4: The relative QS signal productivities of <i>P. aeruginosa</i> isolates used in the present study
Table 5: Different chemicals used in the present study arranged alphabetically
Table 6: Instruments used in the study arranged alphabetically
Table 7: Primary screening resuts for quorum sensing inhibitory activity of different bacterial isolate categories recovered on R ₂ A agar
Table 8: Degrading activity profile of tested Bacillus isolates against AHL signals contained in the extracts of seven P. aeruginosa isolates
Table 9: Primary screening results for quorum sensing inhibitory activity of the recovered Streptomyces isolates 88
Table 10: Degrading activity profile of tested <i>Streptomyces</i> isolates' supernatants against AHL signals contained in the extracts of seven <i>P. aeruginosa</i> isolates 89
Table 11: Comparison of HHL degrading activity and activity spectrum of some tested Bacillus isolates against P. aeruginosa extracts
Table 12: AHL degrading activity of the selected Bacillus isolates as determined by well diffusion method 100
Table 13: Percentage relative activities of crude enzyme extracts when HHL was added to the growth medium as compared to controls 105

Table 14: Catalytic activity rates of crude enzyme extracts of the selected Bacillus isolates using HHL as a substrate 107
Table 15: Kinetic constants, Vmax and Km for each of the tested crude enzyme extracts of the selected <i>Bacillus</i> isolates
Table 16: Enzyme substrate specificity of the crude enzyme extracts of selected Bacillus isolates against different AHL standards 112
Table 17: AHL degrading activity of the tested Streptomyces isolates as determined by well diffusion method 114
Table 18: Effect of temperature on AHL degrading catalytic activity of crude enzyme extract of isolate St62
Table 19: Effect of pH on AHL degrading catalytic activity of crude enzyme extract of isolate St62 119
Table 20: Enzyme substrate specificity of the crude enzyme extract of selected Streptomyces isolate St62 against different AHL standards
Table 21: Result of identification of the selected Bacillus isolates by 16S rRNA gene analysis 127

List of Figures

Figure Page
Figure 1. Phenotypes shown to be regulated by quorum sensing in Gram negative and Gram positive bacteria
Figure 2. QS systems in Gram negative bacteria and in Gram positive bacteria.
Figure 3. Hybrid quorum sensing system found in <i>V. harveyi</i> 14
Figure 4. Synthesis of acyl homoserine lactone
Figure 5. Synthesis of AI-2 signal from SAM in <i>V. harveyi</i> and <i>Salmonella typhimurium</i>
Figure 6. The hierarchical <i>Pseudomonas aeruginosa</i> quorum-sensing systems
Figure 7. Strategies to interefere with quorum sensing
Figure 8. Structures of AHL synthase inhibitors
Figure 9. LuxS inhibitor synthesized by Zhao and his coworkers 29
Figure 10. The possible ways of enzymatic degradation of AHLs 32
Figure 11. AHL lactonases hydrolyze the lactone ring in the homoserine moiety of AHLs, without affecting the rest of the signal molecule structure 33
Figure 12. Acylase hydrolyze the amide bond between the acyl side chain and the homoserine lactone in the AHL molecules generating the free fatty acid and the homoserine lactone
Figure 13. penicillic acid (PA) and patulin produced by <i>Pe. Radicicola</i> and <i>Pe.coprobium</i> , respectively
Figure 14. Some examples of furanones isolated from extracts of <i>D. pulchra</i> 37

Figure 15. A DPD analogue that acts as a potent QS inhibitor
Figure 16. Calibration curve for bovine serum albumin as determined by Lowry's method
Figure 17. Calibration curve showing the relationship between HHL concentrations and diameters of purple color zones formed as determined by well diffusion method. 70
Figure 18. Calibration curve showing the relationship between BHL concentration and the diameters of purple color zones formed as determined by well diffusion method
Figure 19. Calibration curve showing the relationship between C7-HSL concentration and the diameters of purple color zones formed as determined by well diffusion method
Figure 20. Calibration curve showing the relationship between C8-HSL concentration and the diameters of purple color zones formed as determined by well diffusion method
Figure 21. TLC chromatograms of the extracted AHL signals of <i>P. aeruginosa</i> isolates as detected by CV026 biosensor. 82
Figure 22. Categorization of bacterial isolates recovered on R ₂ A agar according to their microscopical characters
Figure 23. Results of primary screening done in microtiter plate as described by Jafra and van-der-Wolf
Figure 24. AHL degrading activity of tested Bacillus isolates at different concentrations of HHL when tested at two culture dilution levels 91
Figure 25. AHL degrading activity of some selected <i>Bacillus</i> isolates at different culture dilutions
Figure 26. AHL degrading activity of some selected <i>Bacillus</i> isolates at different reaction times