

# **INCREASING THE SOLUBILITY AND BIOAVAILABILITY OF A SLIGHTLY SOLUBLE DRUG USING MICROEMULSIONS**

*A thesis submitted in the partial fulfillment of the requirements for the  
PhD Degree of Pharmaceutical Sciences (Pharmaceutics)*

by

**Rania Mohammed Hafez Mohammed Hathout**

Masters of Pharmaceutical Science, 2006, Ain Shams University  
Teaching assistant, Department of Pharmaceutics, Faculty of Pharmacy,  
Ain Shams University

*Under the supervision of*

**Prof. Dr. Ahmed Shawky Geneidi**

*Professor of Pharmaceutics and Industrial Pharmacy  
Faculty of Pharmacy, Ain Shams University*

**Prof. Dr. Nahed Daoud  
Mortada**

*Professor of Pharmaceutics  
Dean of Faculty of Pharmacy  
Ain Shams University*

**Dr. Samar Mansour Holayel**

*Assistant Professor of Pharmaceutics  
Faculty of Pharmacy,  
Ain Shams University*

**Ain Shams University  
Faculty of Pharmacy  
Department of Pharmaceutics  
Cairo  
2010**

***INCREASING THE SOLUBILITY  
AND BIOAVAILABILITY  
OF A SLIGHTLY SOLUBLE  
DRUG USING  
MICROEMULSIONS***

## ***Acknowledgement***

First and foremost thanks to **God** for helping me to fulfill this work.

I want to express my deepest appreciation and sincere gratitude to Prof. Dr. Ahmed Shawky Geneidi, Professor of Pharmaceutics and Industrial Pharmacy, Faculty of Pharmacy, Ain Shams University, for his instructive supervision, kind help and generous attitude throughout the development of this work.

I would also like to express my deepest thanks and sincere appreciation to Prof. Dr. Nahed Daoud Mortada, Professor of Pharmaceutics, Faculty of Pharmacy, Ain Shams University, for her valuable guidance, great support and precious advice and for providing great help for the accomplishment of this work.

I would like to express my thanks and appreciation to Assistant Prof. Dr. Samar Mansour Holayel, Assistant Professor of Pharmaceutics, Faculty of Pharmacy, Ain Shams University for the great help and effort she devoted for the completion of this thesis, for her instructive supervision, valuable advice, encouragement and support.

I would like to express my profound gratitude and appreciation to Prof. Dr. Richard Henry Guy, Professor of pharmaceutical sciences, Department of Pharmacy and Pharmacology, University of Bath, UK, for his great help, advice, support, encouragement and effort he offered for the completion of this thesis, accomplishing the novel work in it and bringing the new ideas into reality.

I also like to thank the members of the department of Pharmaceutics, Ain Shams University especially Dr, Amany Osama, AbdelKader Ali and Maha Nasr for their help and support.

I would also like to thank the staff members of the Department of Pharmacy and Pharmacology, University of Bath at the UK, for their help and support in accomplishing the practical part of this thesis.

I would also like to express my gratitude to all of my friends who supported and encouraged me all through the way.

Finally, no words can express my deepest thanks to my parents for their great support and infinite patience throughout the work accomplished in this thesis.

***Dedication***

*I dedicate this thesis to my dear father, mother and siblings, and to the soul of my grandma.*

## **List of Contents**

<b>Item</b>	<b>Page</b>
<b>List of Abbreviations</b>	VII
<b>List of Tables</b>	VIII
<b>List of Figures</b>	XI
<b>Abstract</b>	XVII
<b>General Introduction</b>	1
<b>Scope of Work</b>	22
<b>Chapter 1: Preparation and Screening of Different Topical Microemulsion Systems</b>	26
<b>Introduction</b>	26
<b>Experimental</b>	34
Methodology	35
I- High performance liquid chromatography (HPLC) for testosterone in ethyl alcohol (95%)	35
II- Determination of the saturation solubility of testosterone in different oils	35
III- Construction of pseudo-ternary phase diagrams	36

<b>Item</b>	<b>Page</b>
-------------	-------------

IV- Evaluation of the prepared microemulsions	36
<b>Results and Discussion</b>	38
I- High performance liquid chromatography (HPLC) of testosterone in ethyl alcohol (95%)	38
II- Saturation solubility of testosterone in different oils	41
III- Pseudo-Ternary phase diagrams	41
IV- Evaluation of the prepared microemulsions	52
<b>Conclusions</b>	57
<b>Chapter 2: Methods Development for the Analysis of Components of a Microemulsion System Using Liquid Chromatography Mass Spectroscopy (LC / MS)</b>	58
<b>Introduction</b>	58
<b>Experimental</b>	63
Methodology	64
I- Development of a novel quantitative method of analysis for oleic acid using LC / MS system	64
II- Development of a novel quantitative method of analysis for Tween20 using LC / MS system	64

<b>Item</b>	<b>Page</b>
III- Development of a novel quantitative method of analysis for Transcutol using LC / MS system	65

<b>Results and Discussion</b>	67
I- Quantitative analysis of oleic acid	67
II- Quantitative analysis of Tween20	71
III- Quantitative analysis of Transcutol	75
<b>Conclusion</b>	80
<b>Chapter 3: <i>Ex-Vivo</i> methods to Assess the Mechanisms of Stratum Corneum Enhancement by Topical Microemulsions</b>	81
<b>Introduction</b>	81
<b>Experimental</b>	93
Methodology	95
I- Preparation of selected microemulsion formulations	95
II- Studing the effects of the microemulsion components on the skin barrier	96
A) ATR-FTIR studies	96
i) Processing	96
<b>Item</b>	<b>Page</b>
ii) Spectral analysis	98
B) Tape stripping followed by ATR-FTIR studies and components analysis across the whole SC depth	98

i) Tape stripping procedure	98
ii) Extraction and analysis of each component in the tape strips	100
iii) Dermatopharmacokinetic comparisons	101
iv) Establishing correlations between FTIR spectrum changes and components concentrations	102
III- Confocal laser scanning microscope (CLSM) imaging	102
<b>Results and Discussion</b>	104
I- ATR-FTIR studies and spectral Analysis	104
II- FTIR changes across the whole SC depth	109
III- Correlations between FTIR changes and components concentrations across the whole SC depth	118
IV- Confocal laser scanning microscope imaging	127
<b>Conclusions</b>	131
<b>Chapter 4: Formulation and Characterization of Testosterone Transdermal Microemulsions</b>	134
<b>Item</b>	<b>Page</b>
<b>Introduction</b>	134
<b>Experimental</b>	140
Methodology	142
I- Preparation of testosterone microemulsions	142



II- Characterization of the prepared microemulsions	142
i) Electrical conductivity measurements	142
ii) Differential scanning calorimetry (DSC)	143
iii) Diffusion-ordered spectroscopy (DOSY)	143
iv) pH measurements	144
v) Rheological measurements	144
vi) Fourier transform infrared spectroscopy (FTIR)	144
vii) $^1\text{H}$ -NMR and $^{13}\text{C}$ -NMR	145
viii) Determination of the alteration in the microemulsion microstructure after drug loading	145
ix) Permeation studies	145

Item	Page
x) Study on the effect of increasing the drug loading on the microemulsion microstructure and the permeation flux	147
<b>Results and Discussion</b>	148
I- Electrical conductivity measurements	148
II- Differential scanning calorimetry (DSC)	153
III- Diffusion-ordered spectroscopy (DOSY)	154
IV- pH measurements	157

V- Rheological measurements	159
VI- Fourier transform infrared spectroscopy (FTIR)	161
VII- <sup>1</sup> H-NMR and <sup>13</sup> C-NMR	175
VIII- <i>Ex-vivo</i> permeation	180
IX- Effect of increasing the drug loading on the microemulsion microstructure	188
<b>Conclusions</b>	192
<b>Summary</b>	195
<b>References</b>	207
<b>Arabic Summary</b>	

### **List of Abbreviations**

Stratum Corneum	SC
Microemulsion	ME
Liquid Chromatography Mass Spectroscopy	LC / MS
Attenuated Total Reflectance Fourier Transform Infrared	ATR-FTIR
Confocal Laser Scanning Microscopy	CLSM
Total Ion Chromatogram	TIC
Single Ion Monitoring	SIM
Mono Ion Chromatogram	MIC
Mass to Charge ratio	m/z
Standard Deviation	S.D.
Limit of Detection	LOD
Limit of Quantitation	LOQ
Coefficient of Variation	C.V.
Retention Time	R <sub>t</sub>
High Performance Liquid Chromatography	HPLC
Internal Reflective Element	IRE
Transepidermal Water Loss	TEWL
Nile Red	NR
Dermatopharmacokinetic	DPK
Differential Scanning Calorimetry	DSC
Nuclear Magnetic Resonance	NMR
Diffusion-ordered Spectroscopy	DOSY

### List of Tables

<b>Table no.</b>	<b>Table Name</b>	<b>Page</b>
<b>1</b>	<b>Formulation considerations for passive transdermal delivery of drugs and their ideal limits.</b>	<b>9</b>
<b>2</b>	<b>Relationship between the concentrations of testosterone in ethyl alcohol (95%) and the peak areas using high performance liquid chromatography assay at 241 nm.</b>	<b>39</b>
<b>3</b>	<b>Effect of the Tween chain length on the water solubilisation parameters of the system oleic acid / Tween / Transcutol / water.</b>	<b>46</b>
<b>4</b>	<b>Dynamic Light Scattering Characterization for selected microemulsions from the system oleic acid / Tween20 / Transcutol / water.</b>	<b>54</b>
<b>5</b>	<b>LC / MS analysis of different concentrations of oleic acid at 210 nm and m/z = 281.</b>	<b>69</b>
<b>6</b>	<b>Validation parameters for the developed oleic acid quantitative method of analysis using LC / MS.</b>	<b>71</b>
<b>7</b>	<b>LC / MS analysis of different concentrations of Tween 20 at 254 nm and m/z = 1151.</b>	<b>73</b>
<b>8</b>	<b>Validation parameters for the developed Tween20 quantitative method of analysis using LC / MS.</b>	<b>75</b>
<b>9</b>	<b>LC / MS analysis of different concentrations of Transcutol at 254 nm and m/z = 135.</b>	<b>77</b>
<b>10</b>	<b>Validation parameters for the developed Transcutol quantitative method of analysis using LC / MS.</b>	<b>79</b>
<b>11</b>	<b>Penetration depth ranges (in <math>\mu\text{m}</math>) of infrared radiation between 4,000 and 625 <math>\text{cm}^{-1}</math> for a selection of substrate materials and angles of incidence (<math>n_1 = 1.6</math>).</b>	<b>85</b>
<b>12</b>	<b>Selected Microemulsion Formulations (% , w/w).</b>	<b>96</b>
<b>13</b>	<b>Partitioning (k) and Rate of Diffusion (<math>D/L^2</math>) of Different Microemulsion Components.</b>	<b>125</b>

<b>Table no.</b>	<b>Table Name</b>	<b>Page</b>
<b>14</b>	<b>Stratum Corneum uptake percentages of the</b>	<b>126</b>

	<b>microemulsion components following their application as pure chemicals or as constituents of microemulsion F.</b>	
<b>15</b>	<b>Conductivity measurements for the selected unloaded and loaded (1%, w/v) microemulsion formulations.</b>	<b>151</b>
<b>16</b>	<b>Diffusion and relative diffusion coefficients of the different microemulsion components obtained using DOSY-NMR.</b>	<b>156</b>
<b>17</b>	<b>pH measurements for the selected unloaded and unloaded (1%, w/v) microemulsion formulations.</b>	<b>158</b>
<b>18</b>	<b>Dynamic viscosity of the selected unloaded and loaded (1%, w/v) microemulsion formulations.</b>	<b>160</b>
<b>19</b>	<b>OH band frequencies of unloaded and loaded (1%, w/v) microemulsion formulations.</b>	<b>162</b>
<b>20</b>	<b><sup>1</sup>H-NMR chemical shifts of different functional groups of unloaded and loaded microemulsion F.</b>	<b>177</b>
<b>21</b>	<b><sup>13</sup>C-NMR chemical shifts of Oleic acid, Tween20 and Transcutol functional groups after incorporation of Drug (1%, w/v) in microemulsion F.</b>	<b>179</b>
<b>22</b>	<b>High performance liquid chromatography analysis of different concentrations of testosterone at 241 nm in PBS.</b>	<b>181</b>
<b>23</b>	<b>Drug Partitioning (KH) and Diffusivity (D/H<sup>2</sup>), Together with Estimated Permeability Coefficients (K<sub>p</sub>) and Steady-State Fluxes (J<sub>ss</sub>) of Testosterone (1%, w/v) from different microemulsions of the system oleic acid / Tween20 / Transcutol / water, across Porcine skin (740 µm).</b>	<b>186</b>
<b>24</b>	<b>Drug Partitioning (KH) and Diffusivity (D/H<sup>2</sup>) Parameters, Together with Estimated Permeability Coefficient (K<sub>p</sub>) and Steady-State Flux (J<sub>ss</sub>) of Testosterone (1% and 3%, w/v) from the microemulsion system oleic acid / Tween20 / Transcutol / water (16%, 32%, 32%, 20% respectively) across Porcine Skin (740 µm).</b>	<b>188</b>

<b>Table no.</b>	<b>Table Name</b>	<b>Page</b>
<b>25</b>	<b>Changes in the physical parameters of microemulsion E after loading with the drug (1% and 3%, w/v).</b>	<b>190</b>
<b>26</b>	<b><sup>1</sup>H-NMR chemical shifts of different functional groups of unloaded and loaded (3%, w/v) microemulsion E.</b>	<b>191</b>

### List of Figures

<b>Figure no.</b>	<b>Figure Name</b>	<b>Page</b>
<b>1</b>	<b>Stratum corneum morphology (brick and mortar-like pattern).</b>	<b>3</b>
<b>2</b>	<b>Macro-routes of drug penetration: (1) across the continuous stratum corneum; (2) through the hair follicles with their associated sebaceous glands or (3) via the sweat ducts.</b>	<b>3</b>
<b>3</b>	<b>Permeation routes through the stratum corneum (i) via the lipid matrix between the corneocytes (intercellular route) and (ii) across the corneocytes and the intercellular lipid matrix (transcellular route).</b>	<b>4</b>
<b>4</b>	<b>Structure of testosterone (17<math>\beta</math>-hydroxyandrost-4-en-3-one, M.W. 288)</b>	<b>17</b>
<b>5</b>	<b>Schematic representation of the three most commonly encountered microemulsion microstructures: (a) oil-in-water, (b) bicontinuous and (c) water-in-oil microemulsion.</b>	<b>27</b>
<b>6</b>	<b>Penetration of oil molecules between the hydrophobic chains of the interfacial surfactant monolayer in a bicontinuous microemulsion.</b>	<b>28</b>
<b>7</b>	<b>Chromatogram of testosterone at wavelength of 241 nm.</b>	<b>38</b>
<b>8</b>	<b>Testosterone calibration curve in ethyl alcohol (95 %) at 241 nm using high performance liquid chromatography.</b>	<b>40</b>
<b>9</b>	<b>Solubility of testosterone in different oils.</b>	<b>41</b>
<b>10</b>	<b>Solubilisation parameters for Oleic acid / Tween20 / Transcutol / Water phase diagram where the mixing ratio of Tween20 : Transcutol is 1:1. ME is the area of water-in-oil, bicontinuous and oil in water microemulsion (one phase region). W<sub>m</sub> is the maximum amount of solubilised water, S<sub>m</sub> is the amount of surfactant needed to obtain maximum water solubilisation.</b>	<b>44</b>
<b>11</b>	<b>Phase diagrams of different oleic acid / Tween /</b>	<b>45</b>