

CHEMISTRY AND BIOLOGY OF
PHENOLICS ISOLATED FROM
Ammannia auriculata
(LYTHRACEAE)

A Thesis Submitted By

Eman Sherien Mostafa

Master of Pharmaceutical Sciences,

Faculty of Pharmacy, Ain Shams University

In partial fulfillment of the
requirements for the degree of

Doctor of Philosophy in Pharmaceutical Sciences

(Pharmacognosy)

Department of Pharmacognosy

Faculty of Pharmacy

Ain Shams University

Abbasia, Cairo, Egypt

2016

Under the Supervision of

Nahla A. Ayoub, Ph.D.

Professor of Pharmacognosy
Dept. of Pharmacognosy
Faculty of Pharmacy
Ain Shams University

Mahmoud Abdel moein Nawwar, Ph.D.

Emeritus Professor of Phytochemistry
Dept. of Phytochemistry and Plant Systematic
National Research Center (N.R.C.)

Soumaya Saad Zaghloul, Ph.D.

Associate Professor of Pharmacognosy
Acting Head of Pharmacognosy Department
Department of Pharmacognosy
Faculty of Pharmacy
MSA University

Omayma A. Eldahshan, Ph.D.

Associate Professor of Pharmacognosy
Acting Head of Pharmacognosy Department
Department of Pharmacognosy
Faculty of Pharmacy
Ain Shams University

Mohamed Abdel aziz El Raey, Ph.D.

Researcher of Phytochemistry
Department of Phytochemistry and Plant Systematic
National Research Center (N.R.C.)

ACKNOWLEDGMENTS

*First of all, I would like to extend due praise and thanks to **Allah** for lightening my way.*

I would like to express my deepest gratitude, sincere thanks and profound appreciation to:

Prof. Dr. Nahla A. Ayoub, Professor of Pharmacognosy, Department of pharmacognosy, Ain-Shams University, (my beloved mum) for her Intellectual and academic support, her guidance, valuable discussions, variable knowledge, scientific advice and extensive efforts throughout this thesis and during the writing and revising stage. I am privileged to be her student not only as a postgraduate but also as an undergraduate at MIU where I still use her well written, simplified and valuable notes as a reference. Really I look up high at her as a model for successful woman in science

Prof. Dr. Mahmoud A. M. Nawwar, Professor of Phytochemistry, National Research Center, NRC, Cairo, (my teacher, my father, my professor) for suggesting the topic of this thesis, I would like to thank him for giving me the chance to work in his lab as an external student and for allowing me to use the lab facilities and for introducing me to international scientific groups through the GESP project where the various biological activities have been analyzed and for his extensive effort in allowing the data to be published to a reputable scientific journal. For his supervising, sponsoring, guidance, variable knowledge, valuable scientific advice and extensive efforts throughout this thesis. Also, for his valuable discussions and generous support specially the NMR interpretations. He has inspired me to become an independent researcher and helped me realize the power of critical reasoning. He also demonstrated what a brilliant and hardworking scientist can accomplish. Finally I would like to send a special thank to him for giving me the honor to be the youngest members of his respectable scientific school. Really, I am very proud to be his student and to get this wonderful chance to work with a scientist like him.

Assoc. Prof. Dr. Soumaya Saad Zaghloul, Associate Professor of Pharmacognosy and Acting Head of Pharmacognosy Department, Faculty of

pharmacy, MSA University, (my soul mum), for her encouragement, sound advice and kind help, especially during writing and revising the thesis. Really, I am proud to be her student and to be a member of her Pharmacognosy teamwork at MSA.

Assoc. Prof. Dr. Omayama A. Eldahshan, Associate Professor of Pharmacognosy and Acting Head of Pharmacognosy Department, Faculty of pharmacy, Ain Shams University, for her continuous support of my Ph.D. especially during the project and the comprehensive exam, for her patience, motivation and immense knowledge. Her guidance helped me in all the time of research, especially during writing and revising of this thesis.

Dr. Mohamed A. El-Raey, Doctor of Phytochemistry, Dept. Of Phytochemistry and Plant Systematic, National Research Center (N.R.C.), for his interest, his patience, his encouragement, his guidance and for his help and support especially during the practical work of the thesis.

My sincere thanks must also go to the members of my thesis advisory and exam committee: Prof. Dr. Meselhy Ragab Meselhy, Prof. of Pharmacognosy, faculty of Pharmacy, Cairo University and Prof. Dr. Helana Naguib Micheal, Professor of Natural Products, Department of Tanning Materials and Leather Technology, N.R.C. They generously gave their time to offer me valuable comments towards improving this work.

Prof. Dr. Ulrike Lindequist, Institut für Pharmazie, LS-Pharmaceutische Biologie, Ernst-Moritz-Arndt-Universität, Greifswald, Germany, for hosting my short stay in Greifswald, allowing me to use the lab facilities and her valuable guidance in the biological assays which were performed at her institute.

Dr. Michael Linscheid, Professor of Analytical and Environmental Chemistry, Institute für Chemie, Humboldt Universität zu Berlin, Germany, for his constructive cooperation during the mass spectroscopic analysis.

PD Dr. Beate Haertel Institut für Pharmazie, Lehrstuhl für Pharmazeutische Biologie, Ernst-Moritz-Arndt Universität, Greifswald, Germany, for her essential assistance, support and advices in the evaluation of the biological activities.

Dr. Susanne Strassenburg, Institut für Pharmazie, Lehrstuhl für Pharmazeutische Biologie, Ernst-Moritz-Arndt Universität, Greifswald, Germany, for her endless help during my short stay in Greifswald, Germany, her essential assistance and valuable guidance in the cell assays.

The thesis was supported and financed by the Alexander von Humboldt foundation through the group linkage program (joint project: "Bioactive phenolics from Egyptian folk medicinal plants", 3.4-Fokoop-DEU/1093980) awarded to U. Lindequist and M. Nawwar.

Dr. Amani N. M. Hashem Associate Professor of Phytochemistry, National Research Center, N.R.C, for her encouragement, sound advice and kind help, during the thesis.

Prof. Dr. Salwa Kawashti, Dr. Mona Marzouk and Dr. Sameh Hussien, Department of Phytochemistry and Plant Systematic, National Research Center, for their efforts in the identification of the plant material.

I am deeply thankful to Dr. Nawal El Degwi Head of Board of Trustees, MSA University for giving me the chance to be a one of her university staff member and providing the time for me to finish my thesis practical work. Also, for her, encouragement and support.

I am deeply thankful to all Ain Shams University staff members who helped me a lot during the post graduate courses, for their cooperation, encouragement and support.

I am deeply thankful to all my colleagues at the NRC, MSA and Ain Shams University, for their help, encouragement and support.

With all the love, I thank my husband, Mohamed. He has been my best friend and great companion, loved, supported, encouraged, entertained and helped me get through this agonizing period in the most positive way.

Most importantly, none of this would have been possible without the love and patience of my dear MUM, to whom this work is dedicated. She has been a constant source of love, concern, support and strength throughout my whole life. I deeply thank her for her unconditional love, support, encouragement and never ending tolerance and patience.

Finally, I would like to express my deepest gratitude and endless love to brother Ahmed and my sisters Hala, Mai, Eman, and Sahar for their unconditional love, support and encouragement and a special point out to my little sun Shine Julie and Kenzy for just being there with their innocent irreplaceable pure smile and to whom I dedicate this work.

CONTENTS

	Page
1. INTRODUCTION	1
2. REVIEW OF LITERATURE	6
3. TAXONOMY	33
4. MATERIAL, APPARATUS AND METHODS	40
5. EXPERIMENTAL	
Phenolic constituents of the flowering whole plant of <i>Ammannia auriculata</i>	
A- Phytochemical Screening of the flowering whole plant of <i>Ammannia auriculata</i>..	56
B- Extraction	
B-1 Qualitative phenolic analysis of <i>Ammannia auriculata</i> extract.....	56
B-2 Chromatographic investigation of <i>Ammannia auriculata</i> extract	57
B-2-1 Paper chromatographic investigation of <i>Ammannia auriculata</i> extract.....	57
B-2-2 Column chromatographic investigation of <i>Ammannia auriculata</i> extract.....	58
C- Isolation of compounds (1-20) from the column fractions (I- X)	60
Paper chromatographic analysis of fractions (I-X).....	60
Fraction I	60
Fraction II	61
Isolation of compounds 1 and 2	61
Identification of compound 1 , 2,3- <i>O</i> -hexahydroxydiphenoyl-(α/β)-glucose	61
Identification of compound 2 , Gallic acid.....	67
Fraction III.....	71
Fraction IV.....	71
Isolation of compounds 3-6	71
Identification of compound 3 , 2,3-di- <i>O</i> -galloyl-(α/β)- ⁴ C ₁ -glucopyranose	72
Identification of compound 4 , 1,6-di- <i>O</i> -galloyl- β - ⁴ C ₁ -glucopyranose	78
Identification of compound 5 , Corilagin	83
Identification of compound 6 , Myricetin-3- <i>O</i> - β - ⁴ C ₁ -glucopyranoside	103
Fraction V.....	111
Isolation of compound 7	111
Identification of compound 7 , Kaempferol-3- <i>O</i> -(6''- <i>p</i> -coumaroyl- β - glucopyranoside)..<	111
Fraction VI.....	121

Isolation of compounds 8 and 9	121
Identification of compound 8 , 1- <i>O</i> -galloyl-2,3- <i>O</i> -Hexahydroxydiphenoyl- ⁴ C ₁ - α -glucopyranose.....	122
Identification of compound 9 , Quercetin-3- <i>O</i> - β -glucopyranoside.....	132
Fraction VII	137
Isolation of compounds 10-13	137
Identification of compound 10 , Quercetin-3- <i>O</i> - β -(6''- <i>O</i> -galloylglucopyranoside).....	138
Identification of compound 11 , Kaempferol-3- <i>O</i> - β -glucopyranoside.....	146
Identification of compound 12 , 6- <i>O</i> -monogalloyl-2,3- <i>O</i> -hexahydroxydiphenoyl-(α/β)- ⁴ C ₁ -glucopyranose.....	151
Identification of compound 13 , Tellimagrandin I.....	162
Fraction VIII	170
Isolation of compounds 14 and 15	170
Identification of compound 14 , 1,2,3-tri- <i>O</i> -galloyl- α - ⁴ C ₁ -glucose.....	170
Identification of compound 15 , 1,2,3-tri- <i>O</i> -galloyl- β - ⁴ C ₁ -glucose.....	176
Fraction IX	181
Isolation of compounds 16-19	181
Identification of compound 16 , 1,4,6-tri- <i>O</i> -galloyl- β - ⁴ C ₁ -glucopyranose.....	181
Identification of compound 17 , New natural product, 2,3,6-tri- <i>O</i> -galloyl-(α/β)- ⁴ C ₁ -glucopyranose.....	191
Identification of compound 18 , New natural product, Kaempferol-3- <i>O</i> - β -(6''-galloylglucopyranoside)-7- <i>O</i> - β -glucopyranoside.....	208
Identification of compound 19 , New natural product, Quercetin-3- <i>O</i> - β -(6''-galloylglucopyranoside)-7- <i>O</i> - β -glucopyranoside.....	227
Fraction X	235
Isolation of compound 20	235
Identification of compound 20 , Gallic acid- 3- <i>O</i> - β - ⁴ C ₁ -glucopyranoside.....	235
C- Biological assay of the extract	243
C-1 Assay for antioxidant activity.....	243
C-2 Assay for cell culture.....	253
C-3 NRU-assay (Cytotoxicity assay).....	255
C-4 Comparison between the antioxidant and cytotoxic activities.....	265

C-5	Cell cycle analysis.....	267
C-6	Discussion.....	276
6.	SUMMARY.....	281
7.	ACHIEVEMENTS AND RECOMMENDATIONS.....	293
8.	REFERENCES	297
9.	ARABIC SUMMARY	

List of Schemes

Scheme (1)	Scheme of work of <i>Ammannia auriculata</i> (Lytheraceae).....	289
------------	---	-----

List of Figures

FIGURE		PAGE
1.	Structure of compounds characterized by GC-MS analysis of hexane extract and VLC fractions of chloroform extract of <i>Ammannia multiflora</i> .	21
2.	<i>Ammannia auriculata</i> herb.	37
3.	<i>Ammannia auriculata</i> flowers.	38
4.	<i>Ammannia auriculata</i> fruits.	39
5.	ESI-MS spectrum of compound (1); 2,3- <i>O</i> -hexahydroxydiphenoyl-(α/β)-glucose.	64
6.	^1H -NMR spectrum of compound (1); 2,3- <i>O</i> -hexahydroxydiphenoyl-(α/β)-glucose.	65
7.	^{13}C -NMR spectrum of compound (1); 2,3- <i>O</i> -hexahydroxydiphenoyl-(α/β)-glucose.	66
8.	ESI-MS spectrum of compound (2); Gallic acid.	68
9.	^1H -NMR spectrum of compound (2); Gallic acid.	69
10.	^{13}C -NMR spectrum of compound (2); Gallic acid.	70
11.	Negative ESI-MS spectrum of compound (3); 2,3-di- <i>O</i> -galloyl-(α/β)- $^4\text{C}_1$ -glucopyranose.	74
12.	Negative ESI-MS spectrum of compound (3a); monogalloyl glucose.	75
13.	^1H -NMR spectrum of compound (3); 2,3-di- <i>O</i> -galloyl-(α/β)- $^4\text{C}_1$ -glucopyranose.	76
14.	^{13}C -NMR spectrum of compound (3); 2,3-di- <i>O</i> -galloyl-(α/β)- $^4\text{C}_1$ -glucopyranose.	77
15.	ESI-MS spectrum of compound (4); 1,6-di- <i>O</i> -galloyl- β - $^4\text{C}_1$ -glucopyranose.	80
16.	^1H -NMR spectrum of compound (4); 1,6-di- <i>O</i> -galloyl- β - $^4\text{C}_1$ -glucopyranose.	81
17.	^{13}C -NMR spectrum of compound (4); 1,6-di- <i>O</i> -galloyl- β - $^4\text{C}_1$ -glucopyranose.	82
18.	Positive FAB-MS spectrum of compound (5); Corilagin.	89
19.	Negative FAB-MS spectrum of compound (5); Corilagin.	90
20.	Positive FAB-MS spectrum of compound (5a); 3,6-hexahydroxydiphenoyl-(α/β)- $^1\text{C}_4$ -glucopyranose.	91
21.	Negative FAB-MS spectrum of compound (5a); 3,6-hexahydroxydiphenoyl-(α/β)- $^1\text{C}_4$ -glucopyranose.	92
22.	^1H -NMR spectrum of ellagic acid.	93
23.	^{13}C -NMR spectrum of ellagic acid.	94
24.	^1H -NMR spectrum of compound (5); Corilagin.	95
25.	^{13}C -NMR spectrum of compound (5); Corilagin.	96
26.	COSY spectrum of compound (5); Corilagin.	97

27.	¹ H-NMR spectrum of compound (5a); 3,6-hexahydroxydiphenoyl-(α/β)- ¹ C ₄ -glucopyranose.	98
28.	¹³ C-NMR spectrum of compound (5a); 3,6-hexahydroxydiphenoyl-(α/β)- ¹ C ₄ -glucopyranose.	99
29.	DEPT 135 spectrum of compound (5a); 3,6-hexahydroxydiphenoyl-(α/β)- ¹ C ₄ -glucopyranose.	100
30.	COSY spectrum of compound (5a); 3,6-hexahydroxydiphenoyl-(α/β)- ¹ C ₄ -glucopyranose.	101
31.	HETCOR spectrum of compound (5a); 3,6-hexahydroxydiphenoyl-(α/β)- ¹ C ₄ -glucopyranose.	102
32.	Negative ESI-MS spectrum of compound (6); Myricetin-3- <i>O</i> - β - ⁴ C ₁ -glucopyranoside.	106
33.	Negative ESI-MS spectrum of compound (6a); Myricetin.	107
34.	¹ H-NMR spectrum of compound (6a); Myricetin.	108
35.	¹ H-NMR spectrum of compound (6); Myricetin-3- <i>O</i> - β - ⁴ C ₁ -glucopyranoside.	109
36.	¹³ C-NMR spectrum of compound (6); Myricetin-3- <i>O</i> - β - ⁴ C ₁ -glucopyranoside.	110
37.	Negative ESI-MS spectrum of compound (7); Kaempferol-3- <i>O</i> -(6''- <i>p</i> -coumaroyl- β -glucopyranoside).	115
38.	¹ H-NMR spectrum of compound (7); Kaempferol-3- <i>O</i> -(6''- <i>p</i> -coumaroyl- β -glucopyranoside).	116
39.	¹³ C-NMR spectrum of compound (7); Kaempferol-3- <i>O</i> -(6''- <i>p</i> -coumaroyl- β -glucopyranoside).	117
40.	¹ H-NMR spectrum of kaempferol.	118
41.	¹ H-NMR spectrum of <i>p</i> -coumaric acid.	119
42.	¹³ C-NMR spectrum of <i>p</i> -coumaric acid.	120
43.	Negative ESI-MS spectrum of compound (8); 1- <i>O</i> -monogalloyl-2,3- <i>O</i> -hexahydroxydiphenoyl- α - ⁴ C ₁ -glucopyranose.	127
44.	Negative ESI-MS spectrum of compound (8a); Ellagic acid.	128
45.	¹ H-NMR spectrum of compound (8); 1- <i>O</i> -monogalloyl-2,3- <i>O</i> -hexahydroxydiphenoyl- α - ⁴ C ₁ -glucopyranose.	129
46.	COSY spectrum of compound (8); 1- <i>O</i> -monogalloyl-2,3- <i>O</i> -hexahydroxydiphenoyl- α - ⁴ C ₁ -glucopyranose.	130
47.	¹³ C-NMR spectrum of compound (8); 1- <i>O</i> -monogalloyl-2,3- <i>O</i> -hexahydroxydiphenoyl- α - ⁴ C ₁ -glucopyranose.	113
48.	Negative ESI-MS spectrum of compound (9); Quercetin-3- <i>O</i> - β -glucopyranoside.	134
49.	¹ H-NMR spectrum of compound (9); Quercetin-3- <i>O</i> - β -glucopyranoside.	135
50.	¹³ C-NMR spectrum of compound (9); Quercetin-3- <i>O</i> - β -	136

	glucopyranoside.	
51.	Negative ESI-MS spectrum of compound (10); Quercetin-3- <i>O</i> - β -(6''- <i>O</i> -galloylglucopyranoside).	142
52.	¹ H-NMR spectrum of quercetin.	143
53.	¹ H-NMR spectrum of compound (10); Quercetin-3- <i>O</i> - β -(6''- <i>O</i> -galloylglucopyranoside).	144
54.	¹³ C-NMR spectrum of compound (10); Quercetin-3- <i>O</i> - β -(6''- <i>O</i> -galloylglucopyranoside).	145
55.	Positive ESI-MS spectrum of compound (11); Kaempferol-3- <i>O</i> - β -glucopyranoside.	148
56.	¹ H-NMR spectrum of compound (11); Kaempferol-3- <i>O</i> - β -glucopyranoside.	149
57.	¹³ C-NMR spectrum of compound (11); Kaempferol-3- <i>O</i> - β -glucopyranoside.	150
58.	Negative ESI-MS spectrum of compound (12); 6- <i>O</i> -monogalloyl-2,3- <i>O</i> -hexahydroxydiphenoyl-(α/β)- ⁴ C ₁ -glucopyranose.	157
59.	ESI-MS spectrum of compound (12c); 6-mono- <i>O</i> -galloyl- ⁴ C ₁ -D-glucopyranose.	158
60.	¹ H-NMR spectrum of compound (12c); 6-mono- <i>O</i> -galloyl- ⁴ C ₁ -D-glucopyranose.	159
61.	¹ H-NMR spectrum of compound (12); 6- <i>O</i> -monogalloyl-2,3- <i>O</i> -hexahydroxydiphenoyl-(α/β)- ⁴ C ₁ -glucopyranose.	160
62.	¹³ C-NMR spectrum of compound (12); 6- <i>O</i> -monogalloyl-2,3- <i>O</i> -hexahydroxydiphenoyl-(α/β)- ⁴ C ₁ -glucopyranose.	161
63.	Negative ESI-MS spectrum of compound (13); Tellimagrandin I.	167
64.	¹ H-NMR spectrum of compound (13); Tellimagrandin I.	168
65.	¹³ C-NMR spectrum of compound (13); Tellimagrandin I.	169
66.	Negative FAB-MS spectrum of compound (14); 1, 2, 3-tri- <i>O</i> -galloyl- α - ⁴ C ₁ -glucose.	173
67.	¹ H-NMR spectrum of compound (14); 1,2,3-tri- <i>O</i> -galloyl- α - ⁴ C ₁ -glucose.	174
68.	¹³ C-NMR spectrum of compound (14); 1,2,3-tri- <i>O</i> -galloyl- α - ⁴ C ₁ -glucose.	175
69.	Negative FAB-MS spectrum of compound (15); 1,2,3-tri- <i>O</i> -galloyl- β - ⁴ C ₁ -glucose.	178
70.	¹ H-NMR spectrum of compound (15); 1,2,3-tri- <i>O</i> -galloyl- β - ⁴ C ₁ -glucose.	179
71.	¹³ C-NMR spectrum of compound (15); 1,2,3-tri- <i>O</i> -galloyl- β - ⁴ C ₁ -glucose.	180
72.	Negative FAB-MS spectrum of compound (16); 1, 4, 6-tri- <i>O</i> -galloyl- β - ⁴ C ₁ -glucopyranose.	185

73.	¹ H-NMR spectrum of compound (16); 1,4,6-tri- <i>O</i> -galloyl- β - ⁴ C ₁ -glucopyranose.	186
74.	COSY spectrum of compound (16); 1,4,6-tri- <i>O</i> -galloyl- β - ⁴ C ₁ -glucopyranose.	187
75.	¹³ C-NMR spectrum of compound (16); 1,4,6-tri- <i>O</i> -galloyl- β - ⁴ C ₁ -glucopyranose.	188
76.	HMBC spectrum of compound (16); 1,4,6-tri- <i>O</i> -galloyl- β - ⁴ C ₁ -glucopyranose.	189
77.	HMQC spectrum of compound (16); 1,4,6-tri- <i>O</i> -galloyl- β - ⁴ C ₁ -glucopyranose.	190
78.	Negative ESI-MS spectrum of compound (17); 2,3,6-tri- <i>O</i> -galloyl-(α/β)- ⁴ C ₁ -glucopyranose.	196
79.	HRESI-MS spectrum of compound (17); 2,3,6-tri- <i>O</i> -galloyl-(α/β)- ⁴ C ₁ -glucopyranose.	197
80.	Negative ESI-MS spectrum of compound (17a); 3,6-tri- <i>O</i> -galloyl-(α/β)- ⁴ C ₁ -glucopyranose.	198
81.	HRESI-MS spectrum of compound (17a); 3,6-tri- <i>O</i> -galloyl-(α/β)- ⁴ C ₁ -glucopyranose.	199
82.	¹ H-NMR spectrum of compound (17a); 3,6-tri- <i>O</i> -galloyl-(α/β)- ⁴ C ₁ -glucopyranose.	200
83.	¹³ C-NMR spectrum of compound (17a); 3,6-tri- <i>O</i> -galloyl-(α/β)- ⁴ C ₁ -glucopyranose.	201
84.	¹ H-NMR spectrum of compound (17); 2,3,6-tri- <i>O</i> -galloyl-(α/β)- ⁴ C ₁ -glucopyranose.	202
85.	¹ H-NMR spectrum of compound (17)''; 2,3,6-tri- <i>O</i> -galloyl-(α/β)- ⁴ C ₁ -glucopyranose.	203
86.	¹³ C-NMR spectrum of compound (17); 2,3,6-tri- <i>O</i> -galloyl-(α/β)- ⁴ C ₁ -glucopyranose.	204
87.	COSY spectrum of compound (17); 2,3,6-tri- <i>O</i> -galloyl-(α/β)- ⁴ C ₁ -glucopyranose.	205
88.	HMBC spectrum of compound (17); 2,3,6-tri- <i>O</i> -galloyl-(α/β)- ⁴ C ₁ -glucopyranose.	206
89.	HMQC spectrum of compound (17); 2,3,6-tri- <i>O</i> -galloyl-(α/β)- ⁴ C ₁ -glucopyranose.	207
90.	Negative ESI-MS spectrum of compound (18a); Kaempferol-3- <i>O</i> - β -(6''- galloylglucopyranoside).	214
91.	¹ H-NMR spectrum of (18a); Kaempferol-3- <i>O</i> - β -(6''- galloylglucopyranoside).	215
92.	¹³ C-NMR spectrum of compound (18a); Kaempferol-3- <i>O</i> - β -(6''- galloylglucopyranoside).	216
93.	¹³ C-NMR spectrum of compound (18a)''; Kaempferol-3- <i>O</i> - β -(6''- galloylglucopyranoside).	217
94.	Negative ESI-MS spectrum of compound (18); Kaempferol 3- <i>O</i> - β -(6''-galloylglucopyranoside)-7- <i>O</i> - β -glucopyranoside.	218
95.	MS-MS spectrum of compound (18); Kaempferol-3- <i>O</i> - β -(6''-galloylglucopyranoside)-7- <i>O</i> - β -glucopyranoside.	219
96.	HRESI-MS spectrum of compound (18); Kaempferol-3- <i>O</i> - β -(6''-galloylglucopyranoside)-7- <i>O</i> - β -	220

	glucopyranoside.	
97.	¹ H-NMR spectrum of compound (18); Kaempferol-3- <i>O</i> - β -(6''-galloylglucopyranoside)-7- <i>O</i> - β -glucopyranoside.	221
98.	¹³ C-NMR spectrum of compound (18); Kaempferol-3- <i>O</i> - β -(6''-galloylglucopyranoside)-7- <i>O</i> - β -glucopyranoside.	222
99.	TCOSY spectrum of compound (18); Kaempferol-3- <i>O</i> - β -(6''-galloylglucopyranoside)-7- <i>O</i> - β -glucopyranoside.	223
100.	COSY spectrum of compound (18); Kaempferol-3- <i>O</i> - β -(6''-galloylglucopyranoside)-7- <i>O</i> - β -glucopyranoside.	224
101.	HSQC spectrum of compound (18); Kaempferol-3- <i>O</i> - β -(6''-galloylglucopyranoside)-7- <i>O</i> - β -glucopyranoside.	225
102.	HMBC spectrum of compound (18); Kaempferol-3- <i>O</i> - β -(6''-galloylglucopyranoside)-7- <i>O</i> - β -glucopyranoside.	226
103.	Negative ESI-MS spectrum of compound (19); Quercetin-3- <i>O</i> - β -(6''-galloylglucopyranoside)-7- <i>O</i> - β -glucopyranoside.	230
104.	MS-MS spectrum of compound (19); Quercetin-3- <i>O</i> - β -(6''-galloylglucopyranoside)-7- <i>O</i> - β -glucopyranoside.	231
105.	HRESI-MS spectrum of compound (19); Quercetin-3- <i>O</i> - β -(6''-galloylglucopyranoside)-7- <i>O</i> - β -glucopyranoside.	232
106.	¹ H-NMR spectrum of compound (19); Quercetin-3- <i>O</i> - β -(6''-galloylglucopyranoside)-7- <i>O</i> - β -glucopyranoside.	233
107.	¹³ C-NMR spectrum of compound (19); Quercetin-3- <i>O</i> - β -(6''-galloylglucopyranoside)-7- <i>O</i> - β -glucopyranoside.	234
108.	Positive FAB-MS spectrum of compound (20); Gallic acid-3- <i>O</i> - β - ⁴ C ₁ -glucopyranoside.	238
109.	¹ H-NMR spectrum of compound (20); Gallic acid-3- <i>O</i> - β - ⁴ C ₁ -glucopyranoside.	239
110.	HSQC spectrum of compound (20); Gallic acid-3- <i>O</i> - β - ⁴ C ₁ -glucopyranoside.	240
111.	¹³ C-NMR spectrum of compound (20); Gallic acid-3- <i>O</i> - β - ⁴ C ₁ -glucopyranoside.	241
112.	HMBC spectrum of compound (20); Gallic acid-3- <i>O</i> - β - ⁴ C ₁ -glucopyranoside.	242
113.	Mechanisms of antioxidant activity.	244
114.	Comparison between the % inhibition of DPPH absorbance for ascorbic acid and <i>Ammannia auriculata</i> extract at 7 different concentrations.	246
115.	Comparison between the radical scavenging activity of ascorbic acid and <i>Ammannia auriculata</i> extract.	246
116.	Principle of the ORAC assay.	248
117.	Radical scavenging activity of <i>Ammannia auriculata</i> extract and its individual fractions compared to trolox.	249
118.	Comparison between the % inhibition of fluorescence decay for <i>Ammannia auriculata</i> extract and the ten different fractions at 8 different concentrations.	250